

Individual patterns of activity predict the response to physical exercise as an intervention in mild to moderate depression

Stefan Spulber¹, Sandra Ceccatelli¹, Yvonne Forsell²

¹ – Department of Neuroscience; ² – Department of Global Public Health, Karolinska Institutet, 171 77 Stockholm, Sweden

Supplementary Material

Calculation of prior inclusion probability for a specific feature X_i


$$\Pr(X_i) = \frac{N \text{ models including } X_i}{N \text{ models possible}} \quad (1)$$

For a given complexity level, the total number of models possible to train is equal to the binomial coefficient $\binom{n}{k}$, and the number of models including a specific feature is the binomial coefficient $\binom{n-1}{k-1}$, where n = total number of features available, and k = model complexity (number of independent variables in the model). Therefore equation (1) can be rewritten as follows:

$$\Pr(X_i) = \frac{\sum_{k=0}^5 \binom{n-1}{k}}{\sum_{k=0}^6 \binom{n}{k}} \quad (2)$$


Solving (2) numerically for $n=26$ yields $\Pr(X_i) = \frac{68\,401}{313\,912} = 0.2179$. The prior inclusion probability of 0.2179 applies to all features, because we trained models for all possible combinations of up to 6 features/model. After pruning the model ensembles, enriched features must have $\text{PIP} > 0.2179$, while depleted features have $\text{PIP} < 0.2179$.

Supplementary Fig. S1 Feature set selection




Correlations between main set of features calculated as daily average (meand), or estimated on full-length recording or average circadian profile (meanp), sequential differences in daily values (sdiff), and variance of daily values (*i.e.*, sequence is ignored; mdiff). The lowest correlation coefficients are observed between meanp and mdiff (arrowheads), suggesting that the information carried by the two feature sets is complementary (non-overlapping).

Supplementary Fig. S2 Feature selection




Top panel: correlations between the two complete feature sets surviving FDR correction. The clusters of highly correlated and conceptually related features were pruned to include preferentially the features with least correlations in the cluster. Bottom panels: correlations between features after feature selection and differences across groups and genders for selected features.

Supplementary Fig. S3 Trained ensembles




Supplementary Fig. S4 Comparison between predicted outcomes for all treatments and subjects.



Dashed lines indicate regression slopes estimated using the Theil-Sen method (for illustration purposes only). Note the predicted response to moderate intensity PE is independent for the predicted response to low or high intensity PE, while the response to low and high intensity PE are inversely correlated.

Supplementary Fig. S5 Predicted improvement in treatment outcome provided each patient is assigned to the PE regime predicted to yield the best response



(A) Evaluation of potential selection bias and estimated improvement. The remission rate was slightly higher in the subset of patients used for model training as compared to the global population assigned to PE as antidepressive intervention. The effect size was estimated as $(\text{best} - \text{random}) / \text{random}$ and was applied on the complete dataset in (B). (B) Illustration of improvement in treatment outcome provided each patient is assigned to the PE regime predicted to yield best response. The relative increase in remission rate between PE* and PE is the effect size calculated in (A). Note the net departure from remission rate for TAU as compared to observed remission rate following random allocation to treatment arms in the original study.