1 Title

- 2 Cardiovascular risk of dementia is associated with brain–behaviour changes in cognitively
- 3 healthy, middle-aged individuals

4 Authors

- 5 Feng Deng^{ab}, Maria-Eleni Dounavi^c, Emanuele R. G. Plini^a, Karen Ritchie^d, Graciela
- 6 Muniz-Terrera ^{ef}, Siobhan Hutchinson ^g, Paresh Malhotra ^h, Craig W Ritchie ^e, Brian Lawlor

7 ^{ab}, Lorina Naci ^{ab*}

8 Affiliations

- 9 ^a Trinity College Institute of Neuroscience, School of Psychology, Trinity College Dublin,
- 10 Dublin, Ireland.

¹¹ ^b Global Brain Health Institute, Trinity College Dublin, Dublin, Ireland

- ^c Department of Psychiatry, School of Clinical Medicine, University of Cambridge,
- 13 Cambridge CB2 0SP, UK
- ¹⁴ ^d U1061 Neuropsychiatry, INSERM, University of Montpellier, Montpellier, France
- ^e Edinburgh Dementia Prevention, University of Edinburgh, Edinburgh, UK
- ¹⁶ ^fDepartment of Social medicine, Ohio University, USA
- ^g Department of Neurology, St James's Hospital, Dublin, Ireland
- ¹⁸ ^h Department of Brain Science, Imperial College Healthcare NHS Trust, UK
- 19 * Corresponding author: NOTE: This preprint reports new research that has not been certified by peer review and should not be used to guide clinical practice.

- 20 Lorina Naci
- 21 School of Psychology
- 22 Trinity College Institute of Neuroscience
- 23 Global Brain Health Institute
- 24 Trinity College Dublin
- 25 Dublin, Ireland
- 26 Email: <u>nacil@tcd.ie</u>

27 Abstract

28	Alzheimer's Disease (AD) neuropathology start decades before clinical manifestations, but
29	whether risk factors are associated with early cognitive and brain changes in midlife remains
30	poorly understood. We examined whether AD risk factors were associated with cognition and
31	functional connectivity (FC) between the Locus Coeruleus (LC) and hippocampus – two key
32	brain structures in AD neuropathology – cross-sectionally and longitudinally in cognitively
33	healthy midlife individuals. Neuropsychological assessments and functional Magnetic
34	Resonance Imaging were obtained at baseline (N=210), and two-years follow-up (N=188).
35	Associations of cognition and FC with apolipoprotein ɛ4 (APOE ɛ4) genotype, family history
36	of dementia, and the Cardiovascular Risk Factors, Aging, and Incidence of Dementia
37	(CAIDE) score were investigated. Cross-sectionally, higher CAIDE scores were associated
38	with worse cognition. Menopausal status interacted with the CAIDE risk on cognition.
39	Furthermore, the CAIDE score significantly moderated the relationship between cognition
40	and LC-Hippocampus FC. Longitudinally, the LC-Hippocampus FC decreased significantly
41	over 2 years. These results shed light on some of the earliest effects of AD risk on cognition
42	and the brain.

- 44 Keywords: Alzheimer's disease; Cognition; Locus Coeruleus; Hippocampus; Functional
- 45 connectivity; Midlife; Cardiovascular risk score

46 1. Introduction

47	Dementia, particularly Alzheimer's disease (AD), is a growing epidemic that presents
48	profound challenges to healthcare systems, families, and societies throughout the world
49	(World Health Organization, 2021). Midlife is a critical period for the beginning of AD
50	pathology (Jansen et al., 2015; Sperling et al., 2011) and potentially a unique disease-altering
51	window prior to the manifestation of substantial brain damage. Therefore, there is an urgent
52	need for risk reduction interventions focused on midlife (Barnes & Yaffe, 2011; Livingston et
53	al., 2017; Ritchie et al., 2010). However, the indicators and brain mechanism of AD in
54	midlife remain poorly understood (Irwin et al., 2018; Ritchie et al., 2017).
<i></i>	To investigate early changes, studies in preclinical AD nonvestions have used risk
55	To investigate early changes, studies in preclinical AD populations have used risk
56	stratification approaches. Key risk factors include the Apolipoprotein E (APOE) ɛ4 genotype
57	(Donix et al., 2012; Liu et al., 2013), the main genetic risk factor for sporadic late-onset AD
58	in the Indo-European population (Lambert et al., 2013), and family history of dementia (FHD)
59	(Berti et al., 2011; Donix et al., 2012; Scarabino et al., 2016). Additionally, there is growing
60	evidence to suggest that up to 40% of all dementia cases are associated with known
61	modifiable risk factors (Livingston et al., 2020). Among dementia risk scores that
62	incorporate lifestyle risk factors (Barnes et al., 2014; Deckers et al., 2015; Kivipelto et al.,
63	2006), the Cardiovascular Risk Factors Aging and Dementia (CAIDE) score has been
64	optimized for midlife (Kivipelto et al., 2006) and validated in a large US population followed
65	longitudinally over 40 years (Exalto et al., 2014).
66	AD is neuropathologically characterized by the accumulation of beta-amyloid (A β) and
67	hyperphosphorylated tau (pTau) (Braak & Braak, 1991; Braak et al., 2011). Recent studies
68	suggest that tau deposition is a key etiological factor that presages sporadic AD (Arnsten et

69 al., 2021; Jacobs et al., 2021; Kametani & Hasegawa, 2018). Analyses of thousands of human

70 brains across the lifespan suggest that tau pathology may begin about a decade before 71 formation of A β plaques (Braak & Del Tredici, 2015). Additionally, tau pathology correlates with progressive grey matter loss to a greater extent than A β (La Joie et al., 2019) and 72 73 cognitive impairment (Giannakopoulos et al., 2003). Neuropathological findings show that the nucleus locus coeruleus (LC) is one of the subcortical regions where tau pathology starts 74 in late young adulthood and early midlife (30–40 years) (Gilvesy et al., 2022; Ehrenberg et al., 75 76 2017; Braak & Braak, 1991) then spreads to the transentorhinal and entorhinal cortex (Braak et al., 2011), to the hippocampal and neocortical association cortex, and finally throughout 77 78 the neocortex.

It has been suggested that early tau pathology in the LC induces LC hyperactivity, thereby 79 promoting its own spread to interconnected brain regions and facilitating the progression of 80 the disease (Weinshenker, 2018). A recent review of the neuropathological literature across 81 the lifespan (Arnsten et al., 2021) suggests that the spread of tau pathology from the LC to 82 83 the medial temporal lobe (MTL) captures the earliest stages of incipient AD progression (but 84 see Kaufman et al. (2018) for a differing account). However, alterations of connectivity between the LC and the hippocampus, due to AD risk factors have not previously been 85 86 studied in midlife preclinical populations.

LC is the earliest site of tau pathology accumulation, and, furthermore, the main brain site of
noradrenaline production (Amaral & Sinnamon 1977), and thus the originating node of the
brain's arousal system (Sara, 2009; Aston-Jones & Cohen 2005). Therefore, early LC
pathology can critically impact arousal and cognitive processes throughout the brain (Mather
& Harley, 2016; Betts, Kirilina, et al., 2019). Previous in vivo studies of the LC have
investigated healthy lifespan adult cohorts (Betts et al., 2017; Clewett et al., 2016; Lee et al.,
2020; Liu et al., 2019; Porat et al., 2022), cognitively unimpaired older adults and preclinical

94 populations with risk for sporadic AD (Jacobs et al., 2021; Prokopiou et al., 2022; Van Egroo et al., 2021), individuals with preclinical autosomal dominant Alzheimer's disease (Jacobs et 95 al., 2022), mild cognitive impairment (Jacobs et al., 2015; Takahashi et al., 2015), and 96 97 Alzheimer's disease (Betts, Cardenas-Blanco, et al., 2019; Dahl et al., 2022; Olivieri et al., 2019; Takahashi et al., 2015) and found significant negative associations of LC integrity with 98 age, age-related cognitive decline (Calarco et al., 2022; Dahl et al., 2019; Liu et al., 2020), 99 AD pathology, such as amyloid (Betts, Cardenas-Blanco, et al., 2019) and tau deposition 100 101 (Dahl et al., 2022; Jacobs et al., 2022), as well as with symptom severity in AD (Sun et al., 102 2020). Prokopiu et al. (2022) also found that lower novelty-related LC activity and LC to hippocampus and parahippocampus functional connectivity were associated with steeper 103 104 beta-amyloid-related cognitive decline. However, by contrast to its key importance to 105 cognitive function and etiological role in AD progression, there is very limited in vivo understanding of LC function disruptions during the early preclinical period. One exception 106 is a recent study of asymptomatic midlife adults, offspring of late-onset AD (O-LOAD) 107 108 patients. Del Cerro and colleagues (2020) found that the connectivity of the LC and 109 cerebellum was significantly associated with delayed recall performance, and reduced in O-LOAD individuals relative to those with no family history. 110

The hippocampus is one of the first regions to demonstrate atrophy during prodromal AD. 111 Neurofibrillary tangle (NFT) accumulation within the MTL, including the hippocampus, is 112 one of the first hallmark pathological manifestations. Recent studies in cognitively healthy 113 114 individuals at risk for late-life AD have found that AD risk impacts the hippocampus in 115 midlife (Dounavi et al., 2022; 2020; Kerchner et al., 2014; for a review see Vilor-Tejedor et al. 2021). Dounavi et al. (2020) reported that the volume of the molecular layer of the 116 hippocampus was reduced in cognitively healthy midlife individuals with APOE ɛ4 genotype. 117 118 A more recent study (Dounavi et al., 2022) reported that the hippocampal fissure was

119 enlarged in middle-aged individuals with high CAIDE dementia risk scores.

Critically, the connectivity between the LC and the hippocampus is important for cognition. 120 121 Animal studies have demonstrated that the co-release of noradrenaline and dopamine from LC terminals in the hippocampus is critical for successful spatial learning and memory 122 (James et al., 2021; Kaufman et al., 2020; Kempadoo et al., 2016). Human studies in healthy 123 124 older adults are consistent with these data and show that higher LC-Hippocampus functional connectivity is significantly associated with better memory (Jacobs et al., 2015). While 125 accumulating evidence indicates that, individually, the LC and hippocampus are affected in 126 127 midlife by risk of late-life AD, it remains unknown whether their interactions and joint role in cognition are affected by AD risk at this stage. Understanding of changes in the interaction of 128 LC and the hippocampus during this preclinical period in at-risk populations has the potential 129 to not only shed light on the brain mechanisms of incipient AD, but also to provide urgently 130 needed early disease biomarkers. 131

To address this gap, we investigated the impact of three risk factors for late-life AD (APOE 132 ε4 genotype, FHD and CAIDE score) on cognition, and its relationship to the LC-133 134 Hippocampus functional connectivity. We tested a cohort of cognitively healthy middle-aged individuals, at baseline (N=210), and two-years follow-up (N=188) with a detailed 135 neuropsychological battery not restricted to functions usually implicated in dementia 136 137 detection in the older adults. Structural and functional Magnetic Resonance Imaging (MRI) data was obtained at both research visits. Our hypotheses were that risk of late-life AD would 138 be significantly associated with cognition, the LC-Hippocampus functional connectivity, and 139 140 their brain-behaviour relationship in middle-aged and cognitively healthy individuals. Furthermore, we expected that risk factors would be associated with longitudinal change of 141 cognition and functional connectivity in this cohort. 142

143

2. Methods 144

2.1 Participants 145

146	PREVENT-Dementia is an ongoing longitudinal multi-site research programme across the
147	UK and Ireland, seeking to identify early biomarkers of AD and elaborate on risk-mechanism
148	interactions for neurodegenerative diseases decades before the cardinal symptoms of
149	dementia emerge. Its protocol has been described in detail elsewhere (Ritchie & Ritchie,
150	2012). In the first PREVENT programme phase, participants were recruited at a single site,
151	via the dementia register database held at the West London National Health Service (NHS)
152	Trust, of the UK National Health Service, the Join Dementia Research website
153	(https://www.joindementiaresearch.nihr.ac.uk/), through public presentations, social media
154	and word of mouth. Procedures involving experiments on human participants were carried
155	out in accord with the ethical standards of the Institutional Review Board of Imperial College
156	London and in accord with the Helsinki Declaration of 1975. Approval for the study was
157	granted by the NHS Research Ethics Committee London Camberwell St Giles. Consented
158	participants were seen at the West London NHS Trust, where they underwent a range of
159	clinical and cognitive assessments (Ritchie & Ritchie, 2012). The cohort comprised
160	cognitively healthy volunteers aged 40-59 years. Here we examined baseline and follow-up
161	data from the West London dataset. 210 individuals (62 male; 148 female) were tested at
162	baseline, with 188 (89.5%) (55 male; 133 female) retained at two-years follow-up (Table 1).
163	Mild cognitive impairment and dementia were ruled out based on detailed clinical assessment
164	on each visit.

At baseline, 2 participants were missing APOE ɛ4 genotype information, and therefore the 165 166 CAIDE scores. At follow-up, 6 participants were missing information for calculating their

167	CAIDE scores, including the 2 missing APOE status. For cognitive tests, 2 participants were
168	missing data at baseline, and 12 at follow-up. Therefore, the cognitive analyses cohort was
169	N=208/N=176 at baseline/follow-up. 193 out of 210 participants finished MRI scanning at
170	baseline, and 169 out of 188 at follow-up. Furthermore, at baseline, 6 were excluded due to
171	incidental findings, and 58 due to not meeting the criterion of $>$ 150 frames with minimal
172	head motion (for details see the following section 2.4 MRI data acquisition and processing).
173	At follow-up, 3 were excluded due to incidental findings and 73 due to < 150 frames with
174	small head motion. The final fMRI analyses cohort was $N = 128/N = 93$ at baseline/follow-up.
175	Please see the Supporting Information (SI) file and STable 1–2 and SFigure 1 for a
176	description of any missing data.
177	
178	2.2 Biological sex
179	While the dichotomies of sex and gender are no longer considered to be sharply discrete, in
180	this study 'sex' was defined as an individual's natal or biological sex and was self-reported in
181	the Brain Injury Screening Questionnaire. The cohort is 70% female, reflecting higher female
182	interest in the study, possibly due to the known female predisposition to developing
183	Alzheimer's disease.
184	
185	2.3 Menopausal Status

Menopausal status was classified based on answers from a pregnancy and menstruation
survey administered during the clinical assessments. Self-reported menopausal status was
determined via answers to the question "Are you postmenopausal?". Those who answered
'Yes' were categorized as postmenopausal, and those who answered 'No' were categorized

as premenopausal. Of the 148/210 female participants, 55 (37.16%) were postmenopausal, 73

191 were premenopausal, and 20 who answered 'don't know'. Those who answered 'Don't Know'

192 (20/210) were excluded from further analyses.

193

- 194 2.4 Assessments
- 195 2.4.1 Risk factors
- 196 APOE ε4 Genotyping

197 The process of APOE ε 4 allele identification is outlined in detail in Ritchie et al. (2017). In

198 brief, genomic DNA was isolated from blood samples and APOE genotyping was performed.

199 All members of the research and clinical teams were blind to the result of APOE genotyping.

200 In this study, APOE ε 4 risk is determined by \geq 1 APOE ε 4 allele. 75/210 carried \geq 1 APOE ε 4

allele (See Table 1). The prevalence of APOE $\varepsilon 2\varepsilon 4$ in our cohort is 0.02% (see SFigure 2).

202 Investigation of the effect of the APOE $\varepsilon 2\varepsilon 4$ allele can be found in the SI. The exclusion of

203 APOE $\epsilon 2\epsilon 4$ carriers did not affect our main results and we therefore report results including

204 APOE $\varepsilon 2\varepsilon 4$ carriers.

205 Family History of Dementia

FHD was determined by a 'yes'/'no' question during clinical visits, where participants were asked whether a parent had a diagnosis of dementia. Participants were asked to include the dementia subtype if known, but answering 'yes' alone categorized a participant as FHD+. The answer 'no' likely captured both participants with no FHD, and participants for whom FHD was unknown. In summary, participants were defined as FHD+ if at least one parent was diagnosed with dementia. Cases where the FHD was unknown or partially known were

not recorded outside of the binary yes/no scoring. 103/210 were FHD+.

213 Cardiovascular Risk Factors, Aging, and Incidence of Dementia (CAIDE) score

214 CAIDE is a composite scale of estimated future dementia risk based on mid-life

- 215 cardiovascular measures (Fayosse et al., 2020; Sindi et al., 2015). It takes into consideration
- the individual's age, sex, educational attainment, APOE ɛ4 genotype, activity level, BMI,
- 217 cholesterol and systolic blood pressure (Kivipelto et al., 2006) and is scored on a range of 0 -
- 18. A higher score indicates greater risk. The CAIDE dementia risk score was calculated at

219 baseline and follow-up.

220

221 2.4.2 Cognitive testing

222 Cognitive function was assessed at baseline and follow up with the COGNITO

223 neuropsychological battery (Ritchie et al., 2014), designed to examine information processing across a wide range of cognitive functions in adults of all ages and not restricted to those 224 functions usually implicated in dementia detection in the older adults. Tests are administered 225 using a tactile screen to capture information processing time as well as response accuracy and 226 require about 40 minutes to complete. The tests, by order of presentation, are: reaction time; 227 reading; comprehension of phonemes, phrases, and syntax; focused and divided attention in 228 both visual and auditory modalities; visual working memory (visual tracking with auditory 229 interference); the Stroop test; immediate, delayed, and recognition trials for verbal recall 230 (name list); delayed recognition of spatial stimuli (faces); visuospatial associative learning; 231 visuospatial span; form perception; denomination of common objects; spatial reasoning; 232 copying of meaningful and meaningless figures; verbal fluency with se-mantic and phonetic 233 prompts; immediate recall of a narrative; immediate recall of a description of the relative 234

position of objects; vocabulary; implicit memory (recognition of new and previously learned
 material).

237 The COGNITO tests are designed to test several aspects of cognition, including attention (task: visual attention), memory (tasks: narrative recall, description recall, implicit memory, 238 name-face association, working memory), language (tasks: phoneme comprehension, verbal 239 fluency) and visuospatial abilities (task: geometric figure recognition) (Ritchie et al., 2014). 240 Based on previous studies (Ritchie et al., 2021; Ritchie et al., 2018; Ritchie et al., 2017), 11 241 summary variables from the COGNITO battery capturing the above functions were used here 242 (for a list see STable 2). 243 Additionally, we used the Visual Short-Term Memory Binding task (VSTMBT) (Parra et al., 244

245 2010), a computer-based task that assesses visual short-term memory binding of single features, e.g., complex shape or color combinations, or feature conjunctions, e.g., shape and 246 color combinations. In the single feature condition, participants must identify whether the test 247 stimuli (three random 6-sided polygons) are the "same" as or "different" to the studied 248 stimuli in terms of shape (shape only) or color (color only). In the binding condition, 249 250 participants are required to correctly identify if both the shape and color of the test stimuli match studied stimuli. The two summary variables from the VSTMBT were the percentage of 251 correctly recognized items from the two conditions. 252

The Addenbrooke's Cognitive Assessment (ACE) III (Hsieh et al., 2013) was recorded at the follow-up session, but not at the baseline. Analyses that focused on the subset of assessments performed at both testing sessions did not include the ACE III.

256

257 2.5 Behavioural data analyses

258 Rotated principal component analysis

259 Rotated principal component analysis (rPCA), a dimensionality reduction technique, was adopted to cluster the above-mentioned 13 cognitive measures into related domains, reduce 260 the number of multiple comparisons between the cognitive tests (Jolliffe & Cadima, 2016). 261 This step maximized power to investigate the impact of risk factors on cognition and brain-262 behaviour relationships. rPCA was conducted by using the psych package (Version 2.0.12) in 263 R software (https://www.r-project.org/), following steps as: (1) Preprocessing. Listwise 264 deletion procedure was used to deal with the missing data. Only participants who had no 265 missing observations were kept (N = 208 at baseline; N = 176 at follow-up). (2) Component 266 267 extraction and estimation. Principal component analysis was used to extract a smaller number of components that capture the most variance of the initial variables, representing the 268 common cognitive processes underlying the 13 original cognitive measures. Scree plots and 269 parallel analysis were used to determine the number of components (Horn, 1965). Parallel 270 analysis generated a set of random correlation matrices (n = 500) by using the same number 271 272 of variables (n = 13) and participants (N = 208 at baseline; N = 176 at follow-up). Then, we compared the solution to the random simulated normal data. The eigenvalues obtained from 273 the actual matrix were compared with those obtained from the randomly generated matrices, 274 275 and the estimated number of components was determined by selecting components with eigenvalues larger than 95th percentile of the randomly generated components' eigenvalues 276 (See also SI). (3) Rotation. To examine the component structure among the initial variables, 277 278 an orthogonal rotation method Varimax that constrains the components to be uncorrelated, 279 was applied to the coefficient matrix (Kaiser, 1958). (4) Component coefficients. Component coefficients (r), reflecting the correlation between each of the components and each of the 280

281 original variables, were thus obtained. Given the sample size in this study (N > 200), component coefficients that were > 0.4 were deemed practically significant (Hair Jr. et al., 282 1998; Stevens, 1992). (5) Component scores. Component scores were calculated by a 283 regression method where the regression weights were found from the inverse of the 284 correlation matrix times the coefficient of each variable on the corresponding components. 285 The coefficients were used to interpret the components, and component scores were used for 286 287 the following analyses. (6) Component similarities. Components were extracted at the baseline and follow-up respectively. To test the similarity of components extracted at each 288 289 study session, to be capturing similar cognitive functions, Tucker's congruence coefficient was calculated (Lorenzo-Seva & ten Berge, 2006). In subsequent analyses, we used these 290 cognitive components, rather than the individual tests to measure the impact of risk factors on 291 292 cognition and brain-behaviour relationships.

293

294 2.6 MRI data acquisition and processing

Imaging data were obtained as part of a multimodal examinations in a 3T Siemens Verio 295 MRI scanner and with 32-channel head coil (https://preventdementia.co.uk/for-researchers/). 296 Resting-state fMRI data were acquired with T2*-weighted echo-planar imaging (EPI) 297 sequence. Total acquisition time for the resting state scan was 11 minutes 06 sec. The eyes 298 were closed during rest. 330 volumes were acquired, and each volume contained 35 slices 299 (interleaved acquisition), with slice thickness of 3 mm (repetition time (TR) = 2000 ms, echo 300 time (TE) = 30ms, flip angle (FA) = 80°, Field of View (FOV) = 192×192 mm², voxel size = 301 3 mm³ isotropic). A 3D T1-weighted magnetization prepared rapid gradient-echo image 302 (MPRAGE, 160 slices, voxel size = 1 mm^3 isotropic, TR = 2300ms, TE = 2.98ms, FOV = 303 240×256 mm², FA = 9°) was also acquired. All scans were repeated after approximately two 304

305 years on the same scanner using the same protocol.

Standard preprocessing procedures for resting-state fMRI data were performed with SPM12 306 307 (Statistical Parametric Mapping, https://www.fil.ion.ucl.ac.uk/spm/software/spm12/) and the AA pipeline software (Cusack et al., 2015) implemented in MATLAB R2019a (The 308 MathWorks, United States). We discarded the first 5 volumes to allow for adaptation to the 309 310 environment and equilibrium of the MR signal. In this pipeline (Figure 1a), we performed slice timing correction, motion correction, co-registration of functional and structural images, 311 normalization into standard MNI space, and spatial smoothing. Spatial normalization was 312 313 performed using SPM12's segment-and-normalize procedure, whereby the T1 structural was segmented into grey matter (GM), white matter (WM) and cerebrospinal fluid (CSF) and 314 normalized to a segmented MNI-152 template. These normalization parameters were then 315 applied to all EPIs. The quality of spatial normalization was visually inspected for each 316 participant and no participants showed a normalization failure. The data were then smoothed 317 318 with a Gaussian kernel of 6mm full width at half maximum. We also applied a general linear 319 model (GLM) that includes 24 head motion parameters -6 rigid-body motion parameters, the first-order temporal derivative of them and quadratic terms of the original parameters and 320 their derivatives (Satterthwaite et al., 2013) – and averaged signals from WM and CSF as 321 covariates to reduce any residual effects of head motion and physiological noise. Lastly, all 322 preprocessed data were temporally band-pass filtered (0.01-0.08 Hz) to remove low-323 frequency drift and high-frequency physiological noise (Salvador et al., 2008; Zuo et al., 324 2010). 325

326

Figure 1 about here please

327

328 Because brain signal from the brainstem is susceptible to head movements, we performed

329 careful measurement and control for head motion, by using a "scrubbing" procedure (Power et al., 2012). We first calculated an index of data quality called framewise displacement (FD) 330 (for details please also see SI), which measures how much the head has changed position 331 from one frame to the next, and can therefore be used to flag frames of suspect quality based 332 on the threshold of FD > 0.2 mm, in order to remove motion-contaminated data before 333 calculating functional connectivity (Power et al., 2012). A participant inclusion criterion was 334 then set to ensure a minimum amount of data remained after scrubbing, i.e., at least 150 335 frames ($\sim 5 \text{ min}$) of data (Figure 1b), based on wide agreement that $\sim 5 \text{ min}$ of fMRI data is an 336 337 adequate for resting state functional connectivity analyses (Van Dijk et al., 2010). After completing the scrubbing procedure, the included participants had $74.0\% \pm 14.8\%$ frames at 338 baseline and $73.5\% \pm 15.0\%$ at follow-up. Finally, we calculated the mean FD across the 339 remaining good frames, which was further regressed in the group-level analyses (Power et al., 340 2014). The mean FD was 0.12 ± 0.01 at baseline and 0.12 ± 0.01 at follow-up. There were no 341 significant associations of mean FD with any of the risk factors or the LC-Hippocampus 342 functional connectivity (See SI). The fMRI-based exclusion criteria are also illustrated in 343 Figure 1b. 344

To ensure high accuracy in the localization of the LC, we created a binary LC mask of 3 mm³ 345 isotropic voxel size that closely matched with two previously published LC maps (Dahl et al., 346 2022; Ye et al., 2021), which were developed in the standard MNI space with 0.5 mm³ 347 isotropic voxel size. In order to achieve the highest precision, instead of resampling the LC 348 maps to match the fMRI data's voxel size (3 mm³) using automatic scripts, we manually drew 349 350 a 3 mm³ binary LC mask in MNI space with reference to the two LC maps. This is because the automatic scripts did not perform well when resampling the two referenced LC maps 351 352 from high to low resolution due to the small size of these two LC regions of interest. The FSL 353 software (https://fsl.fmrib.ox.ac.uk/fsl/fslwiki/) was used for the drawing process, by

following the same procedure as in Plini et al. (2021). The two 0.5 mm³ LC maps were 354 superimposed on a blank 3 mm³ template in standard MNI space. The drawing process was 355 then performed on this blank template by adding or removing voxels to include different 356 anatomical locations of the two referenced maps without crossing the dorsal and medial raphe 357 nuclei, as defined by Beliveau et al. (2015), and the cerebellar vermis white matter. At the 358 end of this procedure, the new bilateral LC mask was symmetrical and consisted of 16 3 mm³ 359 360 voxels, was anatomically inclusive of the 7T LC map (Ye et al., 2021) and the LC meta-map (Dahl et al., 2022), and did not encroach on other pontine nuclei (Figure 1c). 361

362 In an independent study (Dounavi et al., 2022), T1-weighted MPRAGE of this cohort were corrected for field inhomogeneities using the Advanced Normalisation Toolbox (ANTs) N4 363 algorithm (Tustison et al., 2010) and processed using Freesurfer version 7.1.0 (Desikan et al., 364 2006). The *recon-all* pipeline was run for every participant with standard settings. The brain 365 masks and surfaces were inspected following recon-all and manual corrections were applied: 366 367 (a) in the form of erosion of non-brain voxels from the brain mask or non-WM voxels from the WM mask, (b) in the form of filling of areas where the brain was not correctly identified, 368 or (c) with the addition of control points in cases where white matter was not successfully 369 370 identified. Following these steps, we generated individual masks of the hippocampus, which were then applied to the fMRI data to extract the time course of the region (Figure 1d). The 371 functional connectivity (FC) between the LC and the hippocampus was derived using 372 Pearson's correlation coefficient (r) on the denoised time courses (Figure 1d). The r value 373 was then transformed to z value using Fisher's r-to-z transformation to improve normality for 374 375 further statistical analysis. To avoid the formation of artificial anti-correlations (Anderson et al., 2011; Murphy et al., 2009), we did not perform global signal regression. In addition, 376 volume of the hippocampus (HCV) was derived for each participant at baseline and follow-377 378 up as a covariate for functional connectivity analyses.

379

380 2.7 Statistical approach

381 All statistical analyses were performed in R software. The normality of the data was assessed

by combining the visualization of a quantile-quantile plot and the Shapiro–Wilk test.

383 Demographic and clinical information of the study cohort was analyzed across risk groups

using chi-square (χ^2 tests) for categorical variables and Mann Whitney U tests for continuous

385 (discrete) variables, given that they were not normally distributed in this cohort.

To investigate the cross-sectional effects of risk factors (APOE ɛ4 genotype and FHD) on 386 cognition and brain function we used multiple linear regression models. Age, years of 387 388 education were included as covariates for cognition models, and mean FD values and HCV were additionally included as a covariate for models of functional connectivity. The effects of 389 APOE ɛ4 and CAIDE risk factors were modelled independently to avoid modelling the 390 391 variance associated with APOE genotype in the same model twice. Spearman correlation analyses were used to assess the associations between cognition and CAIDE score given the 392 non-normal distribution of CAIDE in our cohort. For the CAIDE model, we did not control 393 for the age, years of education, as they are considered in the CAIDE scoring. 394

For any observed risk effect on cross-sectional changes of cognition or functional 395 396 connectivity, interactions between risk factors and the LC-Hippocampus functional connectivity on cognition were assessed through multiple regression models with the 397 cognition as the dependent variable, the LC-Hippocampus FC, risk factor and the risk × FC 398 interaction as independent variables, and age, years of education, and HCV as covariates. 399 Age, sex, and years of education were not included as covariates when CAIDE was included 400 401 in the models, because CAIDE includes these variables in its scoring. Biological sex was investigated as variables of interest in these analyses especially given the higher proportion of 402

females in the sample. If a sex effect was found, any interactions with risk factors, and the
effect of menopausal status was also considered because the cohort age range [52+/- 11 years]
cover menopause transition in females, which is known to affect cardiovascular health and
cognition.

407 For any observed interactions, we plotted the regression of LC–Hippocampus FC on

408 cognitive performance for each level/value of the risk factor, to interpret the effect (Aiken &

409 West, 1991). For the continuous variable CAIDE, we applied the Johnson-Neyman technique

410 (Bauer & Curran, 2005; Johnson & Fay, 1950), to test the significance of the slopes of the

411 simple regression lines on each value of the CAIDE scores (ranged from 0-18). This

412 examined the CAIDE interval at which the brain-cognition coupling was significant. All

413 continuous variables in the interaction terms, i.e., CAIDE and FC, in the moderation models

414 were mean-centered (subtracting each the mean from each value) to avoid multicollinearity.

Longitudinal changes were first assessed using random intercept linear mixed effect models (Cnaan et al., 1997) for the entire cohort. Specifically, we evaluated longitudinal changes of cognition (Eq. 1) and functional connectivity (Eq. 2) over two years.

418

$$Cognition_{jt} = \beta_0 + \beta_1 (Time_{jt}) + u_j + r_{jt}$$
(1)

$$FC_{jt} = \beta_0 + \beta_1 (Mean FD_{jt}) + \beta_2 (HCV_{jt}) + \beta_3 (Time_{jt}) + u_j + r_{jt}$$
(2)

419

420 For each participant j, measures of cognition/functional connectivity at each time point t

421 served as dependent variables, and the timepoint (a categorical variable with two levels:

422 baseline and follow-up) served as the independent variable. Mean FD and HCV at each time

423 point t were included as covariates for the functional connectivity model only (Eq. 2). u_i

424 represented individual variability in the intercepts, i.e., a random effect.

425 The presence of a longitudinal change for the entire cohort was then further assessed to

426 determine whether it varied in relation to the risk factors, as assessed by the interaction term

427 ($Risk_i \times Time_{it}$). Age at baseline, sex, and years of education, were included as

- 428 covariates for the cognition models that did not include CAIDE (Eq. 3). Mean FD and HCV
- 429 were additionally included as covariates for the functional connectivity models (Eq. 4).

430

$$\begin{aligned} Cognition_{jt} \\ &= \beta_0 + \beta_1 (Age_j) + \beta_2 (Sex_j) + \beta_3 (Education_j) + \beta_4 (Risk_j) \\ &+ \beta_5 (Time_{jt}) + \beta_6 (Risk_j \times Time_{jt}) + u_j \\ &+ r_{jt} \end{aligned}$$

431

$$\begin{aligned} FC_{jt} &= \beta_0 + \beta_1 (Age_j) + \beta_2 (Sex_j) + \beta_3 (Education_j) + \beta_4 (Mean FD_{jt}) \\ &+ \beta_5 (HCV_{jt}) + \beta_6 (Risk_j) + \beta_7 (Time_{jt}) + \beta_8 (Risk_j \times Time_{jt}) \\ &+ u_i + r_{it} \qquad (4) \end{aligned}$$

432

Lastly, the presence of risk related longitudinal change in cognition or FC was further
investigated to determine whether risk factors influence brain-cognition association
longitudinally. Multiple linear regression modelling was used with a percentage change score
[i.e. (follow-up - baseline)/baseline] in cognition as the dependent variable, and a risk factor,
a percentage change in FC, and their interaction as independent variables. Baseline age, sex,
years of education and changes in HCV were included as covariates. Only changes in HCV
were controlled for when assessing CAIDE in the models.

Although the baseline and follow-up datasets were not independent, we considered each in its 440 own right, in addition to testing for longitudinal changes in cognition and functional 441 connectivity. During the two years, a proportion of our participants may have sustained brain 442 health that are vet subthreshold to clinical manifestations, or that may not vet manifest as 443 longitudinal change. Statistical results were corrected for multiple comparisons. 444 445 3. Results 446 3.1 Demographic characteristics 447 Demographic characteristics of the cohort at baseline and follow-up, stratified by APOE E4 448 genotype and FHD, are shown in Table 1. There were no significant differences in age, sex or 449 years of education between the groups. APOE ɛ4 allele genotype was more frequently found 450 in the FHD+ than FHD- group at baseline (p = 0.01) and follow up (p = 0.02). CAIDE scores 451 (including APOE status) were significantly higher for the FHD+ than FHD- group at baseline 452 (p = 0.03) and follow up (p = 0.003). Naturally, CAIDE scores including APOE status were 453 454 significantly higher for the APOE ε 4+ than APOE ε 4- group at baseline (p = 0.0003) and follow up (p = 0.0002), but when APOE status was excluded the CAIDE scores did not differ 455 between the two groups (Table 1). 456 457

458

Table 1 about here please

460 3.2 Cognitive domains

461	We first performed parallel analyses and plotted scree plots to determine the number of
462	components that best represented the original 13 cognitive measures at baseline and follow-
463	up, separately. Results showed that eigenvalues of the first three components were larger than
464	95 th percentile of the randomly generated eigenvalues at both study timepoints (Figure 2a),
465	which indicated that the three component (C) solution best represented the data. The three
466	components were then rotated to be uncorrelated with each other, and could cumulatively
467	explain a total of 41% percentage of the variance, at baseline ($C1 = 17\%$; $C2 = 13\%$; $C3 =$
468	12%) and 40% at follow-up (C1 = 17%; C2 = 12%; C3 = 11%) (Figure 2b).
469	

471

470

The loading values (weights) of the cognitive measures (Figure 3a) reflect the relationships between the original measures and the corresponding components. Measures with higher coefficients were more closely related to the components. Based on the cognitive functions that the highest loading measures tapped into (for details see STable 2), in subsequent analyses we refer to: C1 as 'episodic and relational memory'; C2 as 'working and short-term (single-feature) memory'; C3 as 'verbal, visuospatial functions, and short-term (conjunctive) memory.'

The highest loading measures and their coefficients for each component differed for the
cross-sectional and longitudinal data. For example, verbal functions, visuospatial functions,
and short-term (conjunctive) memory loaded most strongly on the third component at
baseline. At follow-up, implicit memory rather than short-term memory loaded on the third

perpetany.	
All rights reserved. No reuse allowed without pe	armission
All rights reserved. No reuse allowed without pe	CITII331011.

483	component (Figure 3a). We found high similarity between baseline and follow-up for C1 (ϕ
484	= 0.90), and C3 (ϕ = 0.76), with relatively low similarly for C2 (ϕ = 0.40) (Figure 3b).
485	
486	Figure 3 about here please
487	
488	3.3 Impact of risk factors on cognition
489	Episodic and relational memory
490	At baseline, the multiple linear regression model showed a significant positive association
491	between episodic and relational memory with APOE $\varepsilon 4$ allele [β (SE) = 0.28 (0.13), $p < 0.05$],
492	independently of sex, age and years of education (Table 2). At follow-up, there was a trend
493	effect of APOE $\varepsilon 4$ allele [β (SE) = 0.27 (0.15), p = 0.07]. Education was significantly
494	positively associated with this domain at baseline [β (SE) = 0.09 (0.02), $p < 0.0001$] and
495	follow-up [β (SE) = 0.10 (0.02), $p < 0.0001$]. Sex did not interact with the APOE ε 4 allele in
496	its effect on episodic and relational memory. The baseline effect of APEO ɛ4 on episodic and
497	relational memory became a trend effect (SFigure 6) when corrected for multiple
498	comparisons and needs to be interpreted with caution.
499	
500	Table 2 about here please
501	

502 The multiple linear regression models showed no associations between FHD and episodic and relational memory, at either timepoint. Spearman correlation analyses showed no significant 503

associations between the CAIDE score and episodic and relational memory, at either

505 timepoints.

506

507 Working and short-term (single-feature) memory

508 We found no associations between APOE ɛ4, FHD, or CAIDE score with working and short-

509 term (single-feature) memory. Education was significantly positively associated with

510 performance in this domain, independently of the other factors, at baseline [β (SE) = 0.04

511 (0.02), p < 0.05] only.

512

513 Verbal, visuospatial functions, and short-term (conjunctive) memory

We found no associations between APOE ϵ 4 or FHD with verbal, visuospatial functions, and short-term (conjunctive) memory. Age was significantly negatively associated with performance in this domain, independently of the other factors, at baseline [β (*SE*) = -0.04 (0.01), p < 0.005]. The CAIDE score was significantly associated with verbal, visuospatial functions, and short-term (conjunctive) memory at baseline (rho = -0.17, p < 0.05, Figure 4a), and at follow-up (rho = -0.20, p < 0.01, Figure 4b). Higher CAIDE scores were significantly associated with worse performance.

521 As the cohort was 70% female, and the age range (40-60 years) covers the menopausal

522 transition, which is known to affect cardiovascular health and cognition, the effect

523 menopausal status on the relationship between CAIDE score and cognition was investigated.

524 The CAIDE score includes sex in its calculation, with 'male' scored higher. Therefore, the

525 effect of sex on relationship between the CAIDE risk and cognition not investigated.

526	However, there was a significant interaction between CAIDE and menopause on verbal,
527	visuospatial functions and short-term (conjunctive) memory at baseline [β (SE) = -0.20 (0.09),
528	$p < 0.05$] and at follow-up [β (SE) = -0.29 (0.13), $p < 0.05$] (Table 3). Increasing CAIDE was
529	associated with worse cognitive performance in post-menopausal females, but not in pre-
530	menopausal counterparts. This effect was controlled for age effects, as post-menopausal
531	women were overall older than pre-menopausal counterparts (post-: median \pm IQR: 49.0 \pm
532	8.0 vs pre-: 56.0 ± 4.0).
533	
555	
534	Figure 4 about here please
535	Table 3 about here please
536	
537	There was no significant longitudinal change of cognition in any of the three cognitive
538	domains over two years.
539	
557	
540	3.4 Longitudinal change of LC-Hippocampus functional connectivity
541	We found a significant negative association between time and the LC-Hippocampus
542	functional connectivity [β (SE) = -0.06 (0.03), $p < 0.05$], after controlling for mean FD and
543	HCV (Table 4), indicating reduced connectivity at two-years follow-up relative to baseline
544	across the whole cohort (Figure 5). Sex did not interact with time in affecting the LC-
545	Hippocampus functional connectivity.
546	

547	Table 4 about here please
548	Figure 5 about here please
549	
550	3.5 Impact of risk factors on the LC-Hippocampus functional connectivity and brain-
551	behavior relationships
552	There were no significant associations of APOE ɛ4, FHD, or CAIDE scores with the LC-
553	Hippocampus functional connectivity, cross-sectionally, or longitudinally.
554	At baseline, none of the risk factors moderated the relationship between functional
555	connectivity and cognition. At follow-up, the multiple linear regression model with CAIDE,
556	LC–Hippocampus FC, and CAIDE \times LC–Hippocampus FC interaction term as independent
557	variables, and verbal, visuospatial functions, and short-term (conjunctive) memory as the
558	dependent variable showed a significant negative association between the interaction term
559	(CAIDE \times LC–Hippocampus FC) and verbal, visuospatial functions, and short-term
560	(conjunctive) memory [β (SE) = -0.54 (0.22), $p < 0.05$] (Table 5). In individuals with
561	low/high CAIDE scores, higher functional connectivity was associated with better/worse
562	cognition (Figure 6). Menopausal status did not have an effect on this relationship.
563	
564	Table 5 about here please
565	Figure 6 about here please
566	

567 4. Discussion

568	It is well acknowledged that AD neuropathology starts decades before clinical manifestations,
569	but the brain mechanism of sporadic AD and its interactions with risk factors in midlife
570	remain unclear. We examined whether risk factors for late-life AD were associated with
571	cognition, functional connectivity between two of the key structures in AD neurogenesis -
572	the LC and the hippocampus – and the brain-behaviour relationship between them, cross-
573	sectionally and longitudinally in a cohort of middle-aged, cognitively healthy individuals. We
574	found that the CAIDE scores were significantly negatively associated with verbal,
575	visuospatial functions and short-term (conjunctive) memory functions in this cohort.
576	Menopausal status significantly interacted with CAIDE on cognition, independently of age.
577	Post-menopausal women showed worse cognition with increasing CAIDE scores.
578	Furthermore, the CAIDE score significantly moderated the relationship between cognition
579	and the LC-Hippocampus functional connectivity. These results shed light on some of the
580	earliest brain-behaviour alterations due to risk of future AD in cognitively healthy
581	individuals cross-sectionally. We found a significant longitudinal effect of age on the LC-
582	Hippocampus FC, with no longitudinal effects of risk factors on cognition, the LC-
583	Hippocampus functional connectivity, or their relationship. The lack of prominent
584	longitudinal effects of risk factors is likely due to the narrow follow-up time window (2 years)
585	in this relatively young midlife cohort, an estimated 23 years from dementia onset (Dounavi
586	et al., 2022).
50 7	Higher CAIDE george were gignificantly appropriated with worse cognition in verbal

587 Higher CAIDE scores were significantly associated with worse cognition in verbal,

visuospatial functions and short-term (conjunctive) memory, both at baseline and at follow-

589 up. Previous studies from the same cohort have reported significant negative associations of

590 CAIDE scores with visuospatial functions and navigational abilities (Ritchie et al., 2018;

591 Ritchie et al., 2017). We did not observe a longitudinal effect, similarly to other studies of this cohort (Dounavi et al., 2021; Low et al., 2021), possibly due to the relatively young age 592 and the short follow-up window (Ritchie et al., 2018; Ritchie et al., 2017). Post-menopausal 593 women only showed worse cognition with increasing CAIDE scores. The menopausal 594 transition is a middle-age phenomenon with knock-on effects on endocrinal (Weber et al., 595 2014), cardiovascular (Yanes & Reckelhoff, 2011), and metabolic (Anagnostis et al., 2019; 596 597 Rahman et al., 2020) systems that significantly impact brain health (Mosconi et al., 2018; Udeh-Momoh & Watermeyer, 2021). The rapid decline in estrogen, a key hormonal change 598 599 during menopausal transition, is considered to increase vulnerability to AD in females (Scheyer et al., 2018; Udeh-Momoh & Watermeyer, 2021). Post-menopause has been 600 associated with increased levels of A β , elevated tau burden, reduced white matter integrity, 601 602 and glucose hypometabolism (Buckley et al., 2022; Mosconi et al., 2017; Rahman et al., 603 2020). Our results suggest that post-menopausal status exacerbates the effect of cardiovascular AD risk on cognition, rendering females particularly vulnerable to cognitive 604 decline, and conversely, that addressing cardiovascular risk early may protect cognition in 605 post-menopausal females. 606

607 We found that the LC-Hippocampus functional connectivity decreased significantly over two years. The previously reported effect of age on the LC-Hippocampus functional connectivity 608 is inconsistent, possibly due to different age ranges used in different studies. In healthy young 609 and early midlife adults (age range: 18–49), LC functional connectivity to the 610 parahippocampus was negatively associated with age (Zhang et al., 2016). However, another 611 612 study of a lifespan adult cohort (age range: 19-74) found no association of age with the LC-Hippocampus/parahippocampus functional connectivity (Jacobs et al., 2018). In a broader 613 age-range lifespan cohort (age range: 8–83), Song et al. (2021) found a positive quadratic age 614 615 effect (u-shape) on LC-parahippocampal functional connectivity, suggesting an increased

interaction between these two regions in children and older adults. Therefore, our finding
advances the literature, by showing that the LC–Hippocampus functional connectivity
decreases with age in midlife.

Another key question here was whether risk factors impact the LC–Hippocampus functional 619 connectivity and its role in cognition. We found that the CAIDE score significantly 620 moderated this brain-behaviour relationship. Stronger LC-Hippocampus functional 621 connectivity was associated with better cognition for individuals with low CAIDE score, and 622 with worse cognition in those with high scores. The presence of this effect in the follow-up, 623 624 but not the baseline dataset is likely due to the older age of the cohort at follow-up. Previous studies in older adults found that stronger LC-Hippocampus functional connectivity was 625 significantly associated with better cognition (Jacobs et al., 2015). Our finding advances 626 understanding by suggesting that the connectivity between the LC and the hippocampus 627 supports cognition in midlife individuals who have low dementia risk scores. 628

Furthermore, our result suggests that, in the presence of high dementia risk scores, the LC-629 hippocampus functional connectivity is negatively associated with cognition. Previous studies 630 631 have shown that a high CAIDE score, of more than 12, confers a probability of 16.4% for future dementia, with scores above 8 associated with a probability of more than 4% 632 (Kivipelto et al., 2006). In a study of middle-aged individuals with a mean age of 46, 633 participants with a score greater than 8 had a 29% 40-year risk for dementia (Exalto et al., 634 2014). These findings suggest that a proportion of our study's participants, who have a 635 CAIDE score greater than 8, may develop dementia. Assuming that AD pathology is 636 637 underway in this proportion of individuals with high dementia risk scores, our result lends 638 indirect support to the hypothesis that LC hyperactivity and associated hyper-connectivity initiates the spread of pathological tau to MTL (Weinshenker, 2018), and particularly the 639

hippocampus, in the early stages of AD. The ensuing neurodegeneration may explain why 640 hyper-connectivity is associated with worse performance in individuals with high dementia 641 risk scores. However, the neural mechanisms of AD neuropathology genesis remain 642 contested. Given the lack of tau biomarker status for this cohort, future studies that integrate 643 tau and functional connectivity information are required to further test the proposal that tau 644 spread from the LC to the hippocampus via functional connectivity pathways. Furthermore, 645 646 future studies from the continuing longitudinal follow-up of this cohort will validate the role and nature of these changes in individuals who display AD neuropathology as they get older. 647 648 We did not observe any significant effects of risk factors for AD on the LC-Hippocampus connectivity. Previous studies on MCI and AD patients have shown disrupted/lower LC-649 Hippocampus functional connectivity (MCI: Jacobs et al., 2015; age: 65.1 ± 4.5 years) (Liebe 650 et al., 2022; age: 73.3 ± 7.5 years; AD patients: Zhao et al., 2017), compared to age-matched 651 healthy controls. One previous study investigating cognitively healthy middle-aged adults 652 653 with familial risk for late-onset dementia found no effect of AD risk on the LC-Hippocampus connectivity (Del Cerro et al., 2020; age: 50.4 ± 8.3 years). Consistent with this previous 654 study, our null result suggests that AD risk factors are not associated with the LC-655 Hippocampus connectivity in middle-aged individuals who are cognitively healthy. 656 We also observed a trend effect association of APOE ɛ4 allele with better cognition in 657

episodic and relational memory, independently of sex, age and years of education. Previous research has suggested that the APOE ϵ 4 allele, although being associated with poorer health outcomes in old age, may have a positive effect earlier in life. For example, it has been associated with better visual perception (Ritchie et al., 2017), short-term memory (Zokaei et al., 2020), higher intelligence quotient and a more economic use of memory-related neural resources in young healthy humans (Mondadori et al., 2007). These results may also be

interpreted based on the antagonistic pleiotropy hypothesis of aging (William 1957), which 664 proposes that deleterious genes, such as the APOE E4 gene allele, have survived through 665 evolution because they might confer an advantage early in life, when humans are 666 reproductively fit. These results suggest an impact of genetic risk for late-life sporadic AD in 667 cognitively healthy midlife individuals on episodic memory, the hallmark cognitive 668 dysfunction in early clinical dementia and mild cognitive impairment (Irish et al., 2011). 669 Thus, they highlight the importance of studying the origins of Alzheimer's disease in this age 670 group, over 20 years before typical symptom onset. Given the weak effect however, further 671 672 replication in larger samples is warranted to validate these findings. In summary, the PREVENT-Dementia study is a longitudinal multi-site study targeted at 673 identifying early biomarkers for Alzheimer's disease. In this study, we provide early proof 674 that the recruited cohort, with a mean age of 52/54 years at baseline/follow-up, showed a 675 disrupted role of LC-Hippocampus connectivity in cognition with an increasing CAIDE 676 677 score. APOE ɛ4 genotype or FHD did not moderate this brain-behaviour relationship. In other words, when the genetic risk alone was considered, no alterations of brain-behaviour 678 relationships were found. It was only when a risk score (CAIDE) incorporating genetic risk in 679 680 combination with lifestyle factors, sex and age was considered, that such alterations were unravelled. Taken together, these results provide strong evidence that brain-behaviour 681 alterations in individuals with higher CAIDE scores may be driven by lifestyle risk factors 682 included in this dementia risk score (i.e., blood pressure, cholesterol, physical activity, body 683 mass index, years of education).

685 Hence, these findings highlight the importance of considering modifiable risk factors when stratifying risk populations, or potentially designing randomized control trials. Our previous 686 work in this cohort has demonstrated that modifiable lifestyle factors affect cognition, 687

684

particularly in individuals at high risk for late-life AD (Heneghan et al., 2023) and lends
further support to this idea. In fact, a randomized multi-domain control trial in the FINGER
population (mean age 70 years old) over two years demonstrated that the applied lifestyle and
vascular interventions had an impact on cognition (Stephen et al., 2019). Hence, further
investigation of pathological alterations in relation to modifiable risk factors in midlife is
warranted to unveil the sequalae of brain and behaviour alterations leading to dementia.

694 4.1 Strengths and Limitations

Strengths of this study are its large well-characterized middle-aged cohort. State-of-the-art 695 analysis methods and a thorough quality control protocol were applied to the acquired data. 696 Limitations include the short follow-up window of the study and the absence of further well-697 698 established preclinical biomarkers, such as amyloid and tau status. The interpretability of the effect of risk on individual cognitive abilities requires individuation in future studies. 699 Standard MR functional sequences provide limited resolution of the diminutive LC volume, 700 and thus, higher resolution imaging that specifically targets the LC region is required to 701 further validate these findings in future studies. Like most other studies involving mid-life 702 703 and older people in the UK and Ireland, the study population is mainly (95%) of white Caucasian ethnicity, not dissimilar to the historic ethnic mix of older people in the five study 704 sites in this region. The inclusion of other ethnicities and representation from different world 705 706 regions in future studies will ensure true diversity and findings that can be generalized to different populations across the globe. 707

708 Acknowledgement

- F.D. was funded by the Provost PhD Award Scheme from Trinity College Dublin, to
- 710 L.N. L.N. was also funded by a L'Oréal-UNESCO for Women In Science International
- 711 Rising Talent Award, the Welcome Trust Institutional Strategic Support grant, and the Global
- 712 Brain Health Institute Project Grant.
- 713 The PREVENT-Dementia study is supported by the UK Alzheimer's Society (grant numbers
- 178 and 264), the US Alzheimer's Association (grant number TriBEKa-17-519007) and
- 715 philanthropic donations.
- 716 We thank all PREVENT-Dementia participants for their enthusiastic participation in this
- study. We also thank the Research Delivery service at West London NHS Trust and the
- 718 Wolfson Clinical Imaging Facility at Imperial College London for their support in running

the study.

720

- 721 Conflict of interest
- The authors declare no conflict of interest.

724 References:

725	Aiken, L. S., & West, S. G. (1991). Multiple regression: Testing and interpreting interact	ions
726	Newbury Park, CA: Sage.	

- Amaral, D. G., & Sinnamon, H. M. (1977). The locus coeruleus: neurobiology of a central
 noradrenergic nucleus. *Prog Neurobiol*, 9(3), 147-196. <u>https://doi.org/10.1016/0301-</u>
 0082(77)90016-8
- 730 Anagnostis, P., Christou, K., Artzouchaltzi, A. M., Gkekas, N. K., Kosmidou, N., Siolos, P.,
- 731 Paschou, S. A., Potoupnis, M., Kenanidis, E., Tsiridis, E., Lambrinoudaki, I.,
- 732 Stevenson, J. C., & Goulis, D. G. (2019). Early menopause and premature ovarian
- insufficiency are associated with increased risk of type 2 diabetes: a systematic review
 and meta-analysis. *Eur J Endocrinol*, *180*(1), 41-50. <u>https://doi.org/10.1530/eje-18-</u>
 0602
- 736 Anderson, J. S., Druzgal, T. J., Lopez-Larson, M., Jeong, E. K., Desai, K., & Yurgelun-Todd,
- D. (2011). Network anticorrelations, global regression, and phase-shifted soft tissue
 correction. *Hum Brain Mapp*, *32*(6), 919-934. <u>https://doi.org/10.1002/hbm.21079</u>
- Arnsten, A. F. T., Datta, D., Del Tredici, K., & Braak, H. (2021). Hypothesis: Tau pathology
 is an initiating factor in sporadic Alzheimer's disease. *Alzheimer's & Dementia*, *17*(1),
 115-124. https://doi.org/10.1002/alz.12192
- 742 Aston-Jones, G., & Cohen, J. D. (2005). An integrative theory of locus coeruleus-
- norepinephrine function: adaptive gain and optimal performance. *Annu Rev*
- 744 *Neurosci*, 28, 403-450. <u>https://doi.org/10.1146/annurev.neuro.28.061604.135709</u>
- 745 Barnes, D. E., Beiser, A. S., Lee, A., Langa, K. M., Koyama, A., Preis, S. R., Neuhaus, J.,

746	McCammon, R. J., Yaffe, K., Seshadri, S., Haan, M. N., & Weir, D. R. (2014).
747	Development and validation of a brief dementia screening indicator for primary care.
748	Alzheimers Dement, 10(6), 656-665.e651. https://doi.org/10.1016/j.jalz.2013.11.006
749	Barnes, D. E., & Yaffe, K. (2011). The projected effect of risk factor reduction on
750	Alzheimer's disease prevalence. Lancet Neurol, 10(9), 819-828.
751	https://doi.org/10.1016/s1474-4422(11)70072-2
752	Bauer, D. J., & Curran, P. J. (2005). Probing Interactions in Fixed and Multilevel Regression:
753	Inferential and Graphical Techniques. Multivariate behavioral research, 40(3), 373-
754	400. <u>https://doi.org/10.1207/s15327906mbr4003_5</u>
755	Beliveau, V., Svarer, C., Frokjaer, V. G., Knudsen, G. M., Greve, D. N., & Fisher, P. M.
756	(2015). Functional connectivity of the dorsal and median raphe nuclei at rest.
757	NeuroImage, 116, 187-195. https://doi.org/10.1016/j.neuroimage.2015.04.065
758	Betts, M. J., Cardenas-Blanco, A., Kanowski, M., Jessen, F., & Düzel, E. (2017). In vivo
759	MRI assessment of the human locus coeruleus along its rostrocaudal extent in young
760	and older adults. NeuroImage, 163, 150-159.
761	https://doi.org/10.1016/j.neuroimage.2017.09.042
762	Betts, M. J., Cardenas-Blanco, A., Kanowski, M., Spottke, A., Teipel, S. J., Kilimann, I.,
763	Jessen, F., & Düzel, E. (2019). Locus coeruleus MRI contrast is reduced in
764	Alzheimer's disease dementia and correlates with CSF Aß levels. Alzheimers Dement
765	(Amst), 11, 281-285. https://doi.org/10.1016/j.dadm.2019.02.001
766	Betts, M. J., Kirilina, E., Otaduy, M. C. G., Ivanov, D., Acosta-Cabronero, J., Callaghan, M.
767	F., Lambert, C., Cardenas-Blanco, A., Pine, K., Passamonti, L., Loane, C., Keuken, M.
768	C., Trujillo, P., Lüsebrink, F., Mattern, H., Liu, K. Y., Priovoulos, N., Fliessbach, K.,

769	Dahl, M. J., Maaß, A., Madelung, C. F., Meder, D., Ehrenberg, A. J., Speck, O.,
770	Weiskopf, N., Dolan, R., Inglis, B., Tosun, D., Morawski, M., Zucca, F. A., Siebner,
771	H. R., Mather, M., Uludag, K., Heinsen, H., Poser, B. A., Howard, R., Zecca, L.,
772	Rowe, J. B., Grinberg, L. T., Jacobs, H. I. L., Düzel, E., & Hämmerer, D. (2019).
773	Locus coeruleus imaging as a biomarker for noradrenergic dysfunction in
774	neurodegenerative diseases. Brain, 142(9), 2558-2571.
775	https://doi.org/10.1093/brain/awz193
776	Berti, V., Mosconi, L., Glodzik, L., Li, Y., Murray, J., De Santi, S., Pupi, A., Tsui, W., & De
777	Leon, M. J. (2011). Structural brain changes in normal individuals with a maternal
778	history of Alzheimer's. Neurobiol Aging, 32(12), 2325.e2317-2326.
779	https://doi.org/10.1016/j.neurobiolaging.2011.01.001
780	Braak, H., & Braak, E. (1991). Neuropathological stageing of Alzheimer-related changes.
781	Acta Neuropathol, 82(4), 239-259. https://doi.org/10.1007/bf00308809
782	Braak, H., & Del Tredici, K. (2015). Neuroanatomy and pathology of sporadic Alzheimer's
783	disease. Adv Anat Embryol Cell Biol, 215, 1-162.
784	Braak, H., Thal, D. R., Ghebremedhin, E., & Del Tredici, K. (2011). Stages of the pathologic
785	process in Alzheimer disease: age categories from 1 to 100 years. J Neuropathol Exp
786	Neurol, 70(11), 960-969. https://doi.org/10.1097/NEN.0b013e318232a379
787	Buckley, R. F., O'Donnell, A., McGrath, E. R., Jacobs, H. I. L., Lois, C., Satizabal, C. L.,
788	Ghosh, S., Rubinstein, Z. B., Murabito, J. M., Sperling, R. A., Johnson, K. A.,
789	Seshadri, S., & Beiser, A. S. (2022). Menopause Status Moderates Sex Differences in
790	Tau Burden: A Framingham PET Study. Ann Neurol, 92(1), 11-22.
791	https://doi.org/10.1002/ana.26382

792	Cacciaglia, R., Operto, G., Falcón, C., de Echavarri-Gómez, J. M. G., Sánchez-Benavides, G.,
793	Brugulat-Serrat, A., Milà-Alomà, M., Blennow, K., Zetterberg, H., Molinuevo, J. L.,
794	Suárez-Calvet, M., & Gispert, J. D. (2022). Genotypic effects of APOE-ɛ4 on resting-
795	state connectivity in cognitively intact individuals support functional brain
796	compensation. Cereb Cortex. https://doi.org/10.1093/cercor/bhac239
797	Calarco, N., Cassidy, C. M., Selby, B., Hawco, C., Voineskos, A. N., Diniz, B. S., &
798	Nikolova, Y. S. (2022). Associations between locus coeruleus integrity and diagnosis,
799	age, and cognitive performance in older adults with and without late-life depression:
800	An exploratory study. NeuroImage: Clinical, 36, 103182.
801	https://doi.org/10.1016/j.nicl.2022.103182
802	Cnaan, A., Laird, N. M., & Slasor, P. (1997). Using the general linear mixed model to
803	analyse unbalanced repeated measures and longitudinal data. Statistics in medicine,
804	16(20), 2349-2380. https://doi.org/10.1002/(sici)1097-
805	0258(19971030)16:20<2349::aid-sim667>3.0.co;2-e
806	Cusack, R., Vicente-Grabovetsky, A., Mitchell, D. J., Wild, C. J., Auer, T., Linke, A. C., &
807	Peelle, J. E. (2015). Automatic analysis (aa): efficient neuroimaging workflows and
808	parallel processing using Matlab and XML. Front Neuroinform, 8, 90.
809	https://doi.org/10.3389/fninf.2014.00090
810	Clewett, D. V., Lee, T. H., Greening, S., Ponzio, A., Margalit, E., & Mather, M. (2016).
811	Neuromelanin marks the spot: identifying a locus coeruleus biomarker of cognitive
812	reserve in healthy aging. Neurobiol Aging, 37, 117-126.
813	https://doi.org/10.1016/j.neurobiolaging.2015.09.019
814	Dahl, M. J., Mather, M., Düzel, S., Bodammer, N. C., Lindenberger, U., Kühn, S., & Werkle-

815	Bergner, M. (2019). Rostral locus coeruleus integrity is associated with better
816	memory performance in older adults. Nat Hum Behav, 3(11), 1203-1214.
817	https://doi.org/10.1038/s41562-019-0715-2
818	Dahl, M. J., Mather, M., Werkle-Bergner, M., Kennedy, B. L., Guzman, S., Hurth, K., Miller,
819	C. A., Qiao, Y., Shi, Y., Chui, H. C., & Ringman, J. M. (2022). Locus coeruleus
820	integrity is related to tau burden and memory loss in autosomal-dominant Alzheimer's
821	disease. Neurobiol Aging, 112, 39-54.
822	https://doi.org/10.1016/j.neurobiolaging.2021.11.006
823	Deckers, K., van Boxtel, M. P., Schiepers, O. J., de Vugt, M., Muñoz Sánchez, J. L., Anstey,
824	K. J., Brayne, C., Dartigues, J. F., Engedal, K., Kivipelto, M., Ritchie, K., Starr, J. M.,
825	Yaffe, K., Irving, K., Verhey, F. R., & Köhler, S. (2015). Target risk factors for
826	dementia prevention: a systematic review and Delphi consensus study on the evidence
827	from observational studies. Int J Geriatr Psychiatry, 30(3), 234-246.
828	https://doi.org/10.1002/gps.4245
829	Del Cerro, I., Villarreal, M. F., Abulafia, C., Duarte-Abritta, B., Sánchez, S. M., Castro, M.
830	N., Bocaccio, H., Ferrer, I., Menchón, J. M., Sevlever, G., Nemeroff, C. B., Soriano-
831	Mas, C., & Guinjoan, S. M. (2020). Disrupted functional connectivity of the locus
832	coeruleus in healthy adults with parental history of Alzheimer's disease. J Psychiatr
833	Res, 123, 81-88. https://doi.org/10.1016/j.jpsychires.2020.01.018
834	Desikan, R. S., Ségonne, F., Fischl, B., Quinn, B. T., Dickerson, B. C., Blacker, D., Buckner,
835	R. L., Dale, A. M., Maguire, R. P., Hyman, B. T., Albert, M. S., & Killiany, R. J.
836	(2006). An automated labeling system for subdividing the human cerebral cortex on
837	MRI scans into gyral based regions of interest. NeuroImage, 31(3), 968-980.
838	https://doi.org/10.1016/j.neuroimage.2006.01.021

839	Donix, M., Small, G. W., & Bookheimer, S. Y. (2012). Family history and APOE-4 genetic
840	risk in Alzheimer's disease. Neuropsychol Rev, 22(3), 298-309.
841	https://doi.org/10.1007/s11065-012-9193-2
842	Dounavi, ME., Low, A., Mckiernan, E. F., Mak, E., Muniz-Terrera, G., Ritchie, K., Ritchie,
843	C. W., Su, L., & O'Brien, J. T. (2021). Evidence of cerebral hemodynamic
844	dysregulation in middle-aged APOE ɛ4 carriers: The PREVENT-Dementia study.
845	Journal of Cerebral Blood Flow & Metabolism, 41(11), 2844-2855.
846	https://doi.org/10.1177/0271678x211020863
847	Dounavi, ME., Newton, C., Jenkins, N., Mak, E., Low, A., Muniz-Terrera, G., Williams, G.
848	B., Lawlor, B., Naci, L., Malhotra, P., Mackay, C. E., Koychev, I., Ritchie, K.,
849	Ritchie, C. W., Su, L., & O'Brien, J. T. (2022). Macrostructural brain alterations at
850	midlife are connected to cardiovascular and not inherited risk of future dementia: the
851	PREVENT-Dementia study. Journal of Neurology. https://doi.org/10.1007/s00415-
852	<u>022-11061-7</u>
853	Dounavi, M. E., Mak, E., Wells, K., Ritchie, K., Ritchie, C. W., Su, L., & JT, O. B. (2020).
854	Volumetric alterations in the hippocampal subfields of subjects at increased risk of
855	dementia. Neurobiol Aging, 91, 36-44.
856	https://doi.org/10.1016/j.neurobiolaging.2020.03.006
857	Ehrenberg, A. J., Nguy, A. K., Theofilas, P., Dunlop, S., Suemoto, C. K., Di Lorenzo Alho, A.
858	T., Grinberg, L. T. (2017). Quantifying the accretion of hyperphosphorylated tau
859	in the locus coeruleus and dorsal raphe nucleus: the pathological building blocks of
860	early Alzheimer's disease. Neuropathol Appl Neurobiol, 43(5), 393-408.
861	https://doi.org/10.1111/nan.12387

862	Exalto, L. G., Quesenberry, C. P., Barnes, D., Kivipelto, M., Biessels, G. J., & Whitmer, R. A.
863	(2014). Midlife risk score for the prediction of dementia four decades later.
864	Alzheimer's & Dementia, 10(5), 562-570. https://doi.org/10.1016/j.jalz.2013.05.1772
865	Fayosse, A., Nguyen, D. P., Dugravot, A., Dumurgier, J., Tabak, A. G., Kivimäki, M., Sabia,
866	S., & Singh-Manoux, A. (2020). Risk prediction models for dementia: role of age and
867	cardiometabolic risk factors. BMC Med, 18(1), 107. https://doi.org/10.1186/s12916-
868	<u>020-01578-x</u>
869	Giannakopoulos, P., Herrmann, F. R., Bussière, T., Bouras, C., Kövari, E., Perl, D. P.,
870	Morrison, J. H., Gold, G., & Hof, P. R. (2003). Tangle and neuron numbers, but not
871	amyloid load, predict cognitive status in Alzheimer's disease. Neurology, 60(9), 1495-
872	1500. https://doi.org/10.1212/01.wnl.0000063311.58879.01
873	Gilvesy, A., Husen, E., Magloczky, Z., Mihaly, O., Hortobágyi, T., Kanatani, S., Adori, C.
874	(2022). Spatiotemporal characterization of cellular tau pathology in the human locus
875	coeruleus-pericoerulear complex by three-dimensional imaging. Acta Neuropathol.
876	https://doi.org/10.1007/s00401-022-02477-6
877	Hair Jr., J. F., Anderson, R. E., Tatham, R. L., & Black, W. C. (1998). Multivariate Data
878	Analysis (5th ed.). Prentice-Hall: London.
879	Heneghan, A., Deng, F., Wells, K., Ritchie, K., Muniz-Terrera, G., Ritchie, C. W., Lawlor, B.,
880	& Naci, L. (2023). Modifiable Lifestyle Activities Affect Cognition in Cognitively
881	Healthy Middle-Aged Individuals at Risk for Late-Life Alzheimer's Disease. Journal
882	of Alzheimer's disease : JAD, 91(2), 833-846. https://doi.org/10.3233/JAD-220267
883	Horn, J. (1965). A rationale and test for the number of factors in factor analysis.
884	Psychometrika, 30, 179-185. https://doi.org/10.1007/BF02289447

885	Hsieh, S., Schubert, S., Hoon, C., Mioshi, E., & Hodges, J. R. (2013). Validation of the
886	Addenbrooke's Cognitive Examination III in frontotemporal dementia and
887	Alzheimer's disease. Dement Geriatr Cogn Disord, 36(3-4), 242-250.
888	https://doi.org/10.1159/000351671
889	Irish, M., Lawlor, B. A., Coen, R. F., & O'Mara, S. M. (2011). Everyday episodic memory in
890	amnestic mild cognitive impairment: a preliminary investigation. BMC neuroscience,
891	12(1), 1-13. https://doi.org/10.1186/1471-2202-12-80
892	Irwin, K., Sexton, C., Daniel, T., Lawlor, B., & Naci, L. (2018). Healthy Aging and Dementia:
893	Two Roads Diverging in Midlife? Front Aging Neurosci, 10, 275.
894	https://doi.org/10.3389/fnagi.2018.00275
895	Jacobs, H. I., Wiese, S., van de Ven, V., Gronenschild, E. H., Verhey, F. R., & Matthews, P.
896	M. (2015). Relevance of parahippocampal-locus coeruleus connectivity to memory in
897	early dementia. Neurobiol Aging, 36(2), 618-626.
898	https://doi.org/10.1016/j.neurobiolaging.2014.10.041
899	Jacobs, H. I. L., Müller-Ehrenberg, L., Priovoulos, N., & Roebroeck, A. (2018). Curvilinear
900	locus coeruleus functional connectivity trajectories over the adult lifespan: a 7T MRI
901	study. Neurobiology of aging, 69, 167–176.
902	https://doi.org/10.1016/j.neurobiolaging.2018.05.021
903	Jacobs, H. I. L., Becker, J. A., Kwong, K., Engels-Domínguez, N., Prokopiou, P. C., Papp, K.
904	V., Properzi, M., Hampton, O. L., d'Oleire Uquillas, F., Sanchez, J. S., Rentz, D. M.,
905	El Fakhri, G., Normandin, M. D., Price, J. C., Bennett, D. A., Sperling, R. A., &
906	Johnson, K. A. (2021). In vivo and neuropathology data support locus coeruleus
907	integrity as indicator of Alzheimer's disease pathology and cognitive decline. Sci

908	Transl Med, 13(612), eabj2511. https://doi.org/10.1126/scitranslmed.abj2511
909	Jacobs, H. I. L., Becker, J. A., Kwong, K., Munera, D., Ramirez-Gomez, L., Engels-
910	Domínguez, N., Sanchez, J. S., Vila-Castelar, C., Baena, A., Sperling, R. A., Johnson,
911	K. A., Lopera, F., & Quiroz, Y. T. (2022). Waning locus coeruleus integrity precedes
912	cortical tau accrual in preclinical autosomal dominant Alzheimer's disease.
913	Alzheimers Dement. https://doi.org/10.1002/alz.12656
914	James, T., Kula, B., Choi, S., Khan, S. S., Bekar, L. K., & Smith, N. A. (2021). Locus
915	coeruleus in memory formation and Alzheimer's disease. European Journal of
916	Neuroscience, 54(8), 6948-6959. https://doi.org/10.1111/ejn.15045
917	Jansen, W. J., Ossenkoppele, R., Knol, D. L., Tijms, B. M., Scheltens, P., Verhey, F. R.,
918	Visser, P. J., Aalten, P., Aarsland, D., Alcolea, D., Alexander, M., Almdahl, I. S.,
919	Arnold, S. E., Baldeiras, I., Barthel, H., van Berckel, B. N., Bibeau, K., Blennow, K.,
920	Brooks, D. J., Zetterberg, H. (2015). Prevalence of cerebral amyloid pathology in
921	persons without dementia: a meta-analysis. JAMA, 313(19), 1924-1938.
922	https://doi.org/10.1001/jama.2015.4668
923	Johnson, P. O., & Fay, L. C. (1950). The Johnson-Neyman technique, its theory and
924	application. Psychometrika, 15(4), 349-367. <u>https://doi.org/10.1007/BF02288864</u>
925	Jolliffe, I. T., & Cadima, J. (2016). Principal component analysis: a review and recent
926	developments. Philos Trans A Math Phys Eng Sci, 374(2065), 20150202.
927	https://doi.org/10.1098/rsta.2015.0202.
928	Kaiser, H. F. (1958). The varimax criterion for analytic rotation in factor analysis.
929	Psychometrika, 23(3), 187-200. https://doi.org/10.1007/BF02289233

- 930 Kametani, F., & Hasegawa, M. (2018). Reconsideration of Amyloid Hypothesis and Tau
- 931 Hypothesis in Alzheimer's Disease. *Front Neurosci*, *12*, 25.
- 932 <u>https://doi.org/10.3389/fnins.2018.00025</u>
- 933 Kaufman, S. K., Del Tredici, K., Thomas, T. L., Braak, H., & Diamond, M. I. (2018). Tau
- 934 seeding activity begins in the transentorhinal/entorhinal regions and anticipates
- phospho-tau pathology in Alzheimer's disease and PART. Acta Neuropathol, 136(1),
- 936 57-67. <u>https://doi.org/10.1007/s00401-018-1855-6</u>
- 937 Kaufman, A. M., Geiller, T., & Losonczy, A. (2020). A Role for the Locus Coeruleus in
- Hippocampal CA1 Place Cell Reorganization during Spatial Reward Learning.
- 939 *Neuron*, *105*(6), 1018-1026.e1014. <u>https://doi.org/10.1016/j.neuron.2019.12.029</u>
- 940 Kempadoo, K. A., Mosharov, E. V., Choi, S. J., Sulzer, D., & Kandel, E. R. (2016).
- 941 Dopamine release from the locus coeruleus to the dorsal hippocampus promotes
- spatial learning and memory. *Proc Natl Acad Sci US A*, *113*(51), 14835-14840.
- 943 <u>https://doi.org/10.1073/pnas.1616515114</u>
- 944 Kerchner, G. A., Berdnik, D., Shen, J. C., Bernstein, J. D., Fenesy, M. C., Deutsch, G. K., ...

Rutt, B. K. (2014). APOE ε4 worsens hippocampal CA1 apical neuropil atrophy and
episodic memory. *Neurology*, *82*(8), 691-697.

- 947 https://doi.org/10.1212/wnl.00000000000154
- 948 Kivipelto, M., Ngandu, T., Laatikainen, T., Winblad, B., Soininen, H., & Tuomilehto, J.
- 949 (2006). Risk score for the prediction of dementia risk in 20 years among middle aged
- 950 people: a longitudinal, population-based study. *The Lancet Neurology*, *5*(9), 735-741.
- 951 https://doi.org/10.1016/s1474-4422(06)70537-3
- La Joie, R., Ayakta, N., Seeley, W. W., Borys, E., Boxer, A. L., DeCarli, C., Doré, V.,

953	Grinberg, L. T., Huang, E., Hwang, J. H., Ikonomovic, M. D., Jack, C., Jr., Jagust, W.
954	J., Jin, L. W., Klunk, W. E., Kofler, J., Lesman-Segev, O. H., Lockhart, S. N., Lowe,
955	V. J., Rabinovici, G. D. (2019). Multisite study of the relationships between
956	antemortem [(11)C]PIB-PET Centiloid values and postmortem measures of
957	Alzheimer's disease neuropathology. Alzheimers Dement, 15(2), 205-216.
958	https://doi.org/10.1016/j.jalz.2018.09.001
959	Lambert, J. C., Ibrahim-Verbaas, C. A., Harold, D., Naj, A. C., Sims, R., Bellenguez, C.,
960	DeStafano, A. L., Bis, J. C., Beecham, G. W., Grenier-Boley, B., Russo, G., Thorton-
961	Wells, T. A., Jones, N., Smith, A. V., Chouraki, V., Thomas, C., Ikram, M. A.,
962	Zelenika, D., Vardarajan, B. N., Amouyel, P. (2013). Meta-analysis of 74,046
963	individuals identifies 11 new susceptibility loci for Alzheimer's disease. Nat Genet,
964	45(12), 1452-1458. https://doi.org/10.1038/ng.2802
965	Lee, T. H., Kim, S. H., Katz, B., & Mather, M. (2020). The Decline in Intrinsic Connectivity
966	Between the Salience Network and Locus Coeruleus in Older Adults: Implications for
967	Distractibility. Front Aging Neurosci, 12, 2. https://doi.org/10.3389/fnagi.2020.00002
968	Liebe, T., Dordevic, M., Kaufmann, J., Avetisyan, A., Skalej, M., & Müller, N. (2022).
969	Investigation of the functional pathogenesis of mild cognitive impairment by
970	localisation-based locus coeruleus resting-state fMRI. Human brain mapping, 43(18),
971	5630-5642. https://doi.org/10.1002/hbm.26039
972	Liu, C. C., Liu, C. C., Kanekiyo, T., Xu, H., & Bu, G. (2013). Apolipoprotein E and
973	Alzheimer disease: risk, mechanisms and therapy. Nat Rev Neurol, 9(2), 106-118.
974	https://doi.org/10.1038/nrneurol.2012.263
975	Liu, K. Y., Acosta-Cabronero, J., Cardenas-Blanco, A., Loane, C., Berry, A. J., Betts, M. J.,

976	Kievit, R. A., Henson, R. N., Düzel, E., Howard, R., & Hämmerer, D. (2019). In vivo
977	visualization of age-related differences in the locus coeruleus. Neurobiol Aging, 74,
978	101-111. https://doi.org/10.1016/j.neurobiolaging.2018.10.014
979	Liu, K. Y., Kievit, R. A., Tsvetanov, K. A., Betts, M. J., Düzel, E., Rowe, J. B., Cam-CAN,
980	Howard, R., & Hämmerer, D. (2020). Noradrenergic-dependent functions are
981	associated with age-related locus coeruleus signal intensity differences. Nature
982	communications, 11(1), 1712. https://doi.org/10.1038/s41467-020-15410-w
983	Livingston, G., Huntley, J., Sommerlad, A., Ames, D., Ballard, C., Banerjee, S., Brayne, C.,
984	Burns, A., Cohen-Mansfield, J., Cooper, C., Costafreda, S. G., Dias, A., Fox, N.,
985	Gitlin, L. N., Howard, R., Kales, H. C., Kivimäki, M., Larson, E. B., Ogunniyi, A.,
986	Mukadam, N. (2020). Dementia prevention, intervention, and care: 2020 report of the
987	Lancet Commission. The Lancet, 396(10248), 413-446.
988	https://doi.org/10.1016/s0140-6736(20)30367-6
989	Livingston, G., Sommerlad, A., Orgeta, V., Costafreda, S. G., Huntley, J., Ames, D., Ballard,
990	C., Banerjee, S., Burns, A., Cohen-Mansfield, J., Cooper, C., Fox, N., Gitlin, L. N.,
991	Howard, R., Kales, H. C., Larson, E. B., Ritchie, K., Rockwood, K., Sampson, E.
992	L., Mukadam, N. (2017). Dementia prevention, intervention, and care. Lancet,
993	390(10113), 2673-2734. https://doi.org/10.1016/s0140-6736(17)31363-6
994	Lorenzo-Seva, U., & ten Berge, J. M. F. (2006). Tucker's congruence coefficient as a
995	meaningful index of factor similarity. Methodology: European Journal of Research
996	Methods for the Behavioral and Social Sciences, 2(2), 57-64.
997	https://doi.org/10.1027/1614-2241.2.2.57

998 Low, A., Su, L., Stefaniak, J. D., Mak, E., Dounavi, M. E., Muniz-Terrera, G., Ritchie, K.,

999	Ditahia	\mathbf{C} W	Morland	UС	& O'Drian	IT	(2021)	Inharitad	I rick of	dementia and
999	кисте		- WALKUS	по	$\alpha \cup \beta \cap e \cap$	JI		Innernec	LIISK OLO	летнениа апо

- 1000 the progression of cerebral small vessel disease and inflammatory markers in
- 1001 cognitively healthy midlife adults: the PREVENT-Dementia study. *Neurobiol Aging*,
- 1002 98, 124-133. <u>https://doi.org/10.1016/j.neurobiolaging.2020.10.029</u>
- 1003 Mather, M., & Harley, C. W. (2016). The Locus Coeruleus: Essential for Maintaining
- 1004 Cognitive Function and the Aging Brain. *Trends Cogn Sci*, 20(3), 214-226.
- 1005 https://doi.org/10.1016/j.tics.2016.01.001
- 1006 Mondadori, C. R. A., De Quervain, D. J.-F., Buchmann, A., Mustovic, H., Wollmer, M. A.,
- 1007 Schmidt, C. F., Boesiger, P., Hock, C., Nitsch, R. M., Papassotiropoulos, A., & Henke,
- 1008 K. (2007). Better Memory and Neural Efficiency in Young Apolipoprotein E 4
- 1009 Carriers. Cerebral Cortex, 17(8), 1934-1947. <u>https://doi.org/10.1093/cercor/bhl103</u>
- 1010 Mosconi, L., Berti, V., Quinn, C., McHugh, P., Petrongolo, G., Varsavsky, I., Osorio, R. S.,
- 1011 Pupi, A., Vallabhajosula, S., Isaacson, R. S., de Leon, M. J., & Brinton, R. D. (2017).
- 1012 Sex differences in Alzheimer risk: Brain imaging of endocrine vs chronologic aging.
- 1013 *Neurology*, 89(13), 1382-1390. <u>https://doi.org/10.1212/wnl.00000000004425</u>
- 1014 Mosconi, L., Rahman, A., Diaz, I., Wu, X., Scheyer, O., Hristov, H. W., Vallabhajosula, S.,
- 1015 Isaacson, R. S., de Leon, M. J., & Brinton, R. D. (2018). Increased Alzheimer's risk
- 1016 during the menopause transition: A 3-year longitudinal brain imaging study. *PLoS*
- 1017 One, 13(12), e0207885. <u>https://doi.org/10.1371/journal.pone.0207885</u>
- Murphy, K., Birn, R. M., Handwerker, D. A., Jones, T. B., & Bandettini, P. A. (2009). The
 impact of global signal regression on resting state correlations: are anti-correlated
 networks introduced? *NeuroImage*, 44(3), 893-905.
- 1021 https://doi.org/10.1016/j.neuroimage.2008.09.036

- 1022 Olivieri, P., Lagarde, J., Lehericy, S., Valabrègue, R., Michel, A., Macé, P., Caillé, F.,
- 1023 Gervais, P., Bottlaender, M., & Sarazin, M. (2019). Early alteration of the locus
- 1024 coeruleus in phenotypic variants of Alzheimer's disease. Ann Clin Transl Neurol, 6(7),
- 1025 1345-1351. https://doi.org/10.1002/acn3.50818
- 1026 Parra, M. A., Abrahams, S., Logie, R. H., Méndez, L. G., Lopera, F., & Della Sala, S. (2010).
- 1027 Visual short-term memory binding deficits in familial Alzheimer's disease. *Brain*,
- 1028 *133*(9), 2702-2713. <u>https://doi.org/10.1093/brain/awq148</u>
- 1029 Plini, E. R. G., O'Hanlon, E., Boyle, R., Sibilia, F., Rikhye, G., Kenney, J., Whelan, R.,
- 1030 Melnychuk, M. C., Robertson, I. H., & Dockree, P. M. (2021). Examining the Role of
- 1031 the Noradrenergic Locus Coeruleus for Predicting Attention and Brain Maintenance
- 1032 in Healthy Old Age and Disease: An MRI Structural Study for the Alzheimer's
- 1033 Disease Neuroimaging Initiative. *Cells*, 10(7), 1829.
- 1034 https://doi.org/10.3390/cells10071829
- 1035 Porat, S., Sibilia, F., Yoon, J., Shi, Y., Dahl, M. J., Werkle-Bergner, M., Düzel, S.,
- 1036 Bodammer, N., Lindenberger, U., Kühn, S., & Mather, M. (2022). Age differences in
- 1037 diffusivity in the locus coeruleus and its ascending noradrenergic tract. *NeuroImage*,
- 1038 251, 119022. https://doi.org/10.1016/j.neuroimage.2022.119022
- 1039 Power, J. D., Barnes, K. A., Snyder, A. Z., Schlaggar, B. L., & Petersen, S. E. (2012).
- 1040Spurious but systematic correlations in functional connectivity MRI networks arise
- 1041 from subject motion. *NeuroImage*, *59*(3), 2142-2154.
- 1042 <u>https://doi.org/10.1016/j.neuroimage.2011.10.018</u>
- 1043 Power, J. D., Mitra, A., Laumann, T. O., Snyder, A. Z., Schlaggar, B. L., & Petersen, S. E.
- 1044 (2014). Methods to detect, characterize, and remove motion artifact in resting state

1045	fMRI. NeuroImage, 84, 320-341. https://doi.org/10.1016/j.neuroimage.2013.08.048
1046	Prokopiou, P. C., Engels-Domínguez, N., Papp, K. V., Scott, M. R., Schultz, A. P., Schneider,
1047	C., Farrell, M. E., Buckley, R. F., Quiroz, Y. T., El Fakhri, G., Rentz, D. M., Sperling,
1048	R. A., Johnson, K. A., & Jacobs, H. I. L. (2022). Lower novelty-related locus
1049	coeruleus function is associated with $A\beta$ -related cognitive decline in clinically healthy
1050	individuals. Nature Communications, 13(1). https://doi.org/10.1038/s41467-022-
1051	<u>28986-2</u>
1052	Rahman, A., Schelbaum, E., Hoffman, K., Diaz, I., Hristov, H., Andrews, R., Jett, S., Jackson,
1053	H., Lee, A., Sarva, H., Pahlajani, S., Matthews, D., Dyke, J., de Leon, M. J., Isaacson,
1054	R. S., Brinton, R. D., & Mosconi, L. (2020). Sex-driven modifiers of Alzheimer risk:
1055	A multimodality brain imaging study. <i>Neurology</i> , 95(2), e166-e178.
1056	https://doi.org/10.1212/wnl.000000000009781
1057	Ritchie, C. W., & Ritchie, K. (2012). The PREVENT study: a prospective cohort study to
1058	identify mid-life biomarkers of late-onset Alzheimer's disease. BMJ Open, 2(6),
1059	e001893. https://doi.org/10.1136/bmjopen-2012-001893
1060	Ritchie, K., Carrière, I., Gregory, S., Watermeyer, T., Danso, S., Su, L., Ritchie, C. W., &
1061	O'Brien, J. T. (2021). Trauma and depressive symptomatology in middle-aged
1062	persons at high risk of dementia: the PREVENT Dementia Study. Journal of
1063	Neurology, Neurosurgery & Psychiatry, 92(1), 16-21. https://doi.org/10.1136/jnnp-
1064	2020-323823
1065	Ritchie, K., Carrière, I., Howett, D., Su, L., Hornberger, M., O'Brien, J. T., Ritchie, C. W., &
1066	Chan, D. (2018). Allocentric and Egocentric Spatial Processing in Middle-Aged

1067 Adults at High Risk of Late-Onset Alzheimer's Disease: The PREVENT Dementia

1068	Study. Journal of Alzheimer's Disease, 65(3), 885-896. https://doi.org/10.3233/jad-
1069	180432

- 1070 Ritchie, K., Carrière, I., Ritchie, C. W., Berr, C., Artero, S., & Ancelin, M. L. (2010).
- 1071 Designing prevention programmes to reduce incidence of dementia: prospective
- 1072 cohort study of modifiable risk factors. *Bmj*, *341*, c3885.
- 1073 <u>https://doi.org/10.1136/bmj.c3885</u>
- 1074 Ritchie, K., Carrière, I., Su, L., O'Brien, J. T., Lovestone, S., Wells, K., & Ritchie, C. W.
- 1075 (2017). The midlife cognitive profiles of adults at high risk of late-onset Alzheimer's
- 1076 disease: The PREVENT study. *Alzheimer's & Dementia*, 13(10), 1089-1097.
- 1077 <u>https://doi.org/10.1016/j.jalz.2017.02.008</u>
- 1078 Ritchie, K., de Roquefeuil, G., Ritchie, C., Besset, A., Poulain, V., Artero, S., Ancelin, M. L.

1079 (2014). COGNITO: computerized assessment of information processing. *Journal of* 1080 *Psychology & Psychotherapy*, 4(2). https://doi.org/10.4172/2161-0487.1000136

- 1081 Salvador, R., Martínez, A., Pomarol-Clotet, E., Gomar, J., Vila, F., Sarró, S., Capdevila, A.,
- 1082 & Bullmore, E. (2008). A simple view of the brain through a frequency-specific
 1083 functional connectivity measure. *NeuroImage*, *39*(1), 279-289.
- 1084 https://doi.org/10.1016/j.neuroimage.2007.08.018
- Sara, S. J. (2009). The locus coeruleus and noradrenergic modulation of cognition. *Nat Rev Neurosci*, 10(3), 211-223. https://doi.org/10.1038/nrn2573
- 1087 Satterthwaite, T. D., Elliott, M. A., Gerraty, R. T., Ruparel, K., Loughead, J., Calkins, M. E.,
- 1088 Eickhoff, S. B., Hakonarson, H., Gur, R. C., Gur, R. E., & Wolf, D. H. (2013). An
- 1089 improved framework for confound regression and filtering for control of motion
- 1090 artifact in the preprocessing of resting-state functional connectivity data. *NeuroImage*,

1091 *64*, 240-256. <u>https://doi.org/10.1016/j.neuroimage.2012.08.052</u>

- 1092 Scarabino, D., Gambina, G., Broggio, E., Pelliccia, F., & Corbo, R. M. (2016). Influence of
- 1093 family history of dementia in the development and progression of late-onset
- 1094 Alzheimer's disease. Am J Med Genet B Neuropsychiatr Genet, 171b(2), 250-256.
- 1095 https://doi.org/10.1002/ajmg.b.32399
- 1096 Scheyer, O., Rahman, A., Hristov, H., Berkowitz, C., Isaacson, R. S., Diaz Brinton, R., &

Mosconi, L. (2018). Female Sex and Alzheimer's Risk: The Menopause Connection. J
 Prev Alzheimers Dis, 5(4), 225-230. https://doi.org/10.14283/jpad.2018.34

1099 Sindi, S., Calov, E., Fokkens, J., Ngandu, T., Soininen, H., Tuomilehto, J., & Kivipelto, M.

1100 (2015). The CAIDE Dementia Risk Score App: The development of an evidence-

based mobile application to predict the risk of dementia. *Alzheimers Dement (Amst)*,

1102 *I*(3), 328-333. <u>https://doi.org/10.1016/j.dadm.2015.06.005</u>

1103 Song, I., Neal, J., & Lee, T. H. (2021). Age-Related Intrinsic Functional Connectivity

- 1104 Changes of Locus Coeruleus from Childhood to Older Adults. *Brain sciences*, 11(11),
- 1105 1485. https://doi.org/10.3390/brainsci11111485
- 1106 Sperling, R. A., Aisen, P. S., Beckett, L. A., Bennett, D. A., Craft, S., Fagan, A. M., Iwatsubo,

1107 T., Jack, C. R., Kaye, J., Montine, T. J., Park, D. C., Reiman, E. M., Rowe, C. C.,

1108 Siemers, E., Stern, Y., Yaffe, K., Carrillo, M. C., Thies, B., Morrison-Bogorad, M., ...

- 1109 Phelps, C. H. (2011). Toward defining the preclinical stages of Alzheimer's disease:
- 1110 Recommendations from the National Institute on Aging-Alzheimer's Association
- 1111 workgroups on diagnostic guidelines for Alzheimer's disease. *Alzheimer's* &
- 1112 Dementia, 7(3), 280-292. <u>https://doi.org/10.1016/j.jalz.2011.03.003</u>
- 1113 Stephen, R., Liu, Y., Ngandu, T., Antikainen, R., Hulkkonen, J., Koikkalainen, J.,

1114	Kemppainen, N., Lötjönen, J., Levälahti, E., Parkkola, R., Pippola, P., Rinne, J.,
1115	Strandberg, T., Tuomilehto, J., Vanninen, R., Kivipelto, M., Soininen, H., & Solomon,
1116	A. (2019). Brain volumes and cortical thickness on MRI in the Finnish Geriatric
1117	Intervention Study to Prevent Cognitive Impairment and Disability (FINGER).
1118	Alzheimers Res Ther, 11(1), 53. https://doi.org/10.1186/s13195-019-0506-z
1119	Stevens, J. P. (1992). Applied multivariate statistics for the social sciences (2nd ed.).
1120	Hillsdale, NJ: Erlbaum.
1121	Sun, W., Tang, Y., Qiao, Y., Ge, X., Mather, M., Ringman, J. M., & Shi, Y. (2020). A
1122	probabilistic atlas of locus coeruleus pathways to transentorhinal cortex for
1123	connectome imaging in Alzheimer's disease. NeuroImage, 223, 117301.
1124	https://doi.org/10.1016/j.neuroimage.2020.117301
1125	Takahashi, J., Shibata, T., Sasaki, M., Kudo, M., Yanezawa, H., Obara, S., Kudo, K., Ito, K.,
1126	Yamashita, F., & Terayama, Y. (2015). Detection of changes in the locus coeruleus in
1127	patients with mild cognitive impairment and Alzheimer's disease: high-resolution fast
1128	spin-echo T1-weighted imaging. Geriatr Gerontol Int, 15(3), 334-340.
1129	https://doi.org/10.1111/ggi.12280
1130	Tustison, N. J., Avants, B. B., Cook, P. A., Zheng, Y., Egan, A., Yushkevich, P. A., & Gee, J.
1131	C. (2010). N4ITK: improved N3 bias correction. IEEE transactions on medical
1132	imaging, 29(6), 1310-1320. https://doi.org/10.1109/TMI.2010.2046908
1133	Udeh-Momoh, C., & Watermeyer, T. (2021). Female specific risk factors for the
1134	development of Alzheimer's disease neuropathology and cognitive impairment: Call
1135	for a precision medicine approach. Ageing Res Rev, 71, 101459.
1136	https://doi.org/10.1016/j.arr.2021.101459

- 1137 Van Dijk, K. R., Hedden, T., Venkataraman, A., Evans, K. C., Lazar, S. W., & Buckner, R. L.
- 1138 (2010). Intrinsic functional connectivity as a tool for human connectomics: theory,
- properties, and optimization. *Journal of neurophysiology*, 103(1), 297–321.
- 1140 <u>https://doi.org/10.1152/jn.00783.2009</u>
- 1141 Van Egroo, M., van Hooren, R. W. E., & Jacobs, H. I. L. (2021). Associations between locus
- 1142 coeruleus integrity and nocturnal awakenings in the context of Alzheimer's disease
- plasma biomarkers: a 7T MRI study. *Alzheimers Res Ther*, 13(1), 159.
- 1144 <u>https://doi.org/10.1186/s13195-021-00902-8</u>
- 1145 Vilor-Tejedor, N., Evans, T. E., Adams, H. H., González-de-Echávarri, J. M., Molinuevo, J.
- 1146 L., Guigo, R., ... Operto, G. (2021). Genetic Influences on Hippocampal Subfields:
- 1147 An Emerging Area of Neuroscience Research. *Neurol Genet*, 7(3), e591.
- 1148 <u>https://doi.org/10.1212/nxg.000000000000591</u>
- 1149 Weber, M. T., Maki, P. M., & McDermott, M. P. (2014). Cognition and mood in
- 1150 perimenopause: a systematic review and meta-analysis. J Steroid Biochem Mol Biol,
- 1151 *142*, 90-98. <u>https://doi.org/10.1016/j.jsbmb.2013.06.001</u>
- 1152 Weinshenker, D. (2018). Long Road to Ruin: Noradrenergic Dysfunction in
- 1153 Neurodegenerative Disease. *Trends Neurosci*, 41(4), 211-223.
- 1154 <u>https://doi.org/10.1016/j.tins.2018.01.010</u>
- 1155 Williams GC. Pleiotropy, Natural Selection, and the Evolution of Senescence. Evolution.
- 1156 1957;11:398-411. https://doi.org/10.1111/j.1558-5646.1957.tb02911.x
- World Health Organization. (2021). Global status report on the public health response todementia.

- 1159 Yanes, L. L., & Reckelhoff, J. F. (2011). Postmenopausal hypertension. Am J Hypertens,
- 1160 24(7), 740-749. <u>https://doi.org/10.1038/ajh.2011.71</u>
- 1161 Ye, R., Rua, C., O'Callaghan, C., Jones, P. S., Hezemans, F. H., Kaalund, S. S., Tsvetanov, K.
- 1162 A., Rodgers, C. T., Williams, G., Passamonti, L., & Rowe, J. B. (2021). An in vivo
- probabilistic atlas of the human locus coeruleus at ultra-high field. *NeuroImage*, 225,
- 1164 117487. https://doi.org/10.1016/j.neuroimage.2020.117487
- 1165 Zhang, S., Hu, S., Chao, H. H., & Li, C.-S. R. (2016). Resting-state functional connectivity of
- the locus coeruleus in humans: in comparison with the ventral tegmental
- area/substantia nigra pars compacta and the effects of age. *Cerebral Cortex*, 26(8),
- 1168 3413-3427. https://doi.org/10.1093/cercor/bhv172
- 1169 Zhao, S., Rangaprakash, D., Venkataraman, A., Liang, P., & Deshpande, G. (2017).
- 1170 Investigating Focal Connectivity Deficits in Alzheimer's Disease Using Directional
- Brain Networks Derived from Resting-State fMRI. *Frontiers in aging neuroscience*, 9,
- 1172 211. https://doi.org/10.3389/fnagi.2017.00211
- 1173 Zokaei, N., Grogan, J., Fallon, S. J., Slavkova, E., Hadida, J., Manohar, S., . . . Husain, M.
- 1174 (2020). Short-term memory advantage for brief durations in human APOE ε4 carriers.
- 1175 Sci Rep, 10(1), 9503. <u>https://doi.org/10.1038/s41598-020-66114-6</u>
- 1176 Zuo, X. N., Di Martino, A., Kelly, C., Shehzad, Z. E., Gee, D. G., Klein, D. F., Castellanos, F.
- 1177 X., Biswal, B. B., & Milham, M. P. (2010). The oscillating brain: complex and
- reliable. *NeuroImage*, *49*(2), 1432-1445.
- 1179 <u>https://doi.org/10.1016/j.neuroimage.2009.09.037</u>

1180 Figure Legends

1181

1182	Figure 1. The flowchart of functional magnetic resonance imaging (fMRI) data processing. (a)
1183	The standard preprocessing pipeline for fMRI data. (b) Three exclusion criteria were used to
1184	exclude participants. A 'scrubbing' procedure was used to account for the effect of head
1185	movement on functional connectivity analyses (Power et al., 2012). (c) The top panels shows
1186	the study-specific 3mm locus coeruleus (LC) mask (in blue), which closely matched the 7T
1187	probabilistic atlas developed by Ye et al. (2021) and the LC meta-mask developed by Dahl et
1188	al. (2022). The bottom panel shows the study-specific mask superimposed on a participant's
1189	T1w structural data in MNI space. (d) Functional connectivity was calculated between the
1190	bilateral LC (in blue) and the hippocampus (in red). The hippocampus was defined after
1191	recon-all pipeline using Freesurfer version 7.1.0.
1192	

1193

Figure 2. Data reduction of cognitive information. (a) Scree plots and parallel analysis for the 1194 baseline and follow-up datasets. Eigenvalues of the principal components obtained from the 1195 actual data (in blue line) were compared to those of random simulated data (mean values in 1196 grey line and 95th percentile in red line). Eigenvalues larger than 95th percentile of the 1197 randomly generated 500 eigenvalues (simulation) determined the number of components. (b) 1198 1199 The three extracted components were rotated to be uncorrelated with each other. The proportion of variance explained by each component, and the cumulative variance explained 1200 by the three are presented for the baseline and follow-up datasets. Abbreviations: C1, 1201 1202 component 1; C2, component 2; C3, component 3. 1203

1205 Figure 3. Interpretation of data-derived cognitive domains. Component (C) coefficients and Tucker's congruence coefficients between (a) baseline and (b) follow-up, are shown. (a) The 1206 coefficients for 13 original cognitive measures (in rows) on the three components (in 1207 1208 columns). A larger absolute coefficient (darker colors and larger solid circles) represents a closer relationship between the cognitive measure and corresponding component. Cool/warm 1209 colors represent the positive/negative relationships between cognitive measures and 1210 1211 components. Only cognitive measures with a coefficient > 0.4 on each component, highlighted with purple rectangles, were used to interpret the cognitive functions driving each 1212 1213 component. (b) Similarities among the components across baseline (in rows) and follow-up 1214 (in columns) were measured by Tucker's congruence coefficients. The diagonal values 1215 represented the similarities for each of the components with itself across time, and off 1216 diagonal values indicated the similarities for each of the components with the other two 1217 components across time. Cool/warm colors represent the positive/negative relationships with darker colors representing correlation magnitude, as shown in the color-bar scale. 1218 1219 Abbreviations: VSTMBT, visual short-term memory binding test; diff, difference. 1220 1221 Figure 4. The associations between the CAIDE dementia risk score and verbal, visuospatial 1222 1223 functions, and short-term (conjunctive) memory at (a) baseline and (b) follow-up. 1224 Abbreviation: CAIDE, Cardiovascular risk factor, Aging, and Incidence of Dementia. 1225 1226 Figure 5. Longitudinal decline of the LC-Hippocampus functional connectivity over 2 years 1227 in cognitively healthy middle-aged adults. * = p < 0.05. Abbreviations: LC, Locus Coeruleus. 1228 1229 1230

1231 Figure 6. The CAIDE score significantly moderated the brain-behaviour relationship. The relationship was plotted at the fixed values of CAIDE (mean and ± 1 SD) to show the effect 1232 of CAIDE at (a) baseline and (b) follow-up. The LC–Hippocampus functional connectivity 1233 1234 and the verbal, visuospatial functions, and short-term (conjunctive) memory were positively associated for individuals with low CAIDE score, and negatively associated for those with 1235 high scores, at (b) follow-up, but not baseline (a). Johnson-Neyman plots show the specific 1236 range of CAIDE scores where FC was significantly associated with cognition at (c) baseline 1237 and (d) follow-up. The bolded horizontal line shows the range of CAIDE scores of our cohort. 1238 1239 Two blue dashed vertical lines indicate the interval of significant (blue area) and nonsignificant (pink area) brain-behaviour relationships. At follow-up, there was a significant 1240 1241 positive relationship between FC and cognition for individuals with a CAIDE score < 3.25 1242 and a significant negative relationship for individuals with a CAIDE score > 9.65. Abbreviations: LC, locus coeruleus; CAIDE, Cardiovascular risk factor, Aging, and 1243 Incidence of Dementia; FC, functional connectivity; SD, standard deviation; n.s., not 1244 1245 significant.

		Baselin	ne		Follow	z-up
	FHD- (n=107)	FHD+ (n=103)	<i>p</i> (Mann-Whitney U)	FHD- (n=89)	FHD+ (n=99)	<i>p</i> (Mann-Whitney U)
Age (y)	52.0 ± 11.0	53.0 ± 7.0	0.52	54.0 ± 10.0	55.0 ± 6.0	0.22
Years of Education	16.0 ± 5.0	16.0 ± 5.0	0.34	17.0 ± 5.0	16.0 ± 5.0	0.21
CAIDE (incl. APOE status)	5.0 ± 4.0 (n=107)	6.0 ± 2.0 (n=101)	<0.05	6.0 ± 4.0 (n=89)	7.0 ± 3.0 (n=93)	<0.005
			p (Chi-Square)			p (Chi-Square)
Sex (%female)	69.2%	71.8%	0.67	69.7%	71.7%	0.76
APOE ɛ4 (% Carriers)	28.0% (n=107)	44.6% (n=101)	<0.05	28.1% (n=89)	44.3% (n=97)	<0.05
	APOE ε4- (n=133)	APOE ε4+ (n=75)	<i>p</i> (Mann-Whitney U)	APOE ε4- (n=118)	APOE ε4+ (n=68)	<i>p</i> (Mann-Whitney U)
Age (y)	53.0 ± 8.0	52.0 ± 8.0	0.06	55.0 ± 8.0	54.5 ± 8.0	0.08
Years of Education	16.0 ± 5.0	17.0 ± 4.5	0.51	16.0 ± 4.8	17.0 ± 5.0	0.47
CAIDE (incl. APOE status)	5.0 ± 3.0 (n=133)	7.0 ± 3.0 (n=75)	<0.0005	6.0 ± 3.0 (n=118)	7.0 ± 4.0 (n=64)	<0.0005
CAIDE (excl. APOE status)	5.0 ± 2.0 (n=133)	4.0 ± 3.0 (n=75)	0.12	5.0 ± 2.0 (n=118)	4.5 ± 3.0 (n=64)	0.25
			p (Chi-Square)			p (Chi-Square)
Sex (%female)	69.9%	70.7%	0.91	71.2%	69.1%	0.77

1250

- 1251 Note: median ± interquartile range (IQR) was reported for continuous variables. Abbreviations: FHD-, Negative family history of dementia;
- 1252 FHD+, positive family history of dementia; APOE ε4+, Apolipoprotein ε4 genotype positive; APOE ε4-, Apolipoprotein ε4 genotype negative;
- 1253 CAIDE, Cardiovascular Risk Factors, Aging and Dementia.

Table 2. Statistical summary of the multiple regression model testing the relationship

			Deselie			Eallary w	
			Baselin	e		Follow-u	p
Model s	ummara	R^2	F (4, 201)	р	R^2	F (4, 169)	р
	Model summary		9.55	< 0.0001	0.17	8.48	< 0.0001
DV	IV	β	se	р	β	se	р
	APOE ε4	0.28	0.13	< 0.05	0.27	0.15	0.07
Episodic and	Age	-0.02	0.01	0.17	-0.008	0.01	0.55
relational	Sex	0.27	0.14	0.06	0.25	0.16	0.11
memory	Years of education	0.09	0.02	< 0.0001	0.10	0.02	< 0.0001

between APOE and episodic and relational memory

Note: unstandardized coefficient (β) and standard error (se) are reported. DV, dependent variable; IV, independent variable; APOE ϵ 4, Apolipoprotein ϵ 4 genotype.

1255

	_	Baseline			Follow-up		
Model summary –		R^2	F (4, 120)	р	R^2	F (4, 96)	р
		0.07	2.34	0.06	0.06	1.45	0.22
DV	IV	β	se	р	β	se	р
Verbal, visuospatial	CAIDE	0.09	0.06	0.18	0.07	0.07	0.35
functions, and short-	Menopause	1.12	0.58	0.05	1.81	0.87	< 0.05
term (conjunctive)	CAIDE*Menopause	-0.20	0.09	< 0.05	-0.29	0.13	< 0.05
memory	Age	-0.04	0.03	0.10	-0.02	0.03	0.52

1257 Table 3. Statistical summary of the multiple regression model testing the interaction between CAIDE and menopausal status

1258

1259 Note: unstandardized coefficient (β) and standard error (se) are reported. DV, dependent variable; IV, independent variable; APOE ϵ 4,

1260 Apolipoprotein ε4 genotype.

1262

Table 4. Statistical summary of longitudinal linear mixed effect model testing for longitudinal change in the LC–Hippocampus functional connectivity

DV	IV	β	se	р	
	Time	-0.06	0.03	< 0.05	
LC–Hippocampus connectivity	mean FD	0.34	1.42	0.81	
	Hippocampus volume	0.01	0.06	0.87	

Note: unstandardized coefficient (β) and standard error (se) are reported. DV, dependent variable; IV, independent variable; FD, framewise displacement; LC, Locus Coeruleus

1264


Table 5. Regression coefficients of the model testing the relationship between CAIDE, LC-

Hippocampus and verbal, visuospatial functions, and short-term (conjunctive) memory


		Baseline			Follow-up		
Model summary		R^2	F (4, 121)	р	R^2	F (4, 77)	р
		0.07	2.22	0.07	0.07	1.55	0.20
DV	IV	β	se	р	β	se	р
Verbal,	CAIDE	-0.09	0.03	< 0.01	-0.04	0.05	0.45
visuospatial	LC-HC	-0.10	0.34	0.78	0.48	0.53	0.37
functions, and short- term	CAIDE × LC–HC	-0.10	0.14	0.48	-0.54	0.22	< 0.05
(conjunctive) memory	Hippocampus volume	-1.97E-04	2.29E-04	0.39	5.39E-06	3.26E-04	0.99

Note: unstandardized coefficient (β) and standard error (se) are reported. CAIDE and LC-HC have been centred in this model. DV, dependent variable; IV, independent variable; CAIDE, Cardiovascular Risk Factors, Aging and Dementia; LC–HC, Locus Coeruleus–Hippocampus functional connectivity.

(a) Resting-state fMRI preprocessing









- Decline or contraindications to MRI
- Incidental findings

Motion correction Slice-time correction Co-register to T1 Normalization Smooth Regress out covariates Bandpass filtering


(C) Create 3mm LC mask




X = 37

















- - Simulated data (mean)

Simulated data (95th percentile)

- Description recall (accu
 - Narrative recall (accu
- Name-face association (accu
 - Verbal fluency(accu
 - Working memory (latency
 - Working memory (accu
 - VSTMBT shape (accu
- Phoneme comprehension (accu
 - Form perception (late
- nedRxiv preprint doi: https://doi.org/10.1101/2022.09.02.22279541; this version posted February 22, 2024. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in Phoneme comprehension (late
- VSTMBT shape-colour binding (accu
 - Form perception (accu
 - Implicit memory (accu


	C1	C2	C 3
uracy)	0.82	-0.03	0.01
uracy)	0.78	-0.02	0.15
Jracy)	0.55	-0.02	-0.24
uracy)	0.41	0.33	-0.18
y diff)	0.2	-0.74	-0.02
uracy)	0.08	0.67	0.05
uracy)	0.4	0.48	-0.21
uracy)	0.08	0.41	0.11
tency)	0.13	-0.01	0.78
tency)	-0.12	0.09	0.61
Jracy)	0.26	0.25	-0.46
uracy)	0.33	0.29	0.39
uracy)	-0.07	0.15	0.06

C1	C2
0.74	-0.0
0.71	0.1
0.54	-0.1
0.45	-0.0
-0.22	0.4
0.23	0.4
0.56	0.3
0.13	0.4
0.02	0.1
-0.05	-0.1
0.41	0.4
-0.04	0.7
-0.22	0.2


Baseline

Follow-up

		(D)											
;2	C 3												
04	-0.05			Follow-up									
.1	0.14		C1	C2	C3		-0.8						
14	-0.21						-0.6						
05	-0.24	5	0.90	0.35	-0.07		-0.4						
41	-0.1						-0.2						
.4	-0.14	E E			$\sim \sim \sim$		-0						
39	0.02	G gg gg gg gg gg gg gg gg gg gg gg gg gg	0.41	0.40	0.02		-0.2						
41	0.27						-0.4						
19	0.7	B	-0.23	0.11	0.76		0.6						
12	0.78						0.8						
	0.02						1						
.7	0		C1: Episodic and relational memory C2: Working and short-term (single-feature) memory										
.2	-0.36	C3: Verba	al, visuospat short-term (c	ial functions	,								



rho = -0.17, *p* < 0.05 10 CAIDE score




0.40**()** O Ē 0.00-_ -0.40-

-0.80-


medRxiv preprint doi: https://doi.org/10.1101/2022.09.02.22279541; this version posted February 22, 2024. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. All rights reserved. No reuse allowed without permission.

Baseline











Baseline

Interaction:







Follow-up

Interaction:

