

## Appendix A

Table A1 – Socio-demographic, clinical and daily-life activity characteristics of DIALOAD participants aged ≥55 and <55 years

Socio-demographic and clinical characteristics	DIALOAD ≥55 years (n=50)	DIALOAD <55 years (n=8)	p-value
Age (years)	67.7 (SD 7.3)	49.8 (SD 3.7)	<0.001***
Female sex	20 (10)	13 (1)	0.615
Height (cm)	182.3 (SD 10.5)	182.4 (SD 5.0)	0.993
BMI (kg/m <sup>2</sup> )	29.6 (SD 5.4)	30.2 (SD 6.3)	0.809
Type 2 diabetes	88 (44)	38 (3)	<0.001***
Presence of peripheral neuropathy	100 (50)	100 (8)	1.000
Employed	34 (17)	100 (8)	<0.001***
Living alone	48 (24)	25 (2)	0.225
Education level			0.209
- Low	26 (13)	25 (2)	
- Moderate	22 (11)	50 (4)	
- High	52 (26)	25 (2)	
Short Physical Performance Battery score (0-12)	9.0 [IQ 7.0; 10.0] <sup>(n=38)</sup>	12.0 [IQ 11.0; 12.0] <sup>(n=6)</sup>	<0.001***
<b>Daily-life activity characteristics</b>	<b>(n=47)</b>	<b>(n=8)</b>	
Walking			
- Number of steps per day	5213 (SD 3404)	9292 (SD 3165)	0.003**
- Number of episodes per day	349 (SD 182)	560 (SD 561)	0.006**
- Duration per day (hours)	1.1 (SD 0.6)	1.8 (SD 0.6)	0.002**
Standing			
- Number of episodes per day	744 (SD 345)	1201 (SD 501)	0.002**
- Duration per day (hours)	1.8 (SD 0.7)	2.9 (SD 0.8)	<0.001***
Sitting			
- Number of episodes per day	91 (SD 25)	135 (SD 73)	0.127
- Duration per day (hours)	9.4 (SD 2.7)	7.3 (SD 1.5)	0.037*
Lying			
- Number of episodes per day	11 (SD 5)	12 (SD 7)	0.679
- Duration per day (hours)	9.7 (SD 2.8)	8.8 (SD 3.4)	0.384
Total wearing time per day (hours)	22.3 (SD 2.4)	21.4 (SD 3.5)	0.295

Note: Continuous data are mean (SD standard deviation) if normally distributed and median [IQ 25th percentile; 75<sup>th</sup> percentile] if not normally distributed, and discrete data are percentage (number of). Significant differences are depicted as \*\*\* =  $p < 0.001$ , \*\* =  $p < 0.01$  and \* =  $p < 0.05$ . If data was missing for a characteristic, the number of available data is shown in superscript.