

Appendix

Table A1. All adult hospitalisation spells* broken down by whether they were unplanned hospitalisations due to ACSC, by ID status, 2011-19 (%)

	ACSC hospitalisation		Total	N
	No	Yes		
No ID	94.8	5.2	100	579,917
ID	94.8	5.2	100	17,276

* A continuous inpatient (CIP) spell is a continuous period of care within the NHS, regardless of any transfers which may take place.

Table A2. Incidence rates of adult hospitalisations due to an ACSC per 1,000 person-years, by ID status and ACSC, 2011-19 (sorted by decreasing SIR in ID group)

ACSC	Crude incidence rates				ID incidence rates age-sex standardized to 'No ID' population	
	No ID		ID		SIR	SIR 95% CI
	IR	IR 95% CI	IR	IR 95% CI		
Any ACSC	6.07	(6.00, 6.14)	6.10	(5.71, 6.51)	5.94	(5.56, 6.35)
Convulsions and Epilepsy	1.17	(1.14, 1.20)	1.33	(1.15, 1.53)	1.36	(1.18, 1.57)
COPD	1.17	(1.14, 1.20)	1.23	(1.06, 1.43)	1.21	(1.04, 1.40)
Influenza and Pneumonia	0.90	(0.87, 0.92)	0.99	(0.84, 1.17)	0.98	(0.83, 1.15)
Cellulitis	0.50	(0.48, 0.52)	0.41	(0.32, 0.53)	0.43	(0.33, 0.55)
Asthma	0.41	(0.39, 0.43)	0.42	(0.33, 0.55)	0.39	(0.30, 0.51)
Ear, nose, and throat infections	0.37	(0.35, 0.39)	0.39	(0.30, 0.51)	0.37	(0.29, 0.49)

Congestive Heart Failure	0.43	(0.41, 0.45)	0.39	(0.30, 0.51)	0.36	(0.27, 0.47)
Diabetes complications	0.44	(0.42, 0.46)	0.29	(0.22, 0.40)	0.26	(0.19, 0.36)
Dehydration and Gastroenteritis	0.24	(0.22, 0.25)	0.27	(0.20, 0.37)	0.25	(0.18, 0.35)
Angina	0.26	(0.24, 0.27)	0.28	(0.20, 0.38)	0.24	(0.17, 0.34)
Pyelonephritis	0.11	(0.10, 0.12)	0.13u	(0.08, 0.21)u	0.13u	(0.08, 0.20)u
Iron deficiency anaemia	0.10	(0.09, 0.11)	0.08u	(0.05, 0.15)u	0.07u	(0.04, 0.13)u
Perforated/bleeding ulcer	0.04	(0.03, 0.04)	0.05u	(0.02, 0.10)u	0.06u	(0.03, 0.11)u
Gangrene	0.07	(0.07, 0.08)	0.05u	(0.02, 0.10)u	0.04u	(0.02, 0.10)u
Hypertension	0.05	(0.04, 0.06)	0.03u	(0.01, 0.08)u	0.04u	(0.02, 0.09)u
Dental conditions	0.06	(0.05, 0.07)	0.03u	(0.01, 0.08)u	0.03u	(0.01, 0.08)u
Pelvic Inflammatory Disease	0.03	(0.02, 0.03)				
Other Vaccine Preventable	0.01	(0.01, 0.02)				
Nutritional deficiencies						

Note: letter 'u' indicates that the figure may not be reliable due to the small number of cases (between 5-20).

Table A3. Average length of unplanned hospital stay due to ACSC, by ID status and ACSC, 2011-19 (measured in days; stays of adults on Census 2011 day; length refers to hospital spells. Sorted by decreasing mean length of stay in the ID group)

ACSC	No ID			ID		
	Number of spells	median	mean	Number of spells	median	mean
Any ACSC	30690*	1.0	2.5	924	1.0	2.6

Dental conditions	298	1.0	1.7	5	1.0	13.0
Gangrene	351	3.0	7.8	7	4.0	4.7
Influenza and Pneumonia	4342	1.0	3.2	143	1.0	3.8
Cellulitis	2395	1.0	3.1	59	1.0	3.7
Perforated/bleeding ulcer	187	1.0	2.4	7	2.0	3.4
Diabetes complications	2133	1.0	3.4	42	1.0	3.3
COPD	5663	1.0	2.5	177	1.0	2.9
Congestive Heart Failure	2061	1.0	3.3	56	1.0	2.4
Dehydration and Gastroenteritis	1147	1.0	3.2	39	1.0	2.1
Pyelonephritis	521	2.0	2.3	19	1.0	1.9
Convulsions and Epilepsy	5669	1.0	1.7	191	1.0	1.8
Angina	1245	1.0	1.3	40	1.0	1.6
Ear, nose, and throat infections	1794	1.0	1.2	56	1.0	1.4
Asthma	1986	1.0	1.4	61	1.0	1.2
Iron deficiency anaemia	477	1.0	1.9	12	0.0	0.8
Hypertension	238	1.0	1.4	5	0.0	0.4
Other Vaccine Preventable	57	1.0	2.2	<5		
Pelvic Inflammatory Disease	126	2.0	3.1	<5		
Nutritional deficiencies	<5			<5		

* rounded to the nearest 5 to avoid disclosure on the number of Nutritional Deficiencies spells.

Table A4. Adult ACSC mortality rates per 1,000 person-years, by ID status and 'main-cause of death' status, 2011-19 (sorted by decreasing main-cause SMR in ID group)

ACSC	Main-cause						All-cause					
	Crude mortality rate				ID mortality rate age-sex standardized to 'No ID' population		Crude mortality rate				ID mortality rate age-sex standardized to 'No ID' population	
	No ID		ID				No ID		ID			
	CMR	CMR 95% CI	CMR	CMR 95% CI	SMR	SMR 95% CI	CMR	CMR 95% CI	CMR	CMR 95% CI	SMR	SMR 95% CI
Any ACSC	1.03	(1.00, 1.06)	1.67	(1.47, 1.90)	2.62	(2.36, 2.91)	3.90	(3.84, 3.95)	5.66	(5.27, 6.08)	8.79	(8.30, 9.30)
COPD	0.80	(0.78, 0.83)	0.57	(0.46, 0.71)	1.06	(0.90, 1.25)	1.73	(1.70, 1.77)	1.16	(1.00, 1.36)	2.12	(1.88, 2.38)
Convulsions and Epilepsy	0.02	(0.02, 0.02)	0.69	(0.56, 0.85)	0.79	(0.65, 0.95)	0.10	(0.09, 0.11)	2.36	(2.11, 2.63)	2.76	(2.49, 3.06)
Asthma	0.03	(0.02, 0.03)	0.10 _u	(0.06, 0.17)	0.17 _u	(0.12, 0.26)	0.09	(0.08, 0.10)	0.23	(0.16, 0.33)	0.39	(0.30, 0.52)
Hypertension	0.03	(0.03, 0.04)	0.06 _u	(0.03, 0.12)	0.14 _u	(0.09, 0.22)	1.15	(1.12, 1.18)	0.76	(0.63, 0.93)	1.56	(1.36, 1.78)
Congestive Heart Failure	0.08	(0.07, 0.09)	0.07 _u	(0.04, 0.14)	0.13 _u	(0.09, 0.21)	1.15	(1.12, 1.18)	1.25	(1.08, 1.46)	2.18	(1.94, 2.44)
Influenza and Pneumonia	0.03	(0.02, 0.03)	0.07 _u	(0.04, 0.14)	0.12 _u	(0.07, 0.20)	0.04	(0.03, 0.04)	0.08 _u	(0.05, 0.15)	0.13 _u	(0.08, 0.20)
Cellulitis	0.02	(0.02, 0.03)	0.04 _u	(0.02, 0.09)	0.08 _u	(0.04, 0.15)	0.05	(0.04, 0.05)	0.09 _u	(0.05, 0.16)	0.16 _u	(0.10, 0.24)
Pyelonephritis	0.01	(0.00, 0.01)	0.04 _u	(0.02, 0.09)	0.06 _u	(0.03, 0.13)	0.01	(0.01, 0.01)	0.08 _u	(0.05, 0.15)	0.12 _u	(0.07, 0.20)
Dehydration and Gastroenteritis	0.01	(0.00, 0.01)					0.03	(0.03, 0.04)	0.12 _u	(0.07, 0.19)	0.27 _u	(0.20, 0.38)
Angina	0.00 _u	(0.00, 0.00)					0.03	(0.02, 0.03)				
Gangrene							0.02	(0.01, 0.02)	0.04 _u	(0.02, 0.09)	0.06 _u	(0.03, 0.13)
Pelvic Inflammatory Disease							0.00 _u	(0.00, 0.00)				
Nutritional Deficiencies							0.00 _u	(0.00, 0.00)				
Ear, nose and throat infections							0.00 _u	(0.00, 0.00)				
Dental conditions							0.00 _u	(0.00, 0.00)				
Iron deficiency anaemia												
Other Vaccine Preventable												
Perforated/bleeding ulcer												
Diabetes Complications												

Note: letter 'u' indicates that the figure may not be reliable due to the small number of cases (between 5-20).

