

Clinical value of plasma ALZpath pTau217 immunoassay in the assessment of mild cognitive impairment

Sylvain Lehmann, MD^{1*}, Susanna Schraen-Maschke, PhD², Jean-Sébastien Vidal, MD³, Constance Delaby, PhD^{1,4}, Luc Buée PhD², Frédéric Blanc, MD⁵, Claire Paquet, MD⁶, Bernadette Allinquant, PhD⁷, Stéphanie Bombois, MD^{2,8}, Audrey Gabelle, MD⁹, Olivier Hanon, MD^{3*} on behalf of the BALTAZAR study group.

Supplementary data

Page 2: Supplementary Table 1: Distribution of plasma pTau217, plasma pTau181 and CSF pTau181 in A β - and A β + population

Page 3: Supplementary Table 2. Biological values in MCI population.

Page 4: Supplementary Table 3: Risk factors associated with conversion to dementia during follow up

Page 5: Supplementary Table 4: Correlation between plasma pTau181, pTau217 and clinical chemistry analytes

Page 5: Supplementary Figure 1: Correlation between plasma pTau217, plasma pTau181 and hippocampal volume.

Page 6: Supplementary Figure 2: Correlation between plasma pTau217 and pTau181 and CSF pTau181 in MCI according to amyloid status.

Page 7: Supplementary Figure 3: Distribution and ROC curves of plasma pTau217 and pTau181 in MCI according to conversion to AD.

Supplementary Table 1: Distribution of plasma pTau217, plasma pTau181 and CSF pTau181 in A β - and A β + population

Factors	n A β -	Value (Mean(sd))	n A β +	Value (Mean(sd))	A β - vs A β + (p)	Fold change
Plasma pTau217 (pg/mL)	97	0.28 (0.19)	116	0.75 (0.34)	<.0001	2.67
Plasma pTau 181 (pg/mL)	97	2.60 (1.42)	116	3.87 (1.38)	<.0001	1.48
CSF pTau181 (pg/mL)	97	51.9 (16.0)	116	79.1 (32.3)	<.0001	1.52

Values as mean \pm standard deviation in the number (n) of MCI participants in A β - and A β + population with t-student comparison test

Supplementary Table 2. Biological values in MCI population.

	Total n MCI	Value (Mean(sd))	A β -	Value (Mean(sd))	A β +	Value (Mean(sd))	A β - vs A β +(p)	Adjusted age, sex, APOE ϵ 4, education	n MCI Non Converter	Value (Mean(sd))	n MCI Converter	Value (Mean(sd))	Converter vs non converter (p)	Adjusted age, sex, APOE ϵ 4, education
Adiponectin (ug/mL)	472	8.86 (6.22)	97	7.05 (4.25)	116	8.59 (5.92)	0.0279	0.1331	337	8.94 (6.49)	135	8.64 (5.49)	0.6037	0.3337
Fasting glycemia (mmol/L)	439	5.37 (1.19)	90	5.42 (1.15)	114	5.42 (1.16)	0.9977	0.6896	311	5.33 (1.2)	128	5.45 (1.17)	0.3300	0.2035
Triglycerides (mmol/L)	421	5.5 (1.16)	87	5.54 (1.26)	107	5.5 (1.12)	0.8158	0.2670	297	1.22 (0.6)	124	1.22 (0.55)	0.9890	0.8715
Cholesterol total (mmol/L)	419	1.74 (0.52)	87	1.76 (0.49)	108	1.71 (0.51)	0.5282	0.7774	297	5.46 (1.2)	124	5.57 (1.06)	0.3540	0.4530
Cholesterol HDL (mmol/L)	417	3.2 (1)	87	3.25 (1.07)	107	3.25 (0.92)	0.9933	0.1931	296	1.74 (0.53)	123	1.73 (0.48)	0.8324	0.6706
Cholesterol LDL (mmol/L)	421	1.22 (0.59)	87	1.18 (0.48)	108	1.15 (0.69)	0.7116	0.6172	295	3.17 (1.02)	122	3.28 (0.96)	0.2794	0.4081
Creatinine (μ mol/L)	443	78.21 (20.97)	92	80.15 (24.88)	113	78.31 (20.02)	0.5657	0.7440	316	78.36 (20.47)	127	77.84 (22.25)	0.8216	0.4137
eGFR (mL/min/1.73m ²)	444	76.99 (14.7)	92	77.51 (15.55)	114	76.69 (15.34)	0.7061	0.5876	317	76.81 (14.94)	127	77.44 (14.14)	0.6766	0.4420
Prealbumin (mg/dl)	385	0.279 (0.061)	84	0.285 (0.077)	99	0.282 (0.05)	0.7025	0.5216	268	0.28 (0.06)	117	0.28 (0.06)	0.4165	0.3961
Albumin (g/L)	442	40.32 (3.9)	93	39.89 (3.59)	110	40.19 (4.9)	0.6183	0.3942	318	40.52 (3.53)	124	39.81 (4.68)	0.1279	0.1349
Total protein (g/L)	430	70.46 (6.67)	90	70.69 (5.69)	111	70.31 (4.1)	0.5889	0.7118	305	70.33 (7.41)	125	70.75 (4.38)	0.4715	0.4367
CRP (mg/L)	438	3.21 (5.39)	90	4.76 (8.71)	111	2.28 (2.7)	0.0107	0.0120	314	3.05 (4.59)	124	3.63 (7.01)	0.3959	0.1054
Hemoglobin (g/dL)	439	13.8 (1.15)	91	13.76 (1.21)	112	13.84 (0.95)	0.5831	0.2239	312	13.75 (1.19)	127	13.92 (1.04)	0.1594	0.2466
Vitamin B12 (pmol/L)	408	310.9 (131.7)	86	322.4 (156.2)	101	307.5 (108.3)	0.4559	0.2743	291	315.27 (137.18)	101	300 (116.79)	0.2576	0.4200
TSH (mU/L)	391	2.31 (2.69)	83	2.29 (2.73)	101	2.39 (3)	0.8137	0.6952	274	2.25 (2.22)	101	2.43 (3.57)	0.6139	0.6643
Folate (nmol/L)	216	29.45 (47.32)	32	40.44 (71.58)	42	21.69 (24.3)	0.1640	0.0238	156	30.03 (44.11)	42	27.97 (55.17)	0.7963	0.9238
Red-cell folate (nmol/L)	343	1239 (726)	78	1404 (1015)	88	1328 (656)	0.5722	0.9222	240	1225 (737)	88	1271 (701)	0.5796	0.3599

Values in the number (n) of MCI participants for which different data types were available and comparison between non-converter and converters, with t-student or χ^2 and linear regression adjusted for age, sex, and the presence of the APOE ϵ 4 allele; numbers were used to describe categorical variables. mean \pm standard deviation for continuous variables. After Bonferroni adjustment, the significant P value is set at 0.003. Abbreviations: MCI: mild cognitive impairment; eGFR, estimated glomerular filtration rate; sd, standard deviation.

Supplementary Table 3: Risk factors associated with conversion to dementia during follow up

Factors	n	HR Conversion (95%CI)	p	p adjusted (age, sex, ApoE)
Adiponectin	472	0.96 (1.02-1.02)	0.5870	0.2982
Fasting glycemia	439	0.97 (1.27-1.27)	0.1208	0.0540
Triglycerides	421	0.88 (1.64-1.64)	0.2389	0.1680
Cholesterol total	421	0.92 (1.24-1.24)	0.3886	0.1671
Cholesterol HDL	419	0.59 (1.17-1.17)	0.2824	0.8413
Cholesterol LDL	417	0.95 (1.34-1.34)	0.1835	0.1404
Creatinine	443	0.99 (1.01-1.01)	0.7844	0.5356
eGFR	444	0.99 (1.01-1.01)	0.9811	0.7385
Prealbumin	385	0.01 (5.35-5.35)	0.3690	0.4187
Albumin	442	0.90 (0.99-0.99)	0.0193	0.0272
Total protein	430	0.98 (1.04-1.04)	0.7511	0.6173
CRP	438	0.99 (1.05-1.05)	0.2197	0.0452
Hemoglobin	439	0.97 (1.31-1.31)	0.1140	0.1336
Vitamin B12	408	1.00 (1.00-1.00)	0.2269	0.4278
TSH	391	0.97 (1.10-1.10)	0.3234	0.637
Folate	216	0.99 (1.00-1.00)	0.4958	0.8443
Red-cell folate	343	1.00 (1.00-1.00)	0.9336	0.5171



Cox proportional hazard model of conversion to dementia in follow-up before and after adjustment for age, sex, educational level, and the APOE $\epsilon 4$ status. Abbreviations: HR, hazard ratio for conversion; CI: confidence interval.

Supplementary Table 4: Correlation between plasma pTau181, pTau217 and clinical chemistry analytes

	pTau181			pTau217		
	n	Correlation coefficient	Significance Level P	n	Correlation coefficient	Significance Level P
Adiponectin	472	0.074	0.1089	472	0.064	0.164
Fasting glycemia	439	-0.084	0.0795	439	-0.058	0.2219
Triglycerides	421	-0.073	0.1358	421	-0.02	0.6892
Cholesterol total	421	-0.037	0.4432	421	-0.021	0.6712
Cholesterol HDL	419	-0.007	0.8866	419	-0.057	0.242
Cholesterol LDL	417	-0.04	0.4133	417	-0.023	0.6392
Creatinine	444	0.236	<0.0001	444	0.188	0.0001
eGFR	444	-0.24	<0.0001	444	-0.187	0.0001
Prealbumin	385	0.075	0.1408	385	0.08	0.1163
Albumin	442	-0.035	0.4587	442	0.015	0.7591
Total protein	438	-0.126	0.0082	438	-0.118	0.0134
CRP	430	-0.107	0.0262	430	-0.012	0.8069
Hemoglobin	439	0.046	0.3365	439	0.074	0.1203
Vitamin B12	408	0.001	0.9899	408	-0.09	0.0698
TSH	391	-0.031	0.5475	391	0.039	0.4444
Folate	216	-0.03	0.6651	216	-0.075	0.2693
Red-cell folate	343	0.095	0.0805	343	0.013	0.8078

Pearson's correlation coefficient in MCI participants (n) between clinical chemistry analytes and plasma pTau levels.


Supplementary Figure 1: Correlation between plasma pTau217, plasma pTau181 and hippocampal volume.



Legend:

Values of pTau217 (A) and pTau181 (B) in pg/mL were correlated with Hippocampal volume (R+L) (cm³). Pearson's correlation coefficient 'r' are indicated and they did not reach significance.


Supplementary Figure 2: Correlation between plasma pTau217 and pTau181 and CSF pTau181.



Legend:

Correlation between plasma pTau217 (Panel A) and pTau181 (Panel B) and CSF pTau181 in MCI according to amyloid status. Pearson's correlation coefficients 'r' were 0.49 (95% Confidence interval 0.38 to 0.58) and 0.37 (95% Confidence interval 0.26 to 0.48) for plasma pTau217 vs. CSF pTau181, and plasma pTau181 vs. CSF pTau181, respectively.

Supplementary Figure 3: Distribution of plasma pTau217 and pTau181 and ROC curves of plasma pTau217, plasma pTau181 and CSF pTau181 and CSF A β 42/40 in MCI according to conversion to AD.



Legend:

Values of pTau217 (A) and pTau181 (B) in pg/mL are represented in the conversion to AD (-) and (+) populations. (C) ROC curves of plasma pTau217, pTau181 and CSF pTau181 and Abeta42/40 in MCI according to conversion to AD. Respective AUC (95% Confidence interval): plasma pTau217: 0.722 (0.657 to 0.781), plasma pTau181 0.673 (0.606 to 0.736), CSF pTau181: 0.712 (0.646 to 0.771), and CSF A β 42/40: 0.733 (0.668 to 0.791).