

## Supplementary

Criteria considered for gout diagnosis	Included for gout diagnosis	Excluded following exploratory phase
<b>1. ICD-10-GM diagnosis code</b>	M10.00-M10.99: gout	M11.X: other crystal arthropathies.
<b>2. Documents</b> <ul style="list-style-type: none"> <li>Inpatient report (Admission or discharge letter from any type of care)</li> <li>Outpatient report (emergency room, specialist consultation)</li> </ul>	Regular expression query in full text for: <ul style="list-style-type: none"> <li>“gout”, “podagra”, “tophus”, “tophi”, “tophaceous”<sup>†</sup></li> </ul>	Exclusion of documents using a defined set of regular expression <sup>§</sup>
<b>3. Problem list of the EHR</b> (independent of the ICD-10-GM code)	Presence of one of the following keywords: <ul style="list-style-type: none"> <li>“gout”, “gouty”, “podagra”, “tophus”, “tophi”, “tophaceous”<sup>†</sup></li> </ul>	Exclusion of a set list of words See supplementary table 3
<b>4. Joint aspiration result</b>	Presence of monosodium urate crystals	None
<b>5. Medication</b> <ul style="list-style-type: none"> <li>Prescription of urate lowering therapy (ULT)</li> <li>Prescription form containing an ULT</li> </ul>	At least one of: <ul style="list-style-type: none"> <li>Allopurinol (ATC: M04AA01)</li> <li>Febuxostat (ATC: M04AA03)</li> <li>Probenecid (ATC: M04AB01)</li> <li>Lesinurad (ATC: M04AB05)</li> </ul>	Prescription of an ULT and presence of at least one of the following ICD-10 GM code at any time: C81 to C96: Malignant neoplasms, stated or presumed to be primary, of lymphoid, haematopoietic and related tissue
<b>6. Imaging report</b> <ul style="list-style-type: none"> <li>X-ray</li> <li>ultrasound</li> <li>dual-energy computed tomogram</li> </ul>	Regular expression query for the following words: <ul style="list-style-type: none"> <li>“gout”, “podagra”, “tophus”, “tophi”, “tophaceous”, “double contour”<sup>†</sup></li> </ul>	Exclusion of documents using a defined set of regular expression <sup>§</sup>

Supplementary table 1 : Variables and criterion used to assess gout diagnosis. \*Benzbromarone isn't authorized on the Swiss market. Lesinurad was pull out of the market in 2021. Rasburicase isn't authorized for chronic gout treatment. † French translation of the original words used are described in the supplementary table 3. § the R code, as well as a working example has been made available here: [https://gitlab.unige.ch/goutte/register\\_validation](https://gitlab.unige.ch/goutte/register_validation). ICD-10-GM: German Modifications of the International Classification of Diseases, 10<sup>th</sup> revision ATC: Anatomical Therapeutic Chemical Classification

Combination		Variables considered for risk factors	Criterion for gout risk factors (and ICD-10 GM codes)
AND		Sex and age	Women $\geq$ 65 years old Men $\geq$ 40 years old
AND		BMI	$> 25 \text{ kg/m}^2$
AND	OR	Metabolic syndrome	<ul style="list-style-type: none"> <li>• Diabetes or glucose intolerance (E10.X to E14.X, R73.X)</li> <li>• Hypertension (I10.X to I15.X)</li> <li>• Dyslipidaemia (E78.x)</li> <li>• BMI <math>&gt; 30 \text{ kg/m}^2</math></li> </ul>
	OR	Myocardial infarction	I21.X to I25.X
	OR	Chronic kidney disease	N18.X Problem list: Chronic kidney disease G3a or worse (including dialyses)
	OR	Lifestyle at risk to develop a gout	Deleterious use of alcohol (F10.1, F10.2, F10.3)
	OR	Uricaemia	Serum urate $> 500 \mu\text{mol/l}$ (8.4 mg/dl)

Supplementary table 2: Variables and criteria used to assess gout risk factors.

Category	Variables
Anthropometric & demographic data	Age Sex Weight Height BMI First language Country of residency Nationality Insurance class Living/death status Marital status
Clinical pathway	Inpatient gout episode of care Outpatient gout episode of care Departments involved Duration of hospitalisation
Diagnosis	Main diagnosis (inpatient)
Comorbidities (ICD-10-GM diagnostic)	Number of comorbidities Individual comorbidities Hypertension and hypertensive diseases: <ul style="list-style-type: none"> <li>• Essential hypertension: I10.00 – I10.91</li> <li>• Hypertensive cardiopathy: I11.00 – I11.91</li> <li>• Hypertensive nephropathy: I12.00 – I12.91</li> <li>• Hypertensive cardioneuropathy: I13.00-I13.91</li> <li>• Secondary hypertension: I15.00 – I15.91</li> </ul> Dyslipidaemia: <ul style="list-style-type: none"> <li>• Disorders of lipoprotein metabolism and other lipidaemias: E78.0-E78.9</li> </ul> Diabetes <ul style="list-style-type: none"> <li>• Type 1 diabetes: E10.01-E10.91</li> <li>• Type 2 diabetes: E11.01 – E11.91</li> <li>• Malnutrition diabetes: E12.11 – E12.91</li> <li>• Other specified diabetes mellitus: E13.01 – E13.91</li> <li>• Unspecified diabetes mellitus: E14.01 – E14.91</li> </ul> Cardiovascular and ischaemic heart disease: <ul style="list-style-type: none"> <li>• Angina pectoris: I20.00-I20-9</li> <li>• Acute myocardial infarction: I21.0 – I21.9</li> <li>• Recurring myocardial infarction: I22.0-I22.9</li> <li>• Complication of a myocardial infarction: I23.0 – I23.8</li> <li>• Other ischaemic cardiopathy: I24.0-I24.9</li> <li>• Chronic ischaemic cardiopathy: I25.0-I25.9</li> <li>• Cerebral infarction: I63.0 – I63.9</li> <li>• Stroke, not specified: I64</li> </ul>

	<ul style="list-style-type: none"> <li>• Occlusion and stenosis of precerebral arteries, not resulting in cerebral infarction: I65.0 – I65.9</li> <li>• Occlusion and stenosis of cerebral arteries, not resulting in cerebral infarction: I66.0-I66.9</li> <li>• Sequelae of cerebrovascular disease: I69.0 – I69.8</li> <li>• Atherosclerosis: I70.0 – I70.9</li> <li>• Arterial embolism and thrombosis: I74.0 – I74.9</li> <li>• Heart failure: I50.0 – I50.9</li> </ul> <p>Chronic kidney disease (stage ≥3): N18.3 – N18.9  Diseases of liver: K70-K77.8  Psychiatric disorders: F00-F99  Transplanted organ and tissue status: Z94.0 – Z94.9  Malignancies: C00-C97  Disorders of purine and pyrimidine metabolism: E79.1-E79.9</p>
Laboratory data	<p>Serum urate value  Serum creatinine and eGFR  CRP  Erythrocyte sedimentation rate  Liver function tests  Blood cell count</p>
Joint puncture	<p>Joint  Number of joint aspiration  Volume of synovial fluid  Presence of bacteriologic exam, Result of bacteriologic exam  Presence of cell count  Result of cell count  Presence of crystal evaluation  Crystal type in joint aspiration</p>
Chronic drug treatment	<p>Allopurinol  Febuxostat  Probenecid  Lesinurad  Rasburicase</p>
Drug prophylaxis during initiation of ULT	<p>Colchicine  Corticosteroid  NSAID</p>
Acute flare treatment	<p>Colchicine  Intra-articular corticoid injection (triamcinolone acetonide, betamethasone, triamcinolone hexacetonide)  Oral, IV or IM corticoid (dexamethasone, prednisone, methylprednisolone, prednisolone)  Anakinra or canakinumab  NSAID</p>
Gout predisposing drugs	<p>Aspirin  Ciclosporin</p>


	Tacrolimus Loop diuretics (furosemid, torasemid) Thiazide diuretics (hydrochlorothiazide) Thiazide-like diuretics (metolazone, indapamide, chlorthalidone)
Gout protecting drugs	Fenofibrate Losartan SGLT-2 inhibitors
Pain score	Visual analog scale
Fall	Fall event during an acute gout flare ("déclaration de chute")
Rheumatology consultation	Presence or absence
Imaging	Presence of an X-ray of the painful articulation Presence of a CT dual energy of the painful articulation Presence of an ultrasound of the painful articulation Gout-related damage or tophi or double contour sign

*Supplementary Table 3: Data extracted from the electronic health record.*


Query	French translation	English translation
Documents	<ul style="list-style-type: none"> <li>• « goutte »</li> <li>• « podagre »</li> <li>• « tophus »</li> <li>• « tophi »</li> <li>• « tophacée »</li> </ul>	<ul style="list-style-type: none"> <li>• Gout</li> <li>• Podagra</li> <li>• Tophus</li> <li>• Tophi</li> <li>• Tophaceous</li> </ul>
Search in the problem list of the EHR (independent of the ICD-10-GM code)	<ul style="list-style-type: none"> <li>• « goutte »</li> <li>• « goutteux »</li> <li>• « goutteuse »</li> <li>• « podagra »</li> <li>• « tophus »</li> <li>• « tophi »</li> <li>• « tophacée »</li> </ul>	<ul style="list-style-type: none"> <li>• Gout</li> <li>• Gouty</li> <li>• Gouty</li> <li>• Podagra</li> <li>• Tophus</li> <li>• Tophi</li> <li>• Tophaceous</li> </ul>
Imaging report	<ul style="list-style-type: none"> <li>• « goutte »</li> <li>• « podagre »</li> <li>• « tophus »</li> <li>• « tophi »</li> <li>• « tophacée »</li> <li>• « double-contour »</li> </ul>	<ul style="list-style-type: none"> <li>• Gout</li> <li>• Podagra</li> <li>• Tophus</li> <li>• Tophi</li> <li>• Tophaceous</li> <li>• Double contour</li> </ul>

*Supplementary table 4: original translation of the words used in the queries to detect gout patients.*

## Outpatients



## Inpatients



Supplementary Figure 1A and 1B: Upset-plot of the six queries identifying gout patients in the electronic health record of the Geneva University Hospital when using a combination of criteria (Problem list OR Aspiration OR  $\geq 2$  other criteria) and by setting stratification. Rare combinations of criteria were not displayed. ICD-10-GM diagnoses are coded only during an inpatient stay

<b>Criterion present alone</b>	<b>Total</b>	<b>Gout</b>	<b>No gout</b>	<b>Equivocal</b>	<b>Positive predictive value (CI 95%)</b>
ICD-10-GM Codes	5	2	3	0	40.0% (11.8-76.9)
Problem list	8	7	1	0	87.5% (52.9-99.4)
Aspirations	6	4	2	0	66.7% (30.0-90.3)
Drugs	178	58	28	92	32.6% (26.1-39.8)
Documents	143	84	55	4	58.7% (50.5-66.5)
Radiology reports	9	2	5	2	22.2% (6.3-54.7)

*Supplementary table 5 : Absolute numbers of patients and positive predictive value (PPV) of each criterion alone (a patient is detected only by the criterion). PPV was calculated by considering gout present versus gout absent or equivocal meaning equivocal cases were classified as non-gout in the PPV analysis.*



Query (inpatients)	Total	Gout	No gout	Equivocal	Positive predictive value (CI 95%)
<b>Present irrespective of other criteria</b>					
ICD-10-GM Codes	96	89	6	1	92.7% (85.7-96.4)
Problem list	75	74	1	0	98.7% (92.8-99.9)
Joint aspiration	38	37	1	0	97.4% (86.5-99.9)
Drugs	208	202	1	5	97.1% (93.9-98.7)
Documents	167	156	8	3	93.4% (88.6-96.3)
Radiology reports	24	24	0	0	100.0% (86.2-100.0)
<b>Present alone</b>					
ICD-10-GM Codes	0	0	0	0	
Problem list	4	4	0	0	100.0% (51.0-100.0)
Joint aspiration	5	4	1	0	80.0% (37.6-99.0)
Drugs	0	0	0	0	
Documents	0	0	0	0	
Radiology reports	0	0	0	0	
<b>Combination query</b>					
Problem or joint aspirations or $\geq 2$ other criteria	180	168	9	3	93.3% (88.7-96.1)

Supplementary table 6: Positive predictive value according to each query and a combination of queries, among 180 inpatients PPV.

<b>Query (outpatients)</b>	<b>Total</b>	<b>Gout</b>	<b>No gout</b>	<b>Equivocal</b>	<b>Positive predictive value (CI 95%)</b>
<b>Present irrespective of other criteria</b>					
ICD-10-GM Codes	31	30	1	0	96.8% (83.8-99.8)
Problem list	66	63	3	0	95.5% (87.5-98.4)
Joint aspiration	10	9	1	0	90.0% (59.6-99.5)
Drugs	76	76	0	0	100.0% (95.2-100.0)
Documents	73	67	5	1	91.8% (83.2-96.2)
Radiology reports	14	10	3	1	71.4% (45.4-88.3)
<b>Present alone</b>					
ICD-10-GM Codes	0	0	0	0	
Problem list	4	3	1	0	75.0% (30.1-98.7)
Joint aspiration	1	0	1	0	0.0% (0.0-94.9)
Drugs	0	0	0	0	
Documents	0	0	0	0	
Radiology reports	0	0	0	0	
<b>Combination query</b>					
Problem or joint aspirations or $\geq 2$ other criteria	82	74	7	1	90.2% (81.9-95.0)


*Supplementary table 7: Positive predictive value according to each query and a combination of queries, among 82 outpatients PPV. ICD-10-GM codes are not available for inpatient.*

<b>Risk factor</b>	<b>Overall</b>	<b>Alcohol</b>	<b>MI</b>	<b>CKD</b>	<b>MS</b>	<b>Uricemia</b>
<b>Characteristics</b>						
Total number	15646	3249	4818	4188	1906	1485
Age (mean (SD))	73.09 (13.2)	65.80 (13.2)	73.68 (13.0)	79.33 (11.1)	71.75 (10.8)	71.22 (13.6)
BMI (mean (SD))	33.78 (28.7)	32.56 (25.9)	32.85 (29.2)	33.69 (29.7)	38.80 (32.8)	33.27 (22.6)
Number of death (%) at 31.12.2022	6567 (42.0)	1051 (32.3)	1882 (39.1)	2314 (55.3)	622 (32.6)	698 (47.0)
Detected by the register (%)	2588 (16.5)	362 (11.1)	524 (10.9)	817 (19.5)	287 (15.1)	598 (40.3)

Supplementary Table 8: Absolute number of patients with risk factor for gout, their characteristics and number of patients detected by our algorithm detecting gout patients. MI: myocardial infarction. CKD: chronic kidney disease. MS: Metabolic syndrome. BMI: body mass index

<b>Criterion present alone</b>	<b>Total</b>	<b>Gout</b>	<b>No gout</b>	<b>Equivocal</b>	<b>Negative predictive value (CI 95%)</b>
Uricaemia > 500 µmol/l	22	6	16	0	72.7% (51.8 to 86.8)
Chronic kidney disease	86	5	79	2	91.9% (84.1 to 96.0)
Metabolic syndrome	39	3	36	0	92.3% (79.7 to 97.3)
Myocardial infarction	128	1	126	1	98.4% (94.5 to 99.6)
Deleterious use of alcohol	107	2	104	1	97.2% (92.1 to 99.0)

Supplementary table 9 : Absolute numbers of patients and negative predictive value (NPV) of each criterion alone (a patient is detected only by the criterion). NPV was calculated by considering gout present versus gout absent or equivocal meaning equivocal cases were classified as gout in the NPV analysis.



Supplementary figure 2: Evolution of gout diagnostic according to the criterion or combination of criteria to establish the diagnosis at first detection in the register (or second in case of query combination). Other criteria correspond to a combination of  $\geq 2$  criterion among the documents, ICD-10-GM codes, imaging reports and drugs queries.