

Nipah Virus Therapeutics: A Systematic Review to Support Prioritisation for Clinical Trials

*Xin Hui S Chan DPhil,^{1,2,3} Ilsa L Haeusler BMBCh,¹ Bennett J K Choy MBChB,¹
Md Zakiul Hassan MBBS,^{1,2,4} Junko Takata BMBCh,³ Tara P Hurst PhD,¹ Luke M Jones PhD,^{1,5}
Shanghavi Loganathan BMBCh,⁶ Elinor Harriss MSc,⁷ Jake Dunning PhD,^{1,2,8} Prof Joel Tarning PhD,^{9,10}
Prof Miles W Carroll PhD,^{1,5} Prof Peter W Horby PhD,^{1,2} Prof Piero L Olliaro PhD^{1,2}*

¹Pandemic Sciences Institute, Nuffield Department of Medicine, University of Oxford, Oxford, UK

²International Severe Acute Respiratory and Emerging Infection Consortium, University of Oxford, Oxford, UK

³Department of Clinical Infection, Oxford University Hospitals NHS Foundation Trust, Oxford, UK

⁴Programme for Emerging Infections, Infectious Diseases Division, International Centre for Diarrheal Disease Research, Dhaka, Bangladesh

⁵Centre for Human Genetics, Nuffield Department of Medicine, University of Oxford, Oxford, UK

⁶Academic Foundation Programme, Kent Surrey and Sussex Deanery, London, UK

⁷Bodleian Libraries, University of Oxford, Oxford, UK

⁸Department of Infectious Diseases, Royal Free London NHS Foundation Trust, London, UK

⁹Centre for Tropical Medicine and Global Health, Nuffield Department of Medicine, University of Oxford, Oxford, UK

¹⁰Mahidol-Oxford Tropical Medicine Research Unit, Faculty of Tropical Medicine, Mahidol University, Bangkok, Thailand

Corresponding Author:

Dr Xin Hui S Chan
Pandemic Sciences Institute
Old Road Campus Research Building
Roosevelt Drive
Oxford, OX3 7DQ
United Kingdom

+44 1865 612974

Research in context

Evidence before this study

Nipah virus infection is a bat-borne zoonosis with person-to-person transmission, a case fatality rate of 38-75%, and recognised pandemic potential. No therapeutics or vaccines have been licensed to date, and only few candidates are in development. We conducted this systematic review to assess the evidence for the safety and efficacy of therapeutic options (monoclonal antibodies and small molecules) for Nipah virus and other henipaviral diseases to support candidate prioritisation for further evaluation in clinical trials.

We searched bibliographic databases for journal articles, conference abstracts, and patents: PubMed, Ovid Embase, Ovid CAB Abstracts, Ovid Global Health, Scopus, Web of Science (all databases), and the WHO Global Index Medicus. “Henipavirus” or “Nipah” or “Hendra” along with “therapeutics” or “monoclonal” were the title, abstract, and subject heading keywords, with synonyms and variant spellings as additional search terms. We searched trial registries for clinical trials of Henipavirus, Nipah virus, and Hendra virus at all stages of recruitment: Cochrane Central Register of Controlled Trials, ClinicalTrials.gov, and the WHO International Clinical Trials Registry Platform. We searched the Trip database and WHO website for guidelines and reports. All searches were conducted on 30 May 2022. We did not apply language or publication date limits.

Studies were included if they contained primary data on the safety and/or efficacy of monoclonal antibodies (*in vivo*) or small molecules (*in vivo* or *in vitro*) for the treatment and/or prophylaxis of Nipah, Hendra, and related *Henipaviridae*. Almost all had critical or high risk of bias.

Added value of this study

This is the most detailed systematic review and analysis of the Nipah virus therapeutics landscape to date, including all available *in vivo* and related *in vitro* data on the safety, efficacy, and pharmacokinetics of monoclonal antibodies and small molecules with the specific aim of supporting prioritisation for clinical trials. We also present a roadmap for how *in vivo* development of Nipah therapeutics could be strengthened to achieve greater equity, efficiency, and effectiveness.

Implications of the available evidence

At present, there is sufficient evidence to trial only m102.4 and remdesivir for prophylaxis and early treatment of Nipah virus infection. Well-designed clinical efficacy trials as well as *in vivo* pharmacokinetic-pharmacodynamic studies to optimise selection and dosing of therapeutic candidates in animal challenge and natural human infection are needed.

Abstract

Nipah virus disease is a bat-borne zoonosis with person-to-person transmission, a case fatality rate of 38-75%, and recognised pandemic potential. The first reported outbreak occurred in Malaysia and Singapore in 1998, since followed by multiple outbreaks in Bangladesh and India. No therapeutics or vaccines have been licensed to date, and only few candidates are in development. This systematic review aimed to assess the evidence for the safety and efficacy of therapeutic options (monoclonal antibodies and small molecules) for Nipah virus and other henipaviral diseases in order to support candidate prioritisation for further evaluation in clinical trials. At present, there is sufficient evidence to trial only m102.4 and remdesivir (singly and/or in combination) for prophylaxis and early treatment of Nipah virus disease. In addition to well-designed clinical efficacy trials, *in vivo* pharmacokinetic-pharmacodynamic studies to optimise selection and dosing of therapeutic candidates in animal challenge and natural human infection are needed.

149 words

Introduction

Nipah virus disease is a zoonotic infection acquired through contact with or ingestion of contaminated body fluids of infected mammals¹⁻³. Pteropid fruit bats (flying foxes) are the primary reservoir of Nipah virus. Secondary hosts include domestic animals^{4,5} (pigs, horses, cows) and humans. There is person-to-person transmission. Its clinical presentation ranges from asymptomatic⁶ to an acute respiratory syndrome and fatal encephalitis^{1,7}. After an incubation period of four to 14 days, fever, headache, and myalgia may be followed by shortness of breath and cough or confusion and seizures which can rapidly progress to coma within 24 to 48 hours^{1,7}. Disease occurs in all age groups^{1,7}. The case fatality ratio (CFR) is estimated to be between 38 to 75% and debilitating long-term neurological complications, such as paralysis, are common in Nipah survivors^{1,8}.

Nipah virus is part of the genus *Henipavirus* along with Hendra virus which also causes fatal encephalitis and respiratory disease in horses and humans. Both Nipah and Hendra viruses are biosafety level 4 (BSL-4) pathogens requiring the highest level of laboratory containment precautions. The other bat-borne members of the genus (Cedar and Kumasi viruses) are not known to cause human disease⁹.

First identified in 1998, following an outbreak among pig farmers and abattoir workers in Malaysia¹⁰ and Singapore¹¹, Nipah virus is named after the Malaysian village from which the virus was first isolated. 283 cases of encephalitis and 109 deaths were recorded, a CFR of 38.3%¹². This outbreak was halted with mass culling of more than one million pigs and comprehensive modernisation of pig farming practice, including the separation of fruit tree plantations from pig farms¹³. There have been no further Nipah cases in Malaysia and Singapore in the subsequent 25 years, and only one further outbreak of the Nipah virus Malaysia (NiV-M) strain in the Philippines in 2014¹⁴ related to horse slaughter and consumption. Outbreaks of the Nipah virus Bangladesh (NiV-B) strain have been reported in Bangladesh³ and India (West Bengal¹⁵, Kerala¹⁶), with healthcare workers¹⁵ and family¹⁷ caring for infected patients emerging as another important risk group. The highest mortality rates have been recorded in Bangladesh where outbreaks occur almost annually in the winter following harvesting and consumption of contaminated raw date palm sap², a local delicacy. Since 2001, there have been 335 cases with 237 deaths in Bangladesh, a CFR of 70.7%³. The 2023 outbreak in Bangladesh was the largest since 2015 with 14 cases and 10 deaths. A second outbreak occurred less than 6 months later in Kerala, India with six cases and two fatalities¹⁸. Patient outcomes have not improved in 25 years since the first reported outbreaks due in part to the market failure typical of counter-measure development for a high-consequence pathogen¹⁹.

There are no licensed vaccines or therapeutics for Nipah virus infection, and only a few candidates are currently in development²⁰. In recognition of the need for vaccines and therapeutics, Nipah has been a priority disease in the World Health Organization (WHO) Research & Development Blueprint since 2018²¹. Clinical evaluation is limited by the infeasibility of a controlled human infection model and the small number of patients in sporadic outbreaks. Assessment of efficacy is currently reliant on animal challenge studies conducted in BSL-4 facilities.

We conducted this systematic review to assess the evidence for the safety and efficacy of therapeutic options (monoclonal antibodies [mAbs] and small molecules) for Nipah virus and other *Henipaviridae* causing human disease in order to support candidate prioritisation for further evaluation in clinical trials.

Methods

This systematic review was registered prospectively on the PROSPERO database (CRD42022346563) and adheres to the PRISMA 2020 reporting guidelines (appendix).

Search Strategy

We conducted an electronic literature search of the following bibliographic databases for journal articles, conference abstracts, and patents: PubMed, Ovid Embase, Ovid CAB Abstracts, Ovid Global Health, Scopus, Web of Science (all databases), and the WHO Global Index Medicus. “Henipavirus” or “Nipah” or “Hendra” along with “therapeutics” or “monoclonal” were the title, abstract, and subject heading keywords, with synonyms and variant spellings as additional search terms. We searched the following trial registries for clinical trials of Henipavirus, Nipah virus, and Hendra virus at all stages of recruitment: Cochrane Central Register of Controlled Trials, ClinicalTrials.gov, and the WHO International Clinical Trials Registry Platform. We searched the Trip database and WHO website for guidelines and reports. Full search strategies are in the appendix.

All searches were conducted on 30 May 2022. We did not apply language or publication date limits.

References were imported into EndNote and de-duplicated then screened against eligibility criteria. Reference lists of eligible records were checked for additional relevant studies.

Eligibility Criteria

Studies were included if they contained primary data on the safety and/or efficacy of mAbs (*in vivo*) or small molecules (*in vivo* or *in vitro*) for the treatment and/or prophylaxis of Nipah and/or Hendra infections. Studies on candidates without therapeutic applications (e.g. mAbs for diagnostics) or with only *in silico* data were excluded.

Data Extraction

We extracted data on the viruses studied, study characteristics (funder, year, location, design), intervention characteristics (drug, dose, route, administration timepoints), efficacy outcomes (all measures, all timepoints), and safety outcomes (all measures, all timepoints). Study investigators and experts were contacted for further information if necessary.

Data Analysis

The review pilot identified significant heterogeneity in study designs, outcome measures, and reporting. Quantitative data synthesis was deemed not possible. All available data were therefore prespecified to be summarised in tabular format by individual therapeutic candidate as a narrative synthesis prioritising clinical and animal studies.

Quality Assessment

Risk of bias assessment was undertaken for the study designs for which standardised tools exist: Risk of Bias 2 (RoB 2²²) for randomised clinical trials (RCTs), Risk of Bias in Non-randomized Studies of Interventions (ROBINS-I²³) for non-randomised clinical studies, and Systematic Review Centre for Laboratory Animal Experimentation (SYRCLE²⁴) for animal studies.

Review Team & Tools

At least two independent reviewers performed screening (titles and abstracts, then full texts), agreed study eligibility, extracted data, and undertook risk of bias assessment using Covidence (Veritas Health Innovation Ltd, Melbourne, Australia).

Role of the Funding Source

Our funders had no role in study design, data collection, data analysis, data interpretation, or writing of the report. All authors have access to the data in the study and accept responsibility to submit for publication.

Results

Included Studies

We identified 56 eligible studies (Figure 1): 12 of mAbs with clinical and/or animal data²⁵⁻³⁶ (Table 1 & Supplementary Table I), 25 of small molecules with clinical and/or animal data (Table 2 & Supplementary Tables II-III), and 19 of small molecules with *in vitro* data only (Supplementary Table IV).

There was only one clinical trial, a first-in-human phase 1 study in healthy volunteers of m102.4²⁶, a mAb targeting the Hendra virus (HeV) envelope G glycoprotein, conducted in Australia. Of the eight reports of compassionate use for treatment or post-exposure prophylaxis during Hendra or Nipah virus outbreaks, seven were case series of fewer than 10 patients in Australia³⁷, India (Kerala)^{25,38-41}, and Singapore¹¹. The remaining outbreak report was from the two centres where 194 of the 283 cases in the 1998 Malaysia outbreak were treated, the majority with ribavirin⁴².

Of the 23 animal studies, there were seven studies in non-human primates (African green monkeys [AGMs])^{27-29,43-46}, five in ferrets^{30-32,34,47}, and 12 in Syrian golden hamsters^{33,35,36,46,48-55}. All except one involved infectious challenge with Nipah and/or Hendra virus (Supplementary Table V). Nipah virus Malaysia (NiV-M) was the most common challenge strain used in 12 studies^{28,30,32,36,46-50,52,53,55}, followed by six studies using Nipah virus Bangladesh (NiV-B)^{27,33,34,43,44,51} and five HeV^{29,32,35,45,49}.

All animal challenge studies reported death, and time of death, as outcome measures. The majority also reported clinical outcomes (all: signs and symptoms; AGMs only: radiological changes, blood test abnormalities) with day of onset, and a smaller majority reported pathology and virology (detection of RNA, antigen, or live virus by culture) at necropsy. A minority assessed correlation between drug concentrations and survival.

Monoclonal Antibodies

The 12 articles on mAbs reported data on six sets of antibodies from three research groups. Seven articles were on m102.4, an anti-HeV-G glycoprotein antibody, the most advanced candidate in clinical development (Table 1 & Supplementary Table I).

m102.4 was the only Nipah drug candidate with clinical data from an RCT²⁶ (phase 1) and with *in vivo* data from more than one animal species (AGMs²⁷⁻²⁹ and ferrets^{30,31}) challenged with different henipaviruses (NiV-B²⁷, NiV-M^{28,30}, HeV²⁹) (Table 1).

Currently available data in humans support its safety. In a first-in-human dose-escalation randomised placebo-controlled trial of intravenous (IV) m102.4 (single doses of 1-20mg/kg + two doses of 20mg/kg 72 hours apart) in 40 healthy adult volunteers followed up for ~4 months between

2015 and 2016, no serious adverse events were reported²⁶. The frequency of adverse events, of which headache was the most common, were similar between the different treatment and placebo groups²⁶. No anti-m102.4 antibodies were detected²⁶. Prior to this trial, 14 individuals aged 8-59 years had received m102.4 as post-exposure prophylaxis on compassionate grounds for Hendra (n=13) and Nipah virus (n=1) infections in Australia and the USA respectively²⁶. Of these, two individuals experienced infusion-related febrile reactions that were attributed to an early production process of the antibody²⁶. There was also one outbreak report describing a single patient receiving m102.4 as post-exposure prophylaxis in Kerala, India, in 2018²⁵. The patient was reported to have recovered completely, but no further details were provided.

The four animal challenge studies of m102.4 provide evidence of its efficacy in preventing death and severe disease in all treated animals when administered as a single dose in ferrets 10 hours after oronasal NiV-M inoculation³⁰ (n=3) or in a two-dose regimen given 48 hours apart in monkeys starting within five days after intratracheal NiV-M²⁸ (n=12) or HeV²⁹ (n=12) challenge. In comparison, all control animals died within 8-10 days in these three studies. However, the treatment window for the two-dose regimen after NiV-B challenge in monkeys²⁷ was shorter than that for NiV-M and HeV, with only the animals treated within three days (n=6) from inoculation surviving to the end of the study while the monkeys treated on days 5 and 7 (n=2) had similar outcomes to controls (n=2). In these four animal challenge studies, protection from disease was supported by the absence of pathological changes in treated animals on necropsy versus gross pathology in control animals, as well as the correlation of antibody levels with survival, including on day 3 post-challenge.

Additionally, the developers of m102.4 are now also developing h5B.3, an anti-NiV-F glycoprotein antibody. When administered intraperitoneally (IP) to ferrets in a regimen of two 20mg/kg doses given 48 hours apart starting within three days of intranasal challenge with NiV-M (n=6) or HeV (n=3), h5B.3 protected all treated ferrets from severe disease³².

Two articles^{33,34} from 2020 and 2021, from Vanderbilt University in the United States, describe two sets of anti-HeV receptor binding protein antibodies. HENV-26 (n=5) and HENV-32 (n=5) administered as two doses of 15mg/kg IP on days 3 and 5 post-intranasal NIV-B challenge each protected ferrets from death and severe disease compared to controls (n=3)³⁴. HENV-103 and HENV-117 protected all hamsters from intranasal NIV-B challenge in combination (n=5) but not individually (n=5 each) nor as two bispecific antibodies of different designs (n=5 each)³³.

A further two articles^{35,36} from Institut National de la Santé et de la Recherche Médicale (INSERM), France dating from the late 2000s, describe two groups of anti-NiV-F and anti-NiV-G protein antibodies. These were detailed studies of protection, dose titration, and therapeutic window involving 124³⁶ and 54³⁵ hamsters respectively. However, there have been no further studies of these candidates in the subsequent two decades.

Small Molecules

The 25 articles on small molecules with *in vivo* (Table 2 & Supplementary Table II) and *in vitro* (Supplementary Table III) data described 10 potential therapeutics and one group of syndrome-directed broad-spectrum empirical antimicrobials.

Ribavirin

Ten articles were on ribavirin, a repurposed nucleoside analogue prodrug. These were six clinical case series for treatment^{37,39-42} and post-exposure prophylaxis^{37,38} for Nipah³⁸⁻⁴² and Hendra³⁷ outbreaks (Table 2), three animal challenge studies of AGMs (HeV⁴⁵ only) and hamsters (both HeV⁴⁹ and NiV-M^{49,50}) (Table 2), plus three sets of *in vitro* experiments with HeV^{49,56} and NiV-M^{49,50} (Supplementary Table III).

The small numbers of patients treated with ribavirin ($n < 10$ in all except the Malaysia Nipah outbreak⁴²) and pragmatic observational designs of case series precluded definitive statements about clinical efficacy. Dose regimens were different between the four publications^{37,38,40,42} reporting this information and were not reported in the remaining two^{39,41}. All eight healthcare workers in the only post-exposure prophylaxis case series³⁸ of ribavirin did not complete the prescribed course due to adverse effects: six of eight had symptoms (such as fatigue or headache) or transient laboratory abnormalities (increased bilirubin and/or decreased haemoglobin levels).

In the three animal studies, when administered IP^{49,50} or subcutaneously^{45,50} (SC) 24 hours pre-challenge or within 12 hours post-challenge, ribavirin 50-150mg/kg/day delayed time to but did not prevent death or symptoms after NiV-M inoculation in hamsters^{49,50} ($n=17$) or HeV inoculation in AGMs⁴⁵ ($n=6$), compared to untreated controls. Ribavirin 60mg/kg/day neither delayed nor prevented death in HeV-challenged hamsters ($n=5$)⁴⁹. Systemic toxicity from high dose 200mg/kg/day of ribavirin IP was seen in both infected and uninfected (control) hamsters necessitating euthanasia⁴⁹.

In vitro experiments of ribavirin assessed viral replication through virus yield reduction⁵⁶, cytopathic effect⁵⁰, and dose response⁴⁹ assays in NiV-M and HeV infected Vero^{50,56} and HeLa⁴⁹ cells. Ribavirin doses used to achieve 58-fold⁵⁶ or 100% reductions^{49,50} in viral yield were high (50-409 μ M) compared to half-maximal inhibitory concentrations⁴⁹ (IC_{50}) for NiV-M (4.18 μ M) and HeV (4.96 μ M).

Chloroquine

Three articles were on the widely used 4-aminoquinoline antimalarial chloroquine^{47,49,57}: two animal challenge studies^{47,49} (Table 2) and two sets of *in vitro* experiments^{49,57} (Supplementary Table III). Ferrets administered chloroquine 25mg/kg/day IV 24 hours before ($n=3$) and 10 hours after ($n=3$) NiV-M challenge had disease courses identical to controls⁴⁷. NiV-M and HeV inoculated hamsters treated six hours after challenge with chloroquine 50mg/kg IP on alternate days as monotherapy ($n=5$ per virus) or in combination with ribavirin 30mg/kg IP twice a day ($n=5$ per virus) died earlier or at the same time respectively as untreated controls⁴⁹. Chloroquine 50mg/kg/day IP was also ineffective⁴⁹. Higher doses of 100 and 150mg/kg/day of chloroquine IP were consistently lethal by day 2 in both infected and uninfected hamsters⁴⁹.

Remdesivir

The three articles and one abstract on remdesivir, a nucleoside analogue, reported two AGM NiV-B challenge studies assessing IV remdesivir^{43,44} (Table 2) and *in vitro* data from multiple assays on both the IV⁵⁸ and oral⁵⁹ formulations (Supplementary Table III). Remdesivir 10mg/kg/day given from one day post-challenge protected all four AGMs from death, with mild transient respiratory signs in two and detectable viral RNA in brain tissue of one⁴³. Controls all died after respiratory symptoms with detectable viraemia⁴³. Reporter virus, cytopathic effect, and virus yield reduction assays for remdesivir IV (GS5734⁵⁸) and oral (GS441524⁵⁹) as well as viral antigen reduction and minigenome assays for GS5734⁵⁸ were performed in cell types including HeLa and human small airway epithelial

cells. Mean 50% maximal effective concentration (EC_{50}) values were sub-micromolar for both and an order of magnitude lower for GS5734⁵⁸ (0.029-0.066 μ M) than GS441524⁵⁹ (0.19-0.95 μ M).

Favipiravir

The single article on favipiravir⁴⁸, a nucleoside analogue prodrug, contained data from an NiV-M hamster challenge study (Table 2) and *in vitro* assays (Supplementary Table III). Hamsters loaded with 600mg/kg SC immediately post-challenge then given a maintenance dose of 300mg/kg orally twice a day (n=5) or SC daily (n=5) for 13 days all survived without clinical signs or detectable pathology or viral antigen on necropsy while controls all died by day 5-6. The doses used in virus yield (100 μ M) and delayed treatment (250 μ M) assays to attain 100% and 10-fold (at one-hour post-infection) viral reductions, respectively, were high. EC_{50} values for HeV, NiV-B, and NiV-M were 11.7 μ M, 14.8 μ M, and 44.2 μ M respectively⁴⁸.

Others

Six other groups of small molecules were studied, none of which provided complete protection from death at the doses used in animal challenge models. 6-azauridine, the nucleoside analogue metabolite of previously licensed azaribine, delayed mean time to death by ~1 day, but did not prevent death, when given immediately prior to full-dose NiV-M challenge (350 x median lethal dose [LD_{50}]) as a 175mg/kg/day continuous SC infusion for 14 days in hamsters⁵⁰. Rintatolimod, a toll-like receptor 3 agonist, provided partial protection at 3mg/kg/day IP for 10 days, administered from 2 hours after low-dose (35 x LD_{50}) NiV-M inoculation⁵⁰ (Table 2 & Supplementary Table II).

Periodate heparin, an experimental glycosaminoglycan competitive inhibitor of *trans*-infection, protected one of five NiV-M challenged hamsters at a dose of 10mg/kg/day SC for 12 days from day of infection⁵². Despite promising *in vitro* results, experimental cell entry inhibitors like the lectin griffithsin (oxidation resistant and trimeric monomer)⁵¹ and fusion inhibitory lipopeptides (cholesterol and tocopherol-based)^{46,54} administered at 10mg/kg/day intranasally (hamsters^{46,53,54}) or intratracheally (AGMs⁴⁶) prevented death in up to half of each group of NiV-B⁵¹ or NiV-M^{46,53} challenged animals (Table 2). Defective interfering virus particles given IP or intranasal also had partial efficacy in NiV-M challenged hamsters⁵⁵ while virus yield reduction assays had up to an order of magnitude greater reduction in NiV-M than NiV-B infected Vero cells⁶⁰ (Supplementary Table III).

Risk of Bias

Almost all the *in vivo* studies had critical (six of eight case series) or high (18 of 23 animal studies) risks of bias. Only three studies were assessed to have low risk of bias: the one RCT (of healthy volunteers)²⁶, an outbreak report of a single case⁴¹, and an NiV-B challenge study in AGMs⁴³. The remaining five^{25,30,32,52,61} studies had unclear risk of bias (Supplementary Figures I-V).

Discussion

To our knowledge, this is the most detailed review to date of the therapeutics landscape for Nipah and Hendra virus disease with the specific aim of supporting candidate prioritisation for clinical trials. We did not identify any ongoing or completed therapeutic efficacy RCTs for Nipah or Hendra virus infection. There were no data on *in vivo* drug resistance.

Drugs

The pipeline of therapeutics with the potential to be deployed rapidly at the outset of a henipavirus outbreak is currently limited to a few mAbs and repurposed small molecules with efficacy data from animal challenge models (Table 3). The comparative advantages of mAbs and small molecules are summarised in Table 4.

Of the mAbs, only m102.4 has been studied in humans, with safety and pharmacokinetic (PK) data from a phase 1 RCT in healthy adults²⁶. m102.4 is also the only mAb with efficacy (from challenge with NiV-B²⁷, NiV-M^{28,30}, and HeV²⁹) and PK data (without challenge^{29,31}) from two animal species. Further PK studies of mAb candidates to determine minimal doses for efficacy could help to make scale-up more cost-effective.

Of the small molecules, animal efficacy data were supportive for remdesivir and favipiravir, equivocal for ribavirin, and negative for chloroquine. Remdesivir was the only small molecule with *in vivo* data from challenge with NiV-B⁴³, the strain closely related to those causing recent Nipah outbreaks in Bangladesh and India⁶², and has accumulated acceptable safety data from its widespread intravenous use in COVID-19⁶³. While favipiravir prevented death in NiV-M challenged hamsters after a subcutaneous loading dose followed by subcutaneous or oral maintenance doses⁴⁸, and could be an attractive choice for post-exposure prophylaxis with a licensed oral formulation⁶⁴, its non-linear clinical pharmacokinetics seen in Ebola⁶⁵, influenza⁶⁶, and COVID-19⁶⁷ necessitate further dose optimisation prior to inclusion in Nipah trials. This non-linearity is thought to be explained by concentration-dependent aldehyde oxidase inhibition reproducible in non-human primates⁶⁸, and it is unclear if there is an additional infection-specific contribution. PK studies of parenteral (including intravenous⁶⁹) favipiravir in NiV-B inoculated non-human primates would be a key next step in favipiravir evaluation. Notably, favipiravir is associated with teratogenicity in four animal species⁶⁴ and further data on its safety in humans are needed^{70,71}.

Ribavirin prolongs survival but does not prevent death in monkeys and hamsters challenged with HeV⁴⁵ and NiV-M^{49,50}, and is toxic to hamsters at high doses⁴⁹. Issues with clinical tolerability (fatigue, anaemia, and hyperbilirubinaemia)³⁸ are further likely to reduce adherence to a ribavirin-containing post-exposure prophylaxis regimen. Clinical reports of ribavirin in Nipah and Hendra outbreaks were all observational with dosing based on that used for Lassa fever⁴². Recent clinical and PK meta-analyses of ribavirin in Lassa treatment highlight the lack of robust evidence for its efficacy⁷² and that conventional dosing regimens are unlikely to reliably achieve the serum concentrations required to inhibit Lassa virus replication⁷³. PK modelling is ongoing for available ribavirin dosing regimens for Nipah and Hendra. Ribavirin remains part of the Nipah treatment guidelines in India^{74,75} but not Bangladesh⁷⁶. Consultation with Nipah stakeholders would establish whether the potential to delay death by hours to days justifies further use. Chloroquine did not protect ferrets⁴⁷ or hamsters⁴⁹ from NiV-M and was lethal at higher doses in hamsters⁴⁹. The narrow therapeutic window of chloroquine is well-established in clinical practice, where it is a safe and effective antimalarial but has also been employed for rapid self-poisoning in deliberate overdose^{77,78}. Chloroquine should not be used for the treatment or prevention of Nipah or Hendra infection.

Promising *in vitro* efficacy has yet to translate into convincing *in vivo* protection for the experimental small molecules in Table 2. The parent drugs of 6-azauridine (azaribine) and ALS-8112 (lumicitabine) have been withdrawn from market and development respectively due to safety concerns of thrombosis⁷⁹ and paediatric neutropenia⁸⁰. It is unclear whether periodate heparin, fusion lipopeptides, and defective viral particles can be manufactured at scale or are stable for stockpiling.

Treatment Indications & Use Cases

Antiviral drugs, whether mAbs or small molecules, appear to have a narrow temporal window within which they are likely to have clinically relevant efficacy, limiting their use to prophylaxis (pre- and post-exposure) and possibly early treatment¹⁹. They could also play a key role in providing bridging protection prior to vaccine response or availability. The time window for protection post-challenge⁸¹ is shorter with NiV-B than NiV-M in monkeys²⁷ although this has yet to be validated in humans.

Immunomodulators could be used in combination with pathogen-directed antivirals⁸² in later phases of infection when immunopathology is thought to dominate¹⁹ although there are as yet no data on such combinations. Rintatolimod was the only host-directed agent with *in vivo* efficacy data specifically for henipavirus infection identified from this review, providing only partial protection after low dose NiV-M challenge in hamsters⁵⁰.

Drug Evaluation

This paucity of drug candidates and high-quality evidence overall underscores the challenges of clinical development of therapeutics for rare but high-threat infections with pandemic potential.

Despite further Nipah outbreaks in the past year, there remain insufficient cases under the current epidemiological situation to obtain the human phase 3 RCT efficacy data necessary for licensure⁸³ or to attract substantial commercial investment. Alternative approaches similar to the regionally driven end-to-end West African Lassa fever Consortium⁸⁴ framework are needed¹⁹. The requirement for BSL-4 precautions for pre-clinical studies of NiV and HeV also restricts these to a small number of specialist facilities, few of which are located where Nipah or Hendra outbreaks have occurred.

Animal Studies

In the absence of outbreak RCTs, efficacy evaluation of Nipah and Hendra therapeutics is reliant on controlled animal challenge studies. The variable agreement between *in vitro* and *in vivo* efficacy results for most of the small molecules identified in this review emphasises the importance of animal efficacy data for clinical prioritisation.

The United States Food and Drug Administration allows for approval of drugs for conditions which threaten health security under the Animal Rule⁸⁵ when field trials are not possible, provided four criteria are met: 1) sufficient understanding of the pathophysiology of the condition and mechanism of its reduction by the product; 2) efficacy demonstrated in at least two animal species or one species which is a well-characterised model for predicting the product's response in humans; 3) animal study endpoint clearly related to the desired outcome in humans, typically reduction in mortality or major morbidity; and 4) PK and pharmacodynamic (PD) data from animals and humans supporting selection of an effective dose in humans. Anti-infective agents approved under the Animal Rule include: raxibacumab and obitoximab for anthrax, antibiotics like ciprofloxacin for plague, and tecovirimat and brincidofovir for smallpox. The European Medicines Agency has a similar Exceptional Circumstances⁸⁶ mechanism for granting marketing authorisation to medicines where collection of comprehensive efficacy and safety data under normal conditions of use is not possible.

The clinical and pathological features as well as the strengths and limitations of the major animal models in Nipah (AGMs⁸⁷, ferrets⁸⁸, and Syrian Golden Hamsters⁸⁹) have been reviewed⁹⁰ by the Coalition for Epidemic Preparedness Innovations (CEPI). AGMs are closest in physiology to humans

but have less consistent neurological symptoms than hamsters and ferrets⁹⁰. CEPI are also improving these models, particularly through standardising the virus challenge stock⁹⁰. Supporting access to NiV-B strains as well as standardisation of dose and route of challenge for each model could aid comparability across studies. For therapeutics studies, having uninfected controls to assess drug toxicity thresholds, ideally in the context of PK-PD studies to determine *in vivo* EC₅₀ values and concentration-efficacy relationships, is crucial for dose optimisation to derisk human studies.

Clinical Studies

Human data remain essential for safety evaluation⁸⁵. Phase 1 first-in-human safety data need to be collected for new therapeutic agents and existing experience from repurposed agents critically appraised for any potential exacerbation of adverse effects by the pathophysiology of Nipah or Hendra infection prior to deployment in outbreak settings.

Within an outbreak, these therapeutics should be evaluated in well-designed phase 2 clinical trials integrated and sustained in health systems⁹¹ using pre-approved standardised protocols maximising statistical and operational efficiency in assessment of internationally-agreed core outcome measures⁹². Where possible, drug concentrations should be measured at the same time points as efficacy and safety outcomes to characterise and quantify PK-PD relationships, including at different stages of disease. Where RCTs are not possible, observational studies employing enhanced clinical characterisation protocols⁹³ incorporating the same outcome measures could provide higher-quality observational data than is currently available⁹⁴. The long-term neurological sequelae of Nipah encephalitis⁹⁵⁻⁹⁷ also merit more systematic characterisation and potential inclusion as outcomes.

Outbreaks of high-threat infections invoke the ethical duty^{98,99} to conduct inclusive research with speed and rigor. Community and stakeholder engagement⁹⁹, including on design and interventions in trials, are key to support genuinely informed consent and maintain trust in the scientific process¹⁰⁰.

Frameworks & Tools

It is vital that potential therapeutics and their appropriate dosing regimens are selected, optimised, and stockpiled based on all available clinical and pre-clinical evidence well in advance of any outbreak. This continuous iterative process should be guided by disease-specific, and where appropriate product-specific, target product profiles¹⁰¹ (comprising indication, safety, efficacy, route, stability, and affordability characteristics) developed through consensus among all relevant stakeholders, including regulators, end users, and communities. Systems pharmacology, statistical, mathematical, and economic modelling are powerful tools to support decision-making by providing a formal framework for integration of (typically sparse) data from multiple study types, species, and diseases, as well as informing design efficiency of phase 1 and phase 2 RCTs.

Conclusion

At present, there is sufficient evidence to trial only m102.4 (mAb) and remdesivir (small molecule) (singly and/or in combination) for prophylaxis and early treatment of Nipah virus infection. In addition to well-designed RCTs, *in vivo* PK-PD studies to support drug selection and dose optimisation for all high-threat infections are needed.

4500 words

Contributors

XHSC, JD, PWH, and PLO conceptualised this review. XHSC wrote the study protocol and designed the search strategy with EH who conducted the literature search. XHSC piloted the review and screened studies with ILH and SL. XHSC, ILH, BJJC, MZH, JT, LMJ, and TPH extracted and tabulated data. MZH conducted the risk of bias assessment with BJJC. MZH and XHSC produced the figures. XHSC wrote the initial draft with ILH and BJJC. All authors critically reviewed, edited, and approved the final version of the manuscript.

Acknowledgements

XHSC is a United Kingdom (UK) National Institute of Health and Care Research (NIHR) Academic Clinical Lecturer in Infectious Diseases at the University of Oxford. MZH is a Moh Family Foundation Fellow at the Pandemic Sciences Institute, University of Oxford. JD is supported by the Moh Family Foundation, the NIHR Health Protection Research Unit in Emerging and Zoonotic Infections, and the UK Public Health Rapid Support Team Research Programme (grant number IS-RRT-1015-001). JT is supported by the Wellcome Trust (grant number 220211). PWH is the Moh Family Foundation Professor of Emerging Infections and Global Health at the Pandemic Sciences Institute, University of Oxford. PWH and PLO are supported by the UK Foreign and Commonwealth, and Development Office, the Wellcome Trust (grant number 215091/Z/18/Z), and the Bill & Melinda Gates Foundation (grant number OPP1209135).

The authors would like to thank all members of the Pandemic Sciences Institute Henipavirus Programme as well as delegates of the International Pandemic Sciences Conference 2023 and the Nipah Virus 25th Anniversary Symposium, both in Oxford, for illuminating discussions.

References

1. Goh KJ, Tan CT, Chew NK, et al. Clinical features of Nipah virus encephalitis among pig farmers in Malaysia. *N Engl J Med* 2000; **342**(17): 1229-35.
2. Luby SP, Rahman M, Hossain MJ, et al. Foodborne transmission of Nipah virus, Bangladesh. *Emerg Infect Dis* 2006; **12**(12): 1888-94.
3. Nikolay B, Salje H, Hossain MJ, et al. Transmission of Nipah Virus - 14 Years of Investigations in Bangladesh. *N Engl J Med* 2019; **380**(19): 1804-14.
4. Chowdhury S, Khan SU, Cramer G, et al. Serological evidence of henipavirus exposure in cattle, goats and pigs in Bangladesh. *PLoS Negl Trop Dis* 2014; **8**(11): e3302.
5. Islam A, Cannon DL, Rahman MZ, et al. Nipah Virus Exposure in Domestic and Peridomestic Animals Living in Human Outbreak Sites, Bangladesh, 2013-2015. *Emerg Infect Dis* 2023; **29**(2): 393-6.
6. Shete AM, Radhakrishnan C, Pardeshi PG, et al. Antibody response in symptomatic & asymptomatic Nipah virus cases from Kerala, India. *Indian J Med Res* 2021; **154**(3): 533-5.
7. Hossain MJ, Gurley ES, Montgomery JM, et al. Clinical presentation of Nipah virus infection in Bangladesh. *Clin Infect Dis* 2008; **46**(7): 977-84.
8. Sejvar JJ, Hossain J, Sana SK, et al. Long-term neurological and functional outcome in Nipah virus infection. *Annals of Neurology* 2007; **62**(3): 235-42.
9. Broder CC, Wong KT. Henipaviruses. *Neurotropic Viral Infections: Volume 1: Neurotropic RNA Viruses*: Springer International Publishing; 2016: 45-83.
10. Chua KB, Goh KJ, Wong KT, et al. Fatal encephalitis due to Nipah virus among pig-farmers in Malaysia. *Lancet* 1999; **354**(9186): 1257-9.
11. Paton NI, Leo Y, Zaki SR, et al. Outbreak of Nipah-virus infection among abattoir workers in Singapore. *Lancet* 1999; **354**(9186): 1253-6.
12. Chua KB. Epidemiology, surveillance and control of Nipah virus infections in Malaysia. *Malaysian Journal of Pathology* 2010; **32**(2): 69-73.
13. Suit BY, Hassan L, Krauss SE, et al. Mental Model of Malaysian Pig Farmers in Implementing Disease Prevention and Control Practices. *Frontiers in Veterinary Science* 2021; **8** (no pagination).
14. Ching PK, de los Reyes VC, Sucaldito MN, et al. Outbreak of henipavirus infection, Philippines, 2014. *Emerg Infect Dis* 2015; **21**(2): 328-31.
15. Chadha MS, Comer JA, Lowe L, et al. Nipah virus-associated encephalitis outbreak, Siliguri, India. *Emerg Infect Dis* 2006; **12**(2): 235-40.
16. Arunkumar G, Chandni R, Mourya DT, et al. Outbreak Investigation of Nipah Virus Disease in Kerala, India, 2018. *J Infect Dis* 2019; **219**(12): 1867-78.
17. Hassan MZ, Sazzad HMS, Luby SP, et al. Nipah virus contamination of hospital surfaces during outbreaks, Bangladesh, 2013-2014. *Emerging Infectious Diseases* 2018; **24**(1): 15-21.
18. World Health Organization. Disease Outbreak News: Nipah Virus Infection - India. 2023. <https://www.who.int/emergencies/disease-outbreak-news/item/2023-DON490> (accessed 12 February 2024).
19. Hassan MZ, Shirin T, Satter SM, et al. Nipah virus disease: what can we do to improve patient care? *Lancet Infect Dis* 2024.
20. Gomez Roman R, Tornieporth N, Cherian NG, et al. Medical countermeasures against henipaviruses: a review and public health perspective. *Lancet Infect Dis* 2022; **22**(1): e13-e27.
21. Mehand MS, Al-Shorbaji F, Millett P, Murgue B. The WHO R&D Blueprint: 2018 review of emerging infectious diseases requiring urgent research and development efforts. *Antiviral Res* 2018; **159**: 63-7.
22. Sterne JAC, Savovic J, Page MJ, et al. RoB 2: a revised tool for assessing risk of bias in randomised trials. *BMJ* 2019; **366**: l4898.
23. Sterne JA, Hernan MA, Reeves BC, et al. ROBINS-I: a tool for assessing risk of bias in non-randomised studies of interventions. *BMJ* 2016; **355**: i4919.
24. Hooijmans CR, Rovers MM, de Vries RB, Leenaars M, Ritskes-Hoitinga M, Langendam MW. SYRCLE's risk of bias tool for animal studies. *BMC Med Res Methodol* 2014; **14**: 43.

25. Sahay RR, Yadav PD, Gupta N, et al. Experiential learnings from the Nipah virus outbreaks in Kerala towards containment of infectious public health emergencies in India. *Epidemiol Infect* 2020; **148**: e90.
26. Playford EG, Munro T, Mahler SM, et al. Safety, tolerability, pharmacokinetics, and immunogenicity of a human monoclonal antibody targeting the g glycoprotein of henipaviruses in healthy adults: a first-in-human, randomised, controlled, phase 1 study. *Lancet Infectious Diseases* 2020; **20**(4): 445-54.
27. Mire CE, Satterfield BA, Geisbert JB, et al. Pathogenic Differences between Nipah Virus Bangladesh and Malaysia Strains in Primates: Implications for Antibody Therapy. *Scientific Reports* 2016; **6**.
28. Geisbert TW, Mire CE, Geisbert JB, et al. Therapeutic treatment of Nipah virus infection in nonhuman primates with a neutralizing human monoclonal antibody. *Science Translational Medicine* 2014; **6**(242) (no pagination).
29. Bossart KN, Geisbert TW, Feldmann H, et al. A Neutralizing Human Monoclonal Antibody Protects African Green Monkeys from Hendra Virus Challenge. *Science Translational Medicine* 2011; **3**(105).
30. Bossart KN, Zhu Z, Middleton D, et al. A neutralizing human monoclonal antibody protects against lethal disease in a new ferret model of acute Nipah virus infection. *PLoS Pathogens* 2009; **5**(10) (no pagination).
31. Zhu Z, Bossart KN, Bishop KA, et al. Exceptionally potent cross-reactive neutralization of Nipah and Hendra viruses by a human monoclonal antibody. *Journal of Infectious Diseases* 2008; **197**(6): 846-53.
32. Mire CE, Chan Y-P, Borisevich V, et al. A Cross-Reactive Humanized Monoclonal Antibody Targeting Fusion Glycoprotein Function Protects Ferrets Against Lethal Nipah Virus and Hendra Virus Infection. *Journal of Infectious Diseases* 2020; **221**: S471-S9.
33. Doyle MP, Kose N, Borisevich V, et al. Cooperativity mediated by rationally selected combinations of human monoclonal antibodies targeting the henipavirus receptor binding protein. *Cell Reports* 2021; **36**(9) (no pagination).
34. Dong J, Cross RW, Doyle MP, et al. Potent Henipavirus Neutralization by Antibodies Recognizing Diverse Sites on Hendra and Nipah Virus Receptor Binding Protein. *Cell* 2020; **183**(6): 1536-50.e17.
35. Guillaume V, Wong KT, Looi RY, et al. Acute Hendra virus infection: Analysis of the pathogenesis and passive antibody protection in the hamster model. *Virology* 2009; **387**(2): 459-65.
36. Guillaume V, Contamin H, Loth P, et al. Antibody prophylaxis and therapy against Nipah virus infection in hamsters. *Journal of Virology* 2006; **80**(4): 1972-8.
37. Playford EG, McCall B, Smith G, et al. Human Hendra virus encephalitis associated with equine outbreak, Australia, 2008. *Emerging Infectious Diseases* 2010; **16**(2): 219-23.
38. Banerjee S, Niyas VKM, Soneja M, et al. First experience of ribavirin postexposure prophylaxis for Nipah virus, tried during the 2018 outbreak in Kerala, India. *Journal of Infection* 2019; **78**(6): 491-503.
39. Kumar ASA, Sohanlal T, Prasad G, Gupta M, Gopal A. Single-center experience and lessons learnt from management of NIPAH virus outbreak in India. *Open Forum Infectious Diseases* 2019; **6**(Supplement 2): S659-S60.
40. Radhakrishnan C, Renjith TP, Arshad F, et al. Clinical manifestations of Nipah virus-infected patients who presented to the emergency department during an outbreak in Kerala state in India, May 2018. *Clinical Infectious Diseases* 2020; **71**(1): 152-7.
41. Warriar A. A single case outbreak of Nipah Encephalitis from India in May-June 2019. *International Journal of Infectious Diseases* 2020; **101**(Supplement 1): 247.
42. Chong HT, Kamarulzaman A, Tan CT, et al. Treatment of acute Nipah encephalitis with ribavirin. *Annals of Neurology* 2001; **49**(6): 810-3.

43. Lo MK, Feldmann F, Gary JM, et al. Remdesivir (GS-5734) protects African green monkeys from Nipah virus challenge. *Sci Transl Med* 2019; **11**(494).
44. Jordan R, Hogg A, Warren T, et al. Broad-spectrum investigational agent GS-5734 for the treatment of ebola, mers coronavirus and other pathogenic viral infections with high outbreak potential. *Open Forum Infectious Diseases* 2017; **4**(Supplement 1): S737.
45. Rockx B, Bossart KN, Feldmann F, et al. A novel model of lethal Hendra virus infection in African green monkeys and the effectiveness of ribavirin treatment. *Journal of Virology* 2010; **84**(19): 9831-9.
46. Mathieu C, Porotto M, Figueira TN, Horvat B, Moscona A. Fusion Inhibitory Lipopeptides Engineered for Prophylaxis of Nipah Virus in Primates. *J Infect Dis* 2018; **218**(2): 218-27.
47. Pallister J, Middleton D, Crameri G, et al. Chloroquine administration does not prevent nipah virus infection and disease in ferrets. *Journal of Virology* 2009; **83**(22): 11979-82.
48. Dawes BE, Kalveram B, Ikegami T, et al. Favipiravir (T-705) protects against Nipah virus infection in the hamster model. *Scientific Reports* 2018; **8**(1).
49. Freiberg AN, Worthy MN, Lee B, Holbrook MR. Combined chloroquine and ribavirin treatment does not prevent death in a hamster model of Nipah and Hendra virus infection. *Journal of General Virology* 2010; **91**(3): 765-72.
50. Georges-Courbot MC, Contamin H, Faure C, et al. Poly(I)-poly(C12U) but not ribavirin prevents death in a hamster model of Nipah virus infection. *Antimicrob Agents Chemother* 2006; **50**(5): 1768-72.
51. Lo MK, Spengler JR, Krumpke LRH, et al. Griffithsin inhibits Nipah virus entry and fusion and can protect Syrian golden hamsters from lethal Nipah virus challenge. (Special Issue: Twenty years of Nipah virus research.). *Journal of Infectious Diseases* 2020; **221**(Suppl. 4): S480-S92.
52. Mathieu C, Dhondt KP, Chalons M, et al. Heparan sulfate-dependent enhancement of henipavirus infection. *Mbio* 2015; **6**(2) (no pagination).
53. Mathieu C, Augusto MT, Niewiesk S, et al. Broad spectrum antiviral activity for paramyxoviruses is modulated by biophysical properties of fusion inhibitory peptides. *Scientific Reports* 2017; **7**.
54. Porotto M, Rockx B, Yokoyama CC, et al. Inhibition of Nipah Virus Infection In Vivo: Targeting an early stage of paramyxovirus fusion activation during viral entry. *PLoS Pathogens* 2010; **6**(10) (no pagination).
55. Welch SR, Spengler JR, Harmon JR, et al. Defective Interfering Viral Particle Treatment Reduces Clinical Signs and Protects Hamsters from Lethal Nipah Virus Disease. *Mbio* 2022; **13**(2): e0329421.
56. Wright PJ, Crameri G, Eaton BT. RNA synthesis during infection by Hendra virus: an examination by quantitative real-time PCR of RNA accumulation, the effect of ribavirin and the attenuation of transcription. *Archives of Virology* 2005; **150**(3): 521-32.
57. Porotto M, Orefice G, Yokoyama CC, et al. Simulating henipavirus multicycle replication in a screening assay leads to identification of a promising candidate for therapy. *Journal of Virology* 2009; **83**(10): 5148-55.
58. Lo MK, Jordan R, Arvey A, et al. GS-5734 and its parent nucleoside analog inhibit Filo-, Pneumo-, and Paramyxoviruses. *Scientific Reports* 2017; **7**.
59. Lo MK, Shrivastava-Ranjan P, Chatterjee P, et al. Broad-Spectrum In Vitro Antiviral Activity of ODBG-P-RVn: An Orally-Available, Lipid-Modified Monophosphate Prodrug of Remdesivir Parent Nucleoside (GS-441524). *Microbiology Spectrum* 2021; **9**(3) (no pagination).
60. Welch SR, Tilston NL, Lo MK, et al. Inhibition of Nipah virus by defective interfering particles. (Special Issue: Twenty years of Nipah virus research.). *Journal of Infectious Diseases* 2020; **221**(Suppl. 4): S460-S70.
61. Wolf MC, Freiberg AN, Zhang T, et al. A broad-spectrum antiviral targeting entry of enveloped viruses. *Proceedings of the National Academy of Sciences of the United States of America* 2010; **107**(7): 3157-62.

62. Whitmer SLM, Lo MK, Sazzad HMS, et al. Inference of Nipah virus evolution, 1999-2015. *Virus Evol* 2021; **7**(1): veaa062.
63. Chen C, Fang J, Chen S, et al. The efficacy and safety of remdesivir alone and in combination with other drugs for the treatment of COVID-19: a systematic review and meta-analysis. *BMC Infect Dis* 2023; **23**(1): 672.
64. Pharmaceuticals and Medical Devices Agency. Avigan (Favipiravir) Review Report. Tokyo, Japan, 2014.
65. Nguyen TH, Guedj J, Anglaret X, et al. Favipiravir pharmacokinetics in Ebola-Infected patients of the JIKI trial reveals concentrations lower than targeted. *PLoS Negl Trop Dis* 2017; **11**(2): e0005389.
66. Wang Y, Zhong W, Salam A, et al. Phase 2a, open-label, dose-escalating, multi-center pharmacokinetic study of favipiravir (T-705) in combination with oseltamivir in patients with severe influenza. *EBioMedicine* 2020; **62**: 103125.
67. Irie K, Nakagawa A, Fujita H, et al. Population pharmacokinetics of favipiravir in patients with COVID-19. *CPT Pharmacometrics Syst Pharmacol* 2021; **10**(10): 1161-70.
68. Madelain V, Guedj J, Mentre F, et al. Favipiravir Pharmacokinetics in Nonhuman Primates and Insights for Future Efficacy Studies of Hemorrhagic Fever Viruses. *Antimicrob Agents Chemother* 2017; **61**(1).
69. Bixler SL, Bocan TM, Wells J, et al. Efficacy of favipiravir (T-705) in nonhuman primates infected with Ebola virus or Marburg virus. *Antiviral Res* 2018; **151**: 97-104.
70. Pilkington V, Pepperrell T, Hill A. A review of the safety of favipiravir - a potential treatment in the COVID-19 pandemic? *J Virus Erad* 2020; **6**(2): 45-51.
71. Batool S, Vuthaluru K, Hassan A, et al. Efficacy and Safety of Favipiravir in Treating COVID-19 Patients: A Meta-Analysis of Randomized Control Trials. *Cureus* 2023; **15**(1): e33676.
72. Cheng HY, French CE, Salam AP, et al. Lack of Evidence for Ribavirin Treatment of Lassa Fever in Systematic Review of Published and Unpublished Studies(1). *Emerg Infect Dis* 2022; **28**(8): 1559-68.
73. Salam AP, Duvignaud A, Jaspard M, et al. Ribavirin for treating Lassa fever: A systematic review of pre-clinical studies and implications for human dosing. *PLoS Negl Trop Dis* 2022; **16**(3): e0010289.
74. Government of India Ministry of Health and Family Welfare. Clinical Management Protocol for Nipah Virus Disease. 2018.
75. Government of Kerala Department of Health and Family Welfare. Nipah Virus Infection Guidelines for Surveillance, Diagnosis, Treatment, Prevention and Control. 2021.
76. Government of the People's Republic of Bangladesh Ministry of Health and Family Welfare. National Guideline for Management, Prevention and Control of Nipah Virus Infection including Encephalitis. 2018.
77. White NJ, Watson JA, Hoglund RM, Chan XHS, Cheah PY, Tarning J. COVID-19 prevention and treatment: A critical analysis of chloroquine and hydroxychloroquine clinical pharmacology. *PLoS Med* 2020; **17**(9): e1003252.
78. Haeusler IL, Chan XHS, Guerin PJ, White NJ. The arrhythmogenic cardiotoxicity of the quinoline and structurally related antimalarial drugs: a systematic review. *BMC Med* 2018; **16**(1): 200.
79. Drell W, Welch AD. Azaribine-homocystinemia-thrombosis in historical perspective. *Pharmacol Ther* 1989; **41**(1-2): 195-206.
80. Oey A, McClure M, Symons JA, et al. Lumicitabine, an orally administered nucleoside analog, in infants hospitalized with respiratory syncytial virus (RSV) infection: Safety, efficacy, and pharmacokinetic results. *PLoS One* 2023; **18**(7): e0288271.
81. de Wit E, Williamson BN, Feldmann F, et al. Late remdesivir treatment initiation partially protects African green monkeys from lethal Nipah virus infection. *Antiviral Res* 2023; **216**: 105658.
82. Kaufmann SHE, Dorhoi A, Hotchkiss RS, Bartenschlager R. Host-directed therapies for bacterial and viral infections. *Nat Rev Drug Discov* 2018; **17**(1): 35-56.

83. Nikolay B, Ribeiro Dos Santos G, Lipsitch M, et al. Assessing the feasibility of Nipah vaccine efficacy trials based on previous outbreaks in Bangladesh. *Vaccine* 2021; **39**(39): 5600-6.
84. ISARIC. West Africa Lassa fever Consortium (WALC). 2023. <https://isaric.org/research-2/lassa-fever-resources/walc/> (accessed 10 January 2024).
85. United States Food and Drug Administration. Animal Rule Approvals. 2002. <https://www.fda.gov/drugs/nda-and-bla-approvals/animal-rule-approvals> (accessed 4 January 2024).
86. European Medicines Agency. Exceptional Circumstances. 2024. <https://www.ema.europa.eu/en/glossary/exceptional-circumstances> (accessed 10 January 2024).
87. Mire CE, Satterfield BA, Geisbert TW. Nonhuman Primate Models for Nipah and Hendra Virus Countermeasure Evaluation. *Methods Mol Biol* 2023; **2682**: 159-73.
88. Rockx B, Mire CE. Ferret Models for Henipavirus Infection. *Methods Mol Biol* 2023; **2682**: 205-17.
89. Juelich T, Smith J, Freiberg AN. Syrian Golden Hamster Model for Nipah Virus Infection. *Methods Mol Biol* 2023; **2682**: 219-29.
90. Centre for Epidemic Preparedness Innovations. Nipah Virus Assays and Animal Models for Vaccine Development, 2021.
91. Moorthy V, Abubakar I, Qadri F, et al. The future of the global clinical trial ecosystem: a vision from the first WHO Global Clinical Trials Forum. *Lancet* 2023.
92. Williamson PR, Altman DG, Blazeby JM, et al. Developing core outcome sets for clinical trials: issues to consider. *Trials* 2012; **13**: 132.
93. ISARIC Clinical Characterisation Protocol Investigators Group. ISARIC/WHO Clinical Characterisation Protocol. 2023. <https://isaric.net/ccp/> (accessed 9 January 2024).
94. Dunning JW, Merson L, Rohde GGU, et al. Open source clinical science for emerging infections. *Lancet Infect Dis* 2014; **14**(1): 8-9.
95. Lee KE, Umapathi T, Tan CB, et al. The neurological manifestations of Nipah virus encephalitis, a novel paramyxovirus. *Ann Neurol* 1999; **46**(3): 428-32.
96. Tan CT, Goh KJ, Wong KT, et al. Relapsed and late-onset Nipah encephalitis. *Ann Neurol* 2002; **51**(6): 703-8.
97. Sejvar JJ, Hossain J, Saha SK, et al. Long-term neurological and functional outcome in Nipah virus infection. *Ann Neurol* 2007; **62**(3): 235-42.
98. Johnson T, Jamrozik E, Hurst T, Cheah PY, Parker MJ. Ethical issues in Nipah virus control and research: addressing a neglected disease. *J Med Ethics* 2023.
99. World Health Organization. Emergency use of unproven clinical interventions outside clinical trials: ethical considerations, 2022.
100. Meskell P, Biesty LM, Dowling M, et al. Factors that impact on recruitment to vaccine trials in the context of a pandemic or epidemic: a qualitative evidence synthesis. *Cochrane Database Syst Rev* 2023; **9**(9): MR000065.
101. World Health Organization. Target Product Profiles. 2023. <https://www.who.int/observatories/global-observatory-on-health-research-and-development/analyses-and-syntheses/target-product-profile/who-target-product-profiles> (accessed 5 January 2024).
102. British National Formulary. Remdesivir. 2024. <https://bnf.nice.org.uk/drugs/remdesivir/> (accessed 6 January 2024).

Table 1: Nipah and Hendra Virus Therapeutic Monoclonal Antibodies

Drug (mechanism)	Developer	Reference	Study Design	Funder
m102.4 (anti-HeV-G)	Uniformed Services University, USA	Sahay 2020 ²⁵	Clinical: compassionate use post-exposure prophylaxis during Nipah outbreak in Kerala, India (n=1)	USA NIH
		Playford 2020 ²⁶	Clinical: healthy adult volunteers (18-50 years) phase 1 dose-escalation RCT for safety, tolerability, and pharmacokinetics in Brisbane, Australia (n=40)	
		Mire 2016 ²⁷	Animal: AGM challenge with NiV-B for efficacy and safety (n=11) <ul style="list-style-type: none"> 2.5 x 10⁵ PFU intratracheal + 2.5 x 10⁵ PFU intranasal 	
		Geisbert 2014 ²⁸	Animal: AGM challenge with NiV-M for efficacy and safety (n=16) <ul style="list-style-type: none"> 5 x 10⁵ PFU intratracheal 	
		Bossart 2011 ²⁹	Animal: AGM challenge with HeV for efficacy and safety (n=14) <ul style="list-style-type: none"> 4 x 10⁵ TCID₅₀ intratracheal Animal: AGM pharmacokinetics (n=4)	
		Bossart 2009 ³⁰	Animal: Ferret challenge with NiV-M for efficacy (n=8) <ul style="list-style-type: none"> 5 x 10³ TCID₅₀ oronasal 	
		Zhu 2008 ³¹	Animal: Ferret pharmacokinetics (n=4)	
h5B3.1 (anti-NiV-F)		Mire 2020 ³²	Animal: Ferret challenge with NiV-M or HeV for efficacy (n=11) <ul style="list-style-type: none"> 5 x 10³ PFU intranasal 	
HENV-103, HENV-117, HENV-58, HENV-98, HENV-100 (anti-HeV-RBP)	Vanderbilt University, USA	Doyle 2021 ³³	Animal: Hamster challenge with NiV-B for efficacy <ul style="list-style-type: none"> 5 x 10⁶ PFU intranasal 	USA NIH
HENV-26, HENV-32 (anti-HeV-RBP)		Dong 2020 ³⁴	Animal: Ferret challenge with NiV-B for efficacy (n=13) <ul style="list-style-type: none"> 5 x 10³ PFU intranasal 	
NipGIP1.7 & Nip3B10 (anti-NiV-G), NipGIP35 & NipGIP3 (anti-NiV-F)	INSERM, France	Guillaume 2006 ³⁶	Animal: Hamster challenge with NiV-M for efficacy, dose titration, and therapeutic time window (n=124) <ul style="list-style-type: none"> 7.5 x 10² PFU (100 LD₅₀) intraperitoneal 	Aventis Pharma, Bayer Pharma, INSERM & Institut Pasteur
NipGIP35, NipGIP3, NipGIP21, NipGIP7 (anti-NiV-F)		Guillaume 2009 ³⁵	Animal: Hamster challenge with HeV for efficacy and dose titration (n=54) <ul style="list-style-type: none"> 10³ PFU (100 LD₅₀) intraperitoneal 	

AGM = African Green monkey; HeV = Hendra virus; INSERM = Institut National de la Santé et de la Recherche Médicale; LD₅₀ = median lethal dose; NIH = National Institutes of Health; NiV-B = Nipah virus Bangladesh; NiV-M = Nipah virus Malaysia; PFU = plaque-forming units; RBP = receptor binding protein; RCT = randomised controlled trial; TCID₅₀ = median tissue culture infectious dose; USA = United States of America. Please see Supplementary Table I for full table.

Table 2: Nipah and Hendra Virus Therapeutic Small Molecules (Clinical and Animal Studies)

Drug (mechanism)	Reference	Study Design	Funder
Ribavirin (nucleoside analogue prodrug)	Warrier 2020 ⁴¹	Clinical: compassionate use for treatment in Nipah outbreak in Kochi, India, 2019 (n=1)	N/A
	Radhakrishnan 2020 ⁴⁰	Clinical: compassionate use for treatment in Nipah outbreak in Kerala, India, 2018 (n=12: 6 treated, 6 untreated)	
	Banerjee 2019 ³⁸	Clinical: compassionate use for post-exposure prophylaxis of healthcare workers during Nipah outbreak in Kerala, India, 2018 (n=8)	
	Kumar 2019 ³⁹	Clinical: compassionate use for treatment in Nipah outbreak in Kerala, India, 2018 (n=5)	
	Playford 2010 ³⁷	Clinical: compassionate use during Hendra outbreak in Australia, 2008 for treatment (n=2) and post-exposure prophylaxis (n=1)	
	Chong 2001 ⁴²	Clinical: compassionate use for treatment in Nipah outbreak in Malaysia, 1998-99 (n=194: 140 treated, 54 untreated)	
	Rockx 2010 ⁴⁵	Animal: AGM challenge with HeV (n=12) • 4 x 10 ⁵ TCID ₅₀ intratracheal for efficacy	
Ribavirin (nucleoside analogue prodrug) & 6-azauridine (OMP decarboxylase inhibitor) & Rintatolimod (TLR-3 agonist interferon inducer)	Georges-Courbot 2006 ⁵⁰	Animal: Hamster challenge with NiV-M for efficacy (n=18) • 350 x LD ₅₀ intraperitoneal Animal: Hamster challenge with NiV-M for efficacy (n=18) • 35 x LD ₅₀ intraperitoneal	N/A
Ribavirin (nucleoside analogue prodrug) & chloroquine (lysosome alkalisation)	Freiberg 2010 ⁴⁹	Animal: Hamster challenge with NiV-M (n=41) and HeV (n=20) for efficacy (n=85) • 10 ⁴ TCID ₅₀ intraperitoneal	USA NIH
Chloroquine (lysosome alkalisation)	Pallister 2009 ⁴⁷	Animal: Ferret challenge with NiV-M (n=8) for efficacy and pharmacokinetics • 5 x 10 ³ TCID ₅₀ oronasal	USA NIH
Remdesivir (nucleoside analogue)	Lo 2019 ⁴³	Animal: AGM challenge with NiV-B for efficacy (n=8) • 10 ⁵ TCID ₅₀ intranasal + 10 ⁵ TCID ₅₀ intratracheal	USA NIH
	Jordan 2017 ⁴⁴	Animal: AGM challenge with NiV-B for efficacy • Lethal dose (unspecified)	
Favipiravir (nucleoside analogue prodrug)	Dawes 2018 ⁴⁸	Animal: Hamster challenge with NiV-M for efficacy (n=18) • 10 ⁴ PFU intraperitoneal	USA NIH
Griffithsin (GRFT) (fusion and cell entry inhibitor)	Lo 2020 ⁵¹	Animals: Hamster challenge with NiV-B for efficacy (n=65) • 10 ⁷ TCID ₅₀ intranasal	USA NIH & USA CDC
Periodate heparin (competitive inhibitor of <i>trans</i> -infection)	Mathieu 2015 ⁵²	Animal: Hamster challenge with NiV-M for efficacy (n=15) • 500 x LD ₅₀ intraperitoneal	INSERM
Fusion inhibitory lipopeptides (fusion and cell entry inhibitors): VIKI-dPEG4-Chol, VIKI-dPEG4-Toco VG-PEG24-chol VIKI-PEG4-chol	Mathieu 2018 ⁴⁶	Animal: Hamster challenge with NiV-M for efficacy (n=38) • 10 ⁵ PFU (100 x LD ₅₀) intranasal Animal: AGM challenge with NiV-M for efficacy (n=10) • 2 x 10 ⁷ PFU intratracheal Animal: AGM biodistribution (n=4)	USA NIH & INSERM
	Mathieu 2017 ⁵³	Animal: Hamster challenge with NiV-M for efficacy (n=13) • 100 x LD ₅₀ intraperitoneal Animal: Hamster biodistribution (n=6)	
	Porotto 2010 ⁵⁴	Animal: Hamster challenge with NiV (strain unspecified) for efficacy (n=35) • 100 x LD ₅₀ intraperitoneal	
Defective interfering particles (virus-like particles containing defective interfering genomes which inhibit replication): DI-07, DI-10, DI-14, DI-35	Welch 2022 ⁵⁵	Animal: Hamster challenge with NiV-M for efficacy (n=153) • Experiment 1: 10 ⁴ TCID ₅₀ intraperitoneal • Experiment 2: 10 ⁶ TCID ₅₀ intranasal	USA CDC
Ceftriaxone (bacterial cell wall synthesis inhibitor), clarithromycin (bacterial protein synthesis inhibitor), aciclovir (nucleoside analogue)	Paton 1999 ¹¹	Clinical: empirical syndromic treatment during outbreak in Singapore, 1999 (n=11)	N/A

AGM = African Green monkey; CDC = Centres for Disease Control; HeV = Hendra virus; dPEG = discrete Polyethylene Glycol; INSERM = Institut National de la Santé et de la Recherche Médicale; LD₅₀ = median lethal dose; NIH = National Institutes of Health; NiV-B = Nipah virus Bangladesh; NiV-M = Nipah virus Malaysia; OMP = orotidine monophosphate; PFU = plaque-forming units; TCID₅₀ = median tissue culture infectious dose; TLR-3 = toll-like receptor 3; USA = United States of America. Please see Supplementary Table II for full table.

Table 3: Nipah and Hendra Therapeutic Candidates Clinical Prioritisation

Type	Drug	Efficacy	Safety	Feasibility	Clinical Prioritisation & Proposed Further Evaluation
mAb (see also Table 1)	m102.4 (anti-HeV-G)	Protected monkeys from death and pathology given at two doses 48 hours apart starting on day 1 or 3 after NiV-B ²⁷ (n=6) or on day 1, 3, or 5 after NiV-M ²⁸ (n=12) and HeV ²⁹ (n=12) challenge; also protected ferrets from death given 10 hours after NiV-M ³⁰ (n=3) challenge	No SAEs, similar rate of mild TEAEs between treatment and placebo groups, and no anti-m102.4 antibodies in phase 1 RCT in healthy adults (n=30 treated) ²⁶	High cost of goods ²⁰ , limited drug supply, parenteral route only.	Phase 2a post-exposure prophylaxis and/or early treatment RCT during Nipah or Hendra outbreak Shorter treatment window for NiV-B than NiV-M Dose optimisation for cost recommended
	h5B.3 (anti-NiV-F)	Protected ferrets from death but not minor clinical signs given at 20mg/kg in two doses 48 hours apart starting on day 1 or 3 after NiV-M (n=6) or HeV (n=3) challenge ³²	No human studies to date. No safety data reported in animal studies.		Monkey studies with NiV-B challenge Dose optimisation for cost necessary
	HENV-26 (anti-HeV RBP)	Protected ferrets (n=5) from death, symptoms, and viraemia given at 15mg/kg on days 3 and 5 after NiV-B challenge ³⁴			
	HENV-103 + HENV-117 (anti-HeV RBP)	HENV-103 + HENV-117 cocktail (5mg/kg each) protected hamsters (n=5) from death given a day after NiV-B challenge ³³			
Small molecule (see also Table 2)	Remdesivir (nucleoside analogue)	Fully protected monkeys (n=4) from death but not mild respiratory signs (n=2) or focal meningo-encephalitis (n=1) given daily from a day after NiV-B challenge ⁴³ ; EC ₅₀ values 0.029-0.066µM <i>in vitro</i> ⁵⁸	Sinus bradycardia and hepatotoxicity have been observed in humans ¹⁰²	Approved globally for COVID-19. Less affordable ²⁰ , parenteral route with oral formulation in development ⁵⁹ .	Phase 2a post-exposure prophylaxis and/or early treatment RCT during Nipah or Hendra outbreak Dose optimisation for efficacy recommended
	Favipiravir (nucleoside analogue)	Fully protected hamsters (n=10) from death, symptoms, and pathology given daily from time of NiV-M challenge; EC ₅₀ values 11-44µM <i>in vitro</i> ⁴⁸	Lethal toxicity in dogs and monkeys (>1g/kg) ⁶⁴ , teratogenicity across four animal species ⁶⁴ , transient hyperuricaemia in humans ^{64,70}	Approved in Japan for novel influenza ⁶⁴ . Affordable ²⁰ , oral route, but non-linear PK complicates dosing ⁶⁷ .	Monkey study with NiV-B challenge Dose optimisation for efficacy necessary
	Ribavirin (nucleoside analogue)	Delayed time to but did not prevent death given from before or within 12 hours to monkeys (n=6) after HeV ⁴⁵ and hamsters (n=17) after NiV-M ^{49,50} challenge; IC ₅₀ values 4.2-5.0µM <i>in vitro</i> ⁴⁹	Dose-dependent toxicity in hamsters (>100mg/kg) ⁴⁹ and humans ³⁸ limits safety and tolerability	Approved globally for chronic hepatitis C. Affordable ²⁰ but equivocal risk:benefit.	Monkey study with NiV-B challenge Dose optimisation for safety and efficacy critical Time-to-event outcome measure if in phase 2a RCT
	Chloroquine (4-amino- quinoline)	Did not protect ferrets ⁴⁷ (n=6) and hamsters ⁴⁹ (n=19) from death as monotherapy given from before or within 12 hours after NiV-M or HeV challenge	Dose-dependent lethal toxicity in hamsters (>100mg/kg) ⁴⁹ and humans (>3µM plasma) ⁷⁷	Approved globally for malaria. Affordable ²⁰ but unfavourable risk:benefit.	Should not be used for the prophylaxis or treatment of Nipah or Hendra virus infection

EC₅₀ = 50% maximal effective concentration; HeV = Hendra virus; IC₅₀ = 50% maximal inhibitory concentration; mAb = monoclonal antibody; NiV-B = Nipah virus Bangladesh; NiV-M = Nipah virus Malaysia; PK = pharmacokinetics; RBP = receptor binding protein; RCT = randomised controlled trial; SAE = serious adverse event; TEAE = treatment emergent adverse event. Colours represent level of clinical prioritisation: green = high priority, ample evidence for efficacy and safety for phase 2a trial; orange = intermediate priority, further evidence required for efficacy and/or potential major limitations in safety and/or feasibility; red = low priority, no evidence for efficacy and major limitations in safety and/or feasibility

Characteristic	Monoclonal Antibodies	Small Molecules
Efficacy	Highly specific	Less specific
Safety, including pregnancy/lactation and other immunosuppression	Specificity confers low potential for off-target adverse effects. Typically safe in pregnancy/lactation and immunosuppression.	Higher potential for off-target adverse effects. Hepatotoxicity, cardiotoxicity, and teratogenicity often concerns.
Duration of effect	Half-life extending mutations can prolong protection from 1 month to 3-6 months	Elimination half-lives of leading nucleoside analogue candidates between 1-5 hours
Route of administration	Parenteral	Parenteral and/or oral
Pharmacokinetic profile	Typically linear	Can be linear or non-linear
Cost	Higher (USD 1000-2000)	Lower (USD 10 to 1000)
Development Timeline	Shorter	Longer

Table 4: Comparative Advantages of Monoclonal Antibodies and Small Molecules

Figure 1: PRISMA Flow Chart

