

Table S1. Usual nicotine strength used by adult (≥16y) vapers in Great Britain, January 2022-January 2024 – refillable device users

	<i>N</i> ¹	Nicotine strength, % [95% CI] ²						Do not know if it contains nicotine	Contains nicotine but do not know the strength
		No nicotine	6 mg/ml or less	7 to 11 mg/ml	12 to 19 mg/ml	20 mg/ml or more			
All adult vapers (≥16y)	1267	11.8 [9.8–13.8]	39 [35.8–42.1]	13.2 [11.0–15.4]	20.2 [17.6–22.8]	11.5 [9.4–13.6]	0.9 [0.3–1.5]	3.5 [2.4–4.6]	
Age (years)									
16-24	159	15.3 [9.3–21.2]	25.6 [17.7–33.6]	15.3 [9.1–21.4]	20.2 [12.5–27.9]	20.4 [13.3–27.4]	1.6 [0–4.0]	1.7 [0–4.0]	
25-34	284	10.9 [6.8–15.0]	43.1 [36.4–49.7]	15.1 [10.3–20.0]	12.7 [8.2–17.2]	13.5 [9.0–17.9]	1.5 [0–3.1]	3.2 [0.8–5.6]	
35-44	226	12.3 [7.2–17.3]	43.6 [36.2–51.0]	15.2 [9.9–20.6]	19.2 [13.2–25.2]	8.8 [4.6–13.1]	0 [0–0]	0.9 [0–1.9]	
45-54	253	7.7 [4.2–11.2]	44.8 [37.6–52.0]	10.4 [5.6–15.1]	22.9 [16.8–29.0]	7.5 [3.3–11.8]	0.8 [0–1.9]	5.9 [2.3–9.4]	
55-64	219	12.2 [7.3–17.1]	35.6 [28.3–42.9]	10.9 [6.5–15.3]	27.6 [20.8–34.5]	8.9 [4.0–13.8]	0.5 [0–1.5]	4.3 [1.7–6.9]	
≥65	125	15.0 [7.5–22.5]	30.0 [20.5–39.6]	7.4 [2.0–12.8]	29.8 [20.1–39.6]	8.8 [2.3–15.4]	0.7 [0–2.0]	8.2 [3.2–13.3]	
Gender ³									
Men	712	11.8 [9.1–14.4]	39.0 [34.8–43.2]	12.6 [9.7–15.4]	19.1 [15.8–22.5]	12.9 [9.9–15.9]	1.0 [0–1.9]	3.7 [2.1–5.3]	
Women	538	11.8 [8.7–15.0]	39.2 [34.4–44.0]	14.2 [10.7–17.7]	21.3 [17.1–25.5]	9.6 [6.7–12.5]	0.6 [0–1.3]	3.3 [1.8–4.7]	
Occupational social grade									
ABC1 (more advantaged)	751	11.1 [8.7–13.6]	40.2 [36.3–44.1]	13.6 [10.8–16.3]	19.5 [16.4–22.7]	11.3 [8.7–13.9]	0.9 [0.2–1.6]	3.4 [1.9–4.9]	
C2DE (less advantaged)	516	12.3 [9.2–15.5]	38.0 [33.2–42.8]	12.9 [9.5–16.2]	20.7 [16.7–24.7]	11.7 [8.4–14.9]	0.9 [0–1.8]	3.5 [2.0–5.1]	
Nation									
England	907	11.2 [9.1–13.6]	39.2 [35.7–42.7]	13.6 [11.3–16.3]	19.9 [17.1–23.0]	11.9 [9.7–14.5]	1 [0.5–2]	3.3 [2.2–4.7]	
Wales	117	18.8 [12.2–27.9]	31.0 [22.5–41.0]	11.2 [6.1–19.6]	22.9 [15.7–32.3]	10.6 [5.9–18.3]	0 [0–0]	5.4 [2.3–12.0]	
Scotland	243	14.6 [10.1–20.6]	41.4 [34.8–48.3]	9.8 [6.2–15.2]	21.9 [16.7–28.2]	7.6 [4.6–12.3]	0 [0–0]	4.8 [2.4–9.3]	
History of ≥1 diagnosed mental health conditions ⁴									
No	412	10.9 [7.6–14.1]	42.3 [37.0–47.6]	10.8 [7.6–14.0]	19.9 [15.5–24.2]	10.4 [7.0–13.8]	1.3 [0.1–2.5]	4.4 [2.3–6.4]	
Yes	389	11.8 [8.5–15.0]	40.8 [35.3–46.3]	11.7 [8.1–15.3]	21.2 [16.5–25.9]	10.0 [6.7–13.3]	1.1 [0–2.1]	3.5 [1.6–5.5]	
Smoking status									
Long-term (≥1y) ex-smoker	587	12.0 [9.0–15.0]	45.0 [40.3–49.8]	12.9 [9.6–16.1]	21.3 [17.3–25.3]	6.5 [4.1–9.0]	0.5 [0–1.4]	1.7 [0.3–3.1]	
Recent (<1y) ex-smoker	108	12.2 [5.8–18.5]	35.5 [24.9–46.1]	16.2 [7.4–25.0]	20.8 [11.9–29.7]	11.9 [5.2–18.5]	1.6 [0–3.8]	1.9 [0–4.7]	
Current smoker	459	10.1 [7.1–13.2]	34.1 [29.0–39.2]	12.5 [9.0–16.1]	18.9 [14.8–22.9]	16.0 [11.8–20.1]	1.2 [0–2.4]	7.1 [4.7–9.6]	
Never smoker	113	16.0 [7.8–24.1]	32.5 [22.1–42.9]	14.4 [7.2–21.5]	19.4 [10.7–28.1]	17.2 [9.5–24.9]	0.6 [0–1.8]	0 [0–0]	

¹ Unweighted sample size.² Weighted row percentages.³ Participants who described their gender in another way were excluded from analyses by gender due to low numbers.⁴ Mental health conditions were not collected after June 2023, so results are based on aggregated data across January 2022 – June 2023.

Table S2. Usual nicotine strength used by adult (≥16y) vapers in Great Britain, January 2022-January 2024 – disposable device users

	N ¹	Nicotine strength, % [95% CI] ²						Do not know if it contains nicotine	Contains nicotine but do not know the strength
		No nicotine	6 mg/ml or less	7 to 11 mg/ml	12 to 19 mg/ml	20 mg/ml or more			
All adult vapers (≥16y)	804	3.7 [2.4–5.0]	23.3 [20.0–26.6]	3.5 [2.1–5.0]	5.3 [3.5–7.1]	47.9 [44.0–51.8]	1.2 [0.3–2.1]	15.1 [12.4–17.8]	
Age (years)									
16-24	341	1.2 [0.2–2.2]	21.1 [16.0–26.1]	3.1 [1.1–5.1]	5.3 [2.4–8.1]	58.1 [52.1–64.0]	1.2 [0–2.5]	10.1 [6.6–13.6]	
25-34	199	4.6 [1.6–7.6]	25.7 [18.8–32.5]	3.8 [0.7–6.9]	4.5 [1.3–7.6]	44.1 [36.1–52.0]	1.4 [0–3.7]	16.0 [10.4–21.6]	
35-44	112	6.7 [1.7–11.6]	24.8 [15.3–34.3]	5.3 [0.1–10.6]	5.3 [0.6–10.0]	43.0 [32.5–53.5]	1.1 [0–2.8]	13.8 [6.8–20.8]	
45-54	77	4.4 [0–8.8]	24.6 [14.3–34.9]	1.8 [0–4.5]	7.5 [0.6–14.4]	38.7 [25.7–51.8]	0 [0–0]	23.1 [11.6–34.6]	
55-64	50	11.4 [2.0–20.8]	19.5 [7.9–31.1]	3.3 [0–8.2]	4.4 [0–10.5]	25.9 [11.4–40.5]	1.3 [0–4.0]	34.1 [19.4–48.9]	
≥65	25	2.5 [0–7.6]	35.3 [10.1–60.4]	4.9 [0–14.6]	10.3 [0–24.5]	16.4 [0.1–32.7]	3.3 [0–8.2]	27.5 [8.4–46.5]	
Gender ³									
Men	372	2.3 [0.9–3.8]	21.8 [17.1–26.5]	5.1 [2.6–7.7]	6.1 [3.3–9.0]	46.6 [40.7–52.4]	1.1 [0–2.2]	16.9 [12.6–21.2]	
Women	416	4.9 [2.8–7.0]	24.9 [20.1–29.8]	2.1 [0.6–3.7]	4.3 [2.1–6.5]	49.0 [43.6–54.5]	1.3 [0–2.7]	13.3 [9.9–16.8]	
Occupational social grade									
ABC1 (more advantaged)	475	4.6 [2.6–6.6]	21.9 [17.9–25.9]	2.7 [1.1–4.3]	5.2 [2.9–7.5]	48.0 [43.2–52.9]	0.6 [0–1.3]	17.0 [13.3–20.6]	
C2DE (less advantaged)	329	3.0 [1.3–4.8]	24.5 [19.3–29.6]	4.2 [1.9–6.5]	5.4 [2.8–8.0]	47.8 [41.8–53.8]	1.7 [0.2–3.2]	13.5 [9.6–17.4]	
Nation									
England	636	3.6 [2.5–5.3]	23.8 [20.4–27.6]	3.4 [2.2–5.4]	5.5 [3.9–7.8]	47.7 [43.5–52.0]	1.1 [0.5–2.6]	14.7 [12.1–17.9]	
Wales	51	4.0 [0.9–16.4]	16.3 [8.4–29.3]	2.5 [0.6–10.5]	4.7 [1.0–19.6]	59.5 [44.1–73.1]	4.8 [1.3–16.1]	8.0 [3.4–17.7]	
Scotland	117	4.8 [2.0–11.1]	19.1 [12.3–28.4]	4.8 [2.0–11.4]	2.5 [0.9–7.0]	45.5 [35.9–55.5]	0.6 [0.1–3.9]	22.6 [15.1–32.4]	
History of ≥1 diagnosed mental health conditions ⁴									
No	219	5.1 [2.1–8.0]	24.6 [18.2–31.1]	5.1 [1.8–8.4]	4.3 [1.4–7.3]	44.0 [36.7–51.3]	0.7 [0–1.8]	16.1 [10.8–21.3]	
Yes	258	2.8 [0.9–4.7]	24.2 [18.7–29.8]	4.5 [1.7–7.3]	6.0 [2.7–9.4]	51.8 [45.1–58.4]	2 [0.3–3.6]	8.7 [5.0–12.3]	
Smoking status									
Long-term (≥1y) ex-smoker	105	5.9 [1.3–10.4]	19.7 [11.4–27.9]	4.2 [0.3–8.2]	7.5 [1.4–13.6]	50.4 [39.5–61.3]	0.3 [0–0.8]	12.1 [5.1–19.0]	
Recent (<1y) ex-smoker	86	3.1 [0–6.3]	26.9 [16.1–37.7]	0 [0–0]	2.1 [0–5.5]	52.2 [39.7–64.7]	2.7 [0–8.0]	12.9 [4.8–21.1]	
Current smoker	464	3.5 [1.7–5.3]	24.9 [20.3–29.5]	4.2 [2.0–6.3]	6.5 [4–9.1]	44.0 [38.9–49.2]	0.7 [0–1.4]	16.2 [12.5–19.8]	
Never smoker	149	3.3 [0.6–5.9]	18.8 [11.8–25.7]	3.1 [0.4–5.8]	1.9 [0–4.6]	55.3 [46.4–64.1]	2.5 [0–5.1]	15.2 [8.8–21.5]	

¹ Unweighted sample size.² Weighted row percentages.³ Participants who described their gender in another way were excluded from analyses by gender due to low numbers.⁴ Mental health conditions were not collected after June 2023, so results are based on aggregated data across January 2022 – June 2023.

Table S3. Usual nicotine strength used by adult (≥16y) vapers in Great Britain, January 2022-January 2024 – pod device users

	N ¹	Nicotine strength, % [95% CI] ²						Do not know if it contains nicotine	Contains nicotine but do not know the strength
		No nicotine	6 mg/ml or less	7 to 11 mg/ml	12 to 19 mg/ml	20 mg/ml or more			
All adult vapers (≥16y)	262	9.2 [6.1–13.7]	18.4 [13.7–24.2]	9.8 [6.3–14.8]	27.1 [21.7–33.4]	16.3 [11.8–22.0]	2.0 [0.9–4.5]	17.2 [12.3–23.4]	
Age (years)									
16-24	46	7.6 [2.7–19.3]	14.5 [6.6–29.0]	10.0 [4.0–23.1]	15.2 [6.8–30.5]	33.1 [19.9–49.5]	4.2 [1.0–16.2]	15.4 [6.9–30.7]	
25-34	51	2.8 [0.7–10.9]	15.3 [6.8–30.9]	11.1 [4.6–24.5]	24.3 [13.8–39.2]	19.6 [10.1–34.6]	1.6 [0.2–11.1]	25.3 [13.6–42.1]	
35-44	51	10.4 [4.2–23.5]	23.7 [13.4–38.4]	18.1 [8.3–35.1]	23.2 [13.0–37.8]	8.8 [3.0–22.7]	0 [0–0]	15.9 [6.9–32.5]	
45-54	37	14.6 [5.9–31.9]	23.1 [11.0–42.3]	7.9 [2.6–21.2]	34.4 [20.1–52.2]	7.4 [2.1–22.7]	1.0 [0.1–7.7]	11.6 [3.5–32.1]	
55-64	39	21.8 [9.7–42.0]	14.7 [5.8–32.6]	2.3 [0.3–15.8]	38.7 [23.2–56.9]	14.2 [5.7–31.1]	0 [0–0]	8.3 [3.1–20.4]	
≥65	38	4.4 [1.0–17.9]	20.6 [9.3–39.7]	1.3 [0.2–9.3]	39.5 [23.0–58.7]	7.5 [2.3–21.9]	6.6 [1.9–20.3]	20.1 [9.2–38.5]	
Gender ³									
Men	121	5.9 [2.9–11.6]	22.5 [15.1–32.2]	8.3 [4.1–16.0]	26.6 [19.0–35.9]	17.1 [10.6–26.5]	2.6 [0.9–7.4]	16.9 [10.0–27.2]	
Women	134	10.8 [6.1–18.3]	14.7 [9.4–22.3]	11.6 [6.6–19.6]	28.3 [20.6–37.5]	16.2 [10.5–24.3]	0.9 [0.2–4.0]	17.5 [11.2–26.3]	
Occupational social grade									
ABC1 (more advantaged)	163	9.7 [5.8–15.9]	19.9 [13.8–27.7]	8.3 [4.9–13.8]	26.3 [19.8–34.1]	18.7 [12.7–26.6]	1.8 [0.5–5.9]	15.4 [10.4–22.1]	
C2DE (less advantaged)	99	8.8 [4.5–16.3]	16.8 [10.2–26.3]	11.3 [5.8–20.8]	28.0 [19.5–38.4]	13.8 [7.9–23.0]	2.3 [0.8–6.5]	19.0 [11.2–30.4]	
Nation									
England	214	9.6 [6.3–14.4]	17.9 [13–24.2]	9.2 [5.6–14.6]	26.3 [20.5–33.0]	17.5 [12.6–23.6]	1.9 [0.8–4.6]	17.7 [12.5–24.4]	
Wales	16	14.3 [2.6–50.6]	34.5 [10.9–69.4]	25.6 [7.7–58.7]	15.0 [3.8–44.1]	3.7 [0.4–26.2]	0 [0–0]	6.9 [1.4–28.5]	
Scotland	32	0 [0–0]	16.5 [6.4–36.6]	11.4 [3.9–28.9]	50.0 [31.3–68.7]	2.9 [0.4–19.0]	5.6 [1.3–21.0]	13.7 [5.3–30.9]	
History of ≥1 diagnosed mental health conditions ⁴									
No	86	13.4 [7.6–22.5]	19.3 [11.7–30.2]	7.3 [2.9–17.1]	20.9 [13.3–31.3]	16.5 [9.6–26.8]	2.8 [0.9–8.7]	19.7 [11.2–32.4]	
Yes	86	4.6 [1.7–12.1]	19.8 [12.3–30.3]	12.1 [6.0–23.2]	35.7 [25.7–47.2]	14.2 [8.0–24.0]	0.9 [0.1–6.5]	12.6 [7.2–21.1]	
Smoking status									
Long-term (≥1y) ex-smoker	80	12.8 [6.8–22.7]	14.1 [7.8–24.2]	9.6 [4.1–20.8]	40.3 [28.9–53.0]	10.9 [5.3–21.0]	0.4 [0.1–3.2]	11.8 [5.4–23.9]	
Recent (<1y) ex-smoker	25	7.5 [1.7–27.5]	34.9 [16.4–59.3]	14.2 [4.8–35.2]	28.6 [13.3–51.2]	5.2 [0.7–31.4]	4.8 [0.6–29.3]	4.7 [0.6–29.1]	
Current smoker	132	7.3 [3.6–14.1]	18.6 [12.2–27.4]	10.2 [5.4–18.3]	20.5 [14.2–28.8]	19.7 [13.1–28.6]	1.5 [0.5–5]	22.2 [14.8–31.9]	
Never smoker	25	10.9 [3.3–30.7]	13.9 [4.9–33.7]	3.9 [0.8–16.7]	22.1 [8.9–45.4]	24.4 [9.9–48.6]	6.8 [1.5–25.7]	18.0 [6.0–42.9]	

¹ Unweighted sample size.² Weighted row percentages.³ Participants who described their gender in another way were excluded from analyses by gender due to low numbers.⁴ Mental health conditions were not collected after June 2023, so results are based on aggregated data across January 2022 – June 2023.