

Supplemental Material

To the manuscript: Postpartum suicidal ideation in Austria and Germany during the COVID-19 pandemic

C. Florea, J. Preiß, M. Angerer, M. Schabus

Table S1: Summary of results

Suicidality score results and relative risk ratios for suicidal ideation			
Factor	Comparing variables	Suicidality score - Statistical test significance	Change in risk (%)
Age	<33 vs. ≥ 33	$W = 481952, p = 0.119$	-23%
Nr children	≥ 2 vs. 1	$W = 464711, p = 0.003$	+63%
Partnership	no vs. yes	$W = 22005, p = 0.796$	+19%
Education	0 vs. 1 vs. 2 vs. 3	$\chi^2 = 4.786, df = 3, p = 0.188$	
	low (0 or 1) vs. high (2 or 3)	$W = 309742, p = 0.151$	+30%
Income	low vs. med. vs. high	$\chi^2 = 6.705, p = 0.082$	
	low vs. med.	$W = 414753, p = 0.001$	+71%
	high vs. med.	$W = 76831, p = 0.951$	-2%
Social support	poor vs. good	$W = 180082, p < 0.001$	+265%
Pandemic Repercussions Score (PRS)	>2 (neg. influence) vs. ≤ 2 (pos. influence)	$W = 494425, p < 0.001$	+163%
PRS - Pregnancy	bad vs. no influence	$W = 223595, p = 0.462$	+18%
	good vs. no influence	$W = 63969, p = 0.583$	-14%
PRS - Birth	bad vs. no influence	$W = 349682, p = 0.07$	+41%
	good vs. no influence	$W = 80315, p = 0.486$	+22%
PRS - Health and dev. Child	bad vs. no influence	$W = 418176, p < 0.001$	+126%
	good vs. no influence	$W = 87153, p = 0.341$	-32%
PRS - Financial security	bad vs. no influence	$W = 426778, p < 0.001$	+94%
	good vs. no influence	$W = 64400, p = 0.698$	+15%
PRS - Partnership	bad vs. no influence	$W = 219888, p < 0.001$	+223%
	good vs. no influence	$W = 220134, p = 0.601$	-13%
Pandemic Distress Score (PDS)	high (> 5) vs. low (≤ 5)	$W = 392738, p = 0.006$	-39%
PDS - Close people sick	slight vs. moderate worry	$W = 235353, p < 0.001$	+112%
	strong vs. moderate worry	$W = 209236, p = 0.072$	+48%
PDS - Self sick	slight vs. moderate worry	$W = 289404, p = 0.015$	+65%
	strong vs. moderate worry	$W = 75046, p = 0.544$	+20%
PDS - COVID-19 baby	slight vs. moderate worry	$W = 159154, p = 0.806$	+39%
	strong vs. moderate worry	$W = 150540, p = 0.439$	+5%

Stress (PSS)	high vs. moderate	$W = 139437, p < 0.001$	+350%
	low vs. moderate	$W = 316365, p < 0.001$	-92%
Access to a midwife	moderately/strongly restricted vs. not/slightly restricted	$W = 313822, p = 0.079$	+38%
Birth type	cesarean vs. natural	$W = 342461, p = 0.407$	+16%
Birth weight	low vs. normal vs. high	$\chi^2 = 0.81, df = 3, p = 0.847$	
	low vs. normal	$W = 65192, p = 0.938$	-4%
	high vs. normal	$W = 150028, p = 0.843$	+5%
Gestational age	preterm vs. term vs. postterm	$\chi^2 = 0.89, df = 3, p = 0.829$	
	preterm vs. term	$W = 123222, p = 0.379$	-28%
	postterm vs. term	$W = 108266, p = 0.909$	+5%
Quarantine	yes vs. no	$W = 309052, p = 0.661$	-9%
Partner support	poor vs. good	$W = 132568, p < 0.001$	+171%
Help from child's grandparents	no vs. yes	$W = 436705, p < 0.001$	+94%
Help from family and friends	no vs. yes	$W = 401164, p < 0.001$	+128%
Effective coping mechanism	no vs. yes	$W = 290587, p < 0.001$	+132%

□ white = no significant change in suicidality score but may still represent a risk or protective factor as shown by the risk change); □ yellow = significant change in suicidality score and an increase in risk between 1-99%; □ orange = significant change in suicidality score and an increase in risk between 100-199%; □ red = significant change in suicidality score and an increase in risk between 200-299%; purple □ = significant change in suicidality score and an increase in risk > 300%; □ Green = protective factor. In the case of trends ($p < 0.1$), the row is left white, and the p value is written down.

Correlations between worry that oneself/family and friends/the child would get COVID-19 and the agreement with the authorities on the restrictions and on COVID-19 severity

Spearman correlations:

	Worry about others getting sick	Worry about oneself getting sick	Worry about baby getting sick
Perception of COVID-19 severity	$p < 0.001, rho = 0.57$	$p < 0.001, rho = 0.53$	$p < 0.001, rho = 0.48$
Agreement with restrictions	$p < 0.001, rho = 0.55$	$p < 0.001, rho = 0.51$	$p < 0.001, rho = 0.49$

A moderation analysis using a linear model to test if one's opinion on the restrictions would moderate the relationship between worry for self/others/baby getting sick and the suicidal ideation was not possible. That is because the relationship between the level of worry and the suicidal ideation was not linear, with the moderate levels of worry having the lowest scores of suicidal ideation, while not worrying at all or worrying very much had higher scores of suicidal ideation.