

Table S1. Exclusion criteria and variable definitions	
Variable	Definition
Hypertension	Systolic blood pressure ≥ 130 mmHg OR diastolic blood pressure ≥ 80 mmHg OR current use of an antihypertensive medication
Diabetes	Hemoglobin A1c (HbA1c) $\geq 6.5\%$ (44.0 mmol/mol) OR current use of insulin OR current use of an anti-hyperglycemic medication
Class III Obesity	Body mass index (BMI) ≥ 40 kg/m ²
Hyperlipidemia	Total cholesterol ≥ 240 mg/dL (6.5 mmol/L) OR current use of a lipid-lowering medication
Chronic kidney disease	Estimated glomerular filtration rate < 60 ml/min per 1.73m ² or urinary albumin-to-creatinine ratio ≥ 30 mg/g (3 mmol/mol)
History of cardiovascular disease	Ever been told by a medical professional that you have had coronary heart disease, angina pectoris, heart failure, a stroke, or a heart attack.
History of cancer	Ever been told by a medical professional that you have had a cancer or malignancy of any kind.
History of respiratory disease	Ever been told by a medical profession that you have chronic bronchitis, emphysema, asthma, or chronic obstructive pulmonary disease.
History of liver disease	Ever been told by a medical professional that you have had any kind of liver condition.
History of arthritis	Ever been told by a medical professional that you have arthritis.
History of dialysis use	Underwent dialysis in the last 12 months.
Hepatitis B	Positive hepatitis B core antibody with duplicate testing.
Hepatitis C	Positive hepatitis C core antibody with duplicate testing.
Human immunodeficiency virus (HIV)	Positive HIV-1 or HIV-2 antibody with duplicate testing and/or confirmatory testing.
Pregnancy	Positive urine pregnancy test.

Table S2. Variable Definitions for Social Determinants of Health			
Variable	Domain[†]	NHANES Question[‡] / Variable Definition	Variable Levels
Poverty to Income Ratio	Economic Stability	Total family income was reported by participants as a range of values (in dollars). NHANES then calculated a family poverty-to-income ratio.	<100%, 100-300%, and ≥300% of the federal poverty level.
Educational Attainment	Education Access and Quality	What is the highest grade or level of school you have completed, or the highest degree you have received?	Below high school, high school or general education diploma, and above high school.
Food Insecurity	Economic Stability	Participants completed an 18-item U.S. Household Food Security Survey Module.	Food secure (zero affirmative responses) and food insecure (≥1 positive response).
Employment	Economic Stability	What kind of work did you do in the last week?	Employed (currently working, student, or retired) and unemployed (not currently working).
Health Insurance	Health Care Access and Quality	Are you covered by health insurance or some other kind of health care plan?	Insured and uninsured.
Routine source of care	Health Care Access and Quality	Is there a place that you usually go when you are sick or need advice about your health? What kind of place do you go most often?	Routine source of care (there is at least one source of care that is not the emergency room) and lack of a routine source of care (no source of care or only rely on the emergency room).
Acculturation	Social and Community Context	What country were you born in? If you were not born in the U.S., how long have you been in the U.S.	U.S. born, non-U.S. born and lived in the U.S. for ≥10 years, and non-U.S. born and lived in the U.S. <10 years.
Cumulative SDOH score	All of the above	One point value was assigned to each adverse social determinant: PIR <100%, below high school education, food insecurity, unemployment, lack of health insurance, lack of a routine source of healthcare, born in the U.S.	A cumulative score was obtained by summing the number of points (range 0-7).
[†] As defined by Healthy People 2030. Available at https://health.gov/healthypeople/priority-areas/social-determinants-health [‡] Details available at https://wwwn.cdc.gov/nchs/nhanes/default.aspx			

Table S3. Missingness prior to imputation	
	N (%)
Smoking status	12 (0.1)
Hemoglobin A1c	19 (0.2)
Body mass index	105 (1.0)
Systolic blood pressure	541 (4.9)
Total cholesterol	876 (8.0)
Serum creatinine	900 (8.2)

Table S4. Participant Characteristics Prior to Imputation (N=10,320).

	Overall	Non-Hispanic White	Non-Hispanic Black	Hispanic†	Other
N	10,320	4,116	1,698	3,433	1,073
Age	35.0 (11.2)	36.2 (11.6)	32.2 (10.0)	32.9 (9.9)	33.6 (9.8)
Female sex	52.0	51.9	52.6	50.1	56.1
Race or ethnicity					
Non-Hispanic White	64.5	100	0	0	0
Mexican American	12.2	0	0	62.6	0
Other Hispanic	7.3	0	0	37.4	0
Non-Hispanic Black	9.1	0	100	0	0
Other Race	6.9	0	0	0	100
Education					
Below High School	15.0	7.7	17.8	39.5	9.3
High School/GED	21.8	21.9	26.0	22.0	14.8
Above High School	63.3	70.4	56.2	38.5	75.9
Poverty-to-Income Ratio					
<100% FPL	14.2	8.5	24.3	29.1	15.1
100-300% FPL	35.7	31.8	44.3	44.3	30.3
≥300 FPL	50.1	59.7	31.3	23.6	54.5
Food insecurity	22.3	14.4	36.7	43.5	18.0
Uninsured	25.0	17.3	32.5	49.1	19.3
Lack of employment	19.2	17.6	21.8	22.7	20.9
Lack a regular source of care	26.1	21.2	28.2	40.5	28.7
Acculturation					
US Born	82.0	94.8	89.0	43.8	39.1
US Born, Lived in US ≥10 years	10.7	2.9	6.6	35.6	31.9
Non-US Born, Lived in US <10 years	7.4	2.4	4.5	20.6	29.0
Cumulative SDOH score	2.5 (1.1)	2.4 (1.0)	2.9 (1.3)	2.7 (1.3)	1.9 (1.2)
Smoking status					
Never	61.2	57.1	68.0	67.8	72.0
Former	16.5	18.6	6.9	14.9	13.0
Current	22.3	24.3	25.1	17.2	15.0
Heavy alcohol use	15.3	14.9	9.9	21.8	7.7
Intravenous drug use	6.8	8.2	3.4	5.1	2.8
Body Mass Index, kg/m ²	26.2 (4.8)	25.9 (4.7)	27.2 (5.3)	27.2 (4.6)	24.8 (4.6)
Hemoglobin A1c, %	5.2 (0.3)	5.2 (0.3)	5.3 (0.4)	5.3 (0.3)	5.3 (0.3)
Total Cholesterol, mg/dL	181.8 (29.5)	182.8 (29.4)	176.3 (30.5)	181.5 (29.5)	180.4 (28.8)

Systolic Blood Pressure, mmHg	111.5 (8.8)	111.6 (9.1)	112.9 (8.6)	111.3 (8.8)	109.9 (9.2)
eGFR, ml/min per 1.73m ²	106.6 (15.9)	104.1 (15.6)	102.3 (16.8)	114.7 (13.8)	111.6 (13.8)

† Hispanic includes both Mexican American and Other Hispanic.

Abbreviations: eGFR, estimated glomerular filtration rate; FPL, federal poverty level; GED, general education diploma.

	Overall	Non-Hispanic White	Non-Hispanic Black	Hispanic†	Other
N	11,413	4,464	1,967	3,752	1,230
Age	34.9 (11.2)	36.1 (11.7)	32.4 (10.1)	32.9 (9.8)	33.8 (10.0)
Female sex	51.9	52.0	51.9	50.0	55.9
Race or ethnicity					
Non-Hispanic White	63.7	100	0	0	0
Mexican American	12.2	0	0	62.6	0
Other Hispanic	7.3	0	0	37.4	0
Non-Hispanic Black	9.6	0	100	0	0
Other Race	7.2	0	0	0	100
Education					
Below High School	15.4	7.9	18.0	40.8	9.9
High School/GED	21.9	22.1	26.4	21.7	14.3
Above High School	62.7	70.0	55.6	37.5	75.9
Poverty-to-Income Ratio					
<100% FPL	14.8	9.0	24.6	29.7	15.9
100-300% FPL	35.7	31.8	43.8	47.2	31.2
≥300 FPL	49.5	59.3	31.6	23.1	52.9
Food insecurity	22.4	14.5	36.0	43.6	18.1
Uninsured	25.3	17.5	32.2	49.9	19.1
Lack of employment	19.6	18.0	22.8	22.6	21.5
Lack a regular source of care	26.4	21.5	28.2	41.0	28.4
Acculturation					
US Born	81.4	94.7	88.2	43.3	37.6
US Born, Lived in US ≥10 years	10.9	3.0	6.8	35.9	32.7
Non-US Born, Lived in US <10 years	7.7	2.4	5.0	20.8	29.7
Cumulative SDOH score	2.5 (1.1)	2.4 (1.0)	2.9 (1.3)	2.8 (1.3)	1.9 (1.2)
Smoking status					
Never	61.6	57.4	68.5	68.0	72.2
Former	16.0	18.1	6.6	14.7	12.8
Current	22.4	24.5	24.9	17.3	15.0
Heavy alcohol use	20.9	20.0	17.3	27.7	16.4
Intravenous drug use	6.5	8.0	3.1	5.0	2.5
Body Mass Index, kg/m ²	26.2 (4.8)	25.9 (4.8)	27.2 (5.3)	27.2 (4.6)	24.8 (4.6)
Hemoglobin A1c, %	5.2 (0.3)	5.2 (0.3)	5.3 (0.4)	5.3 (0.3)	5.3 (0.3)
Total Cholesterol, mg/dL	181.8 (29.6)	182.8 (29.4)	177.0 (30.4)	181.6 (29.7)	180.7 (29.0)

Systolic Blood Pressure, mmHg	111.6 (9.1)	111.7 (9.1)	113.0 (8.8)	111.3 (8.8)	110.3 (9.2)
eGFR, ml/min per 1.73m ²	106.4 (16.2)	104.1 (15.8)	102.5 (17.0)	114.4 (14.2)	110.8 (14.8)

† Hispanic includes both Mexican American and Other Hispanic.

Abbreviations: eGFR, estimated glomerular filtration rate; FPL, federal poverty level; GED, general education diploma; SDOH, social determinants of health.

Table S6. Trends in Age-Adjusted* Prevalence of Social Determinants of Health, 1999-2018.						
	Overall	1999-2002	2003-2008	2009-2014	2015-2018	p-trend
Poverty-to-Income Ratio						
<100% FPL	11.8 (10.8, 12.8)	10.6 (8.6, 12.6)	11.3 (9.4, 13.2)	14.1 (12.3, 15.9)	10.3 (8.2, 12.3)	0.10
100-300% FPL	33.2 (31.1, 35.3)	34.4 (29.4, 39.4)	38.1 (34.9, 41.2)	30.6 (26.4, 34.8)	32.5 (28.4, 36.6)	0.91
≥300% FPL	55.1 (52.7, 57.4)	55.0 (49.4, 60.5)	50.6 (47.1, 54.1)	55.3 (50.6, 60.0)	57.2 (52.5, 61.9)	0.46
Educational Attainment						
Below high school	15.5 (14.1, 17.0)	19.8 (16.1, 23.5)	16.4 (13.8, 19.0)	15.4 (12.8, 18.0)	12.1 (9.1, 15.2)	<0.001
High school/GED	19.9 (18.1, 21.6)	20.3 (15.5, 25.1)	22.4 (19.7, 25.1)	17.4 (14.8, 20.0)	20.3 (15.5, 25.1)	0.50
Above high school	64.5 (62.2, 66.8)	60.0 (55.2, 64.7)	61.1 (57.9, 64.3)	67.2 (63.7, 70.7)	67.5 (61.1, 74.0)	0.02
Food Insecure	17.5 (16.2, 18.9)	11.5 (9.6, 13.4)	15.5 (13.4, 17.6)	19.1 (17.2, 21.2)	23.1 (18.9, 27.3)	<0.001
Unemployed	18.9 (17.3, 20.5)	18.7 (15.9, 21.5)	17.0 (14.8, 19.3)	19.4 (17.0, 21.8)	19.7 (15.5, 23.9)	0.53
Uninsured	20.5 (19.0, 21.9)	19.3 (16.0, 22.6)	23.3 (20.3, 26.3)	22.9 (20.2, 25.7)	15.1 (12.3, 18.0)	0.02
No usual source of care	21.2 (19.9, 22.5)	21.1 (18.4, 23.8)	22.5 (19.6, 25.3)	19.0 (17.0, 21.1)	23.1 (20.4, 25.8)	0.67
* Age-standardized to the 2010 U.S. Census						
Abbreviations: FPL, federal poverty level; GED, general education diploma.						

Table S7. Adjusted cumulative incidence of mortality and absolute risk difference by social determinant of health				
	10-year adjusted cumulative incidence (95% CI)	15-year adjusted cumulative incidence (95% CI)	10-year risk difference (95% CI)	15-year risk difference (95% CI)
Poverty-to-Income Ratio (PIR)				
<100% FPL	3.2% (1.5-4.9)	5.6% (2.8-8.5)		
100-300% FPL	2.0% (1.2-2.7)	3.5% (2.1-4.9)	0.6% (-0.03-1.6) ^a	1.1% (-0.7-3.0) ^a
≥300% FPL	1.3% (0.7-2.0)	2.4% (1.1-3.7)	1.9% (0.06-3.7) ^b	3.2% (0.08-6.3) ^b
Educational Attainment				
Below high school	3.0% (1.9-4.2)	5.2% (3.2-7.3)		
High school/GED	2.1% (0.9-3.3)	3.7% (1.6-5.8)	0.9% (-0.7-2.5) ^c	1.5% (-1.3-4.4) ^c
Above high school	1.4% (1.0-1.9)	2.5% (1.6-3.4)	1.6% (0.4-2.8) ^d	2.7% (0.4-4.9) ^d
Employment Status				
Unemployed	2.4% (1.5-3.3)	4.1% (2.7-5.5)	0.4% (-0.4-1.4)	0.8% (-0.8-2.5)
Employed	1.9% (1.4-2.3)	3.3% (2.3-4.2)		
Food Security				
Food insecure	2.9% (1.7-4.1)	4.9% (2.7-7.2)	1.2% (-0.06-2.4)	2.0% (-0.04-4.4)
Food secure	1.7% (1.3-2.1)	2.9% (2.2-3.7)		
Health Insurance				
Uninsured	2.4% (1.4-3.5)	4.2% (2.2-6.2)	0.6% (-0.5-1.7)	1.0% (-1.0-3.2)
Insured	1.8% (1.4-2.2)	3.1% (2.3-4.0)		
Access to Care				
Lack of a routine source of healthcare	2.4% (1.2-3.5)	4.1% (2.1-6.0)	0.5% (-0.7-1.7)	0.8% (-1.3-3.0)
Routine source of healthcare	1.9% (1.4-2.3)	3.2% (2.3-4.2)		
Nativity and Acculturation				
U.S. Born	2.3% (1.7-3.0)	4.0% (2.8-5.2)		
Non-U.S. Born, Lived in U.S. ≥ 10 years	1.8% (0.8-2.8)	3.1% (1.5-4.7)	0.5% (-0.6-1.7) ^e	0.9% (-1.1-2.9) ^e
Non-U.S. Born, Lived in U.S. < 10 years	0.7% (0.0-1.4)	1.2% (0.0-2.5)	1.6% (0.6-2.6) ^f	2.8% (0.9-4.6) ^f
^a Risk difference represents: PIR <100% – PIR 100-300%				
^b Risk difference represents: PIR <100% – PIR >300%				
^c Risk difference represents: below high school – above high school				
^d Risk difference represents: below high school – high school				
^e Risk difference represents: U.S. Born – Non-U.S. Born, Lived in the U.S. ≥ 10 years				
^f Risk difference represents: U.S. Born – Non-U.S. Born, Lived in the U.S. < 10 years				

Models adjusted for age, sex, race or ethnicity, smoking status, body mass index, hemoglobin A1c, total cholesterol, systolic blood pressure, eGFR, survey year, alcohol use, and injection drug use.

Abbreviations: CI, confidence interval; FPL, federal poverty level; GED, general education diploma; PIR, poverty to income ratio; U.S., United States.

Table S8. Associations between social determinants of health and mortality.					
	N †	Crude ^a	Model 1 ^b	Model 2 ^c	Model 3 ^d
		HR [95% CI]	HR [95% CI]	HR [95% CI]	HR [95% CI]
Poverty-to-Income Ratio	10,433				
<100% FPL		1.64 [1.02, 2.62]	3.12 [1.86, 5.24]	2.40 [1.38, 4.18]	2.48 [1.43, 4.32]
100-300% FPL		1.19 [0.82, 1.71]	1.72 [1.19, 2.49]	1.49 [1.04, 2.14]	1.52 [1.06, 2.19]
≥300% FPL		Referent	Referent	Referent	Referent
Education	11,402				
Below high school		2.23 [1.60, 3.11]	2.72 [1.84, 4.01]	2.17 [1.49, 3.16]	2.22 [1.52, 3.21]
High school/GED		1.54 [1.04, 2.30]	1.73 [1.18, 2.54]	1.52 [1.05, 2.21]	1.52 [1.03, 2.23]
Above high school		Referent	Referent	Referent	Referent
Lack of employment	11,407	1.07 [0.75, 1.52]	1.34 [0.94, 1.93]	1.28 [0.89, 1.83]	1.28 [0.90, 1.84]
Food insecurity	11,138	1.32 [0.95, 1.83]	2.10 [1.47, 3.02]	1.73 [1.20, 2.49]	1.73 [1.20, 2.51]
Nativity and Acculturation	10,394				
U.S. Born		5.10 [2.24, 11.60]	3.75 [1.56, 9.06]	3.53 [1.46, 8.52]	3.44 [1.42, 8.34]
U.S. Born, Lived in U.S. ≥10 years		4.66 [1.82, 11.96]	2.70 [1.04, 6.99]	2.68 [1.04, 6.89]	2.67 [1.03, 6.88]
Non-US Born, Lived in U.S. <10 years		Referent	Referent	Referent	Referent
Lack of health insurance	11,359	1.07 [0.77, 1.51]	1.67 [1.17, 2.37]	1.37 [0.94, 1.99]	1.36 [0.93, 1.98]
Lack of regular source of care	11,413	1.08 [0.74, 1.58]	1.47 [0.99, 2.18]	1.31 [0.88, 1.94]	1.29 [0.88, 1.91]
Cumulative SDOH score	11,413	1.25 [1.12, 1.41]	1.45 [1.27, 1.64]	1.32 [1.15, 1.52]	1.33 [1.16, 1.52]

† Sample size varies due to variable participant non-response to social determinants of health questions.

^a Crude

^b Model 1: adjusted for age, sex, race or ethnicity, and survey year.

^c Model 2: adjusted for Model 1 + smoking status, body mass index, hemoglobin A1c, total cholesterol, systolic blood pressure, eGFR.

^d Model 3: adjusted for Model 2 + alcohol use, injection drug use.

Abbreviations: eGFR, estimated glomerular filtration rate; FPL, federal poverty level; GED, general education diploma; HR, hazard ratio; SDOH, social determinants of health; U.S., United States.