

Title: Non-genetic mother-infant factors and Human milk oligossacharides composition/profile in breast milk of Ugandan mothers

First Author: Tonny Jimmy Owalla

Supplementary Table 1: Median and Percent HMO concentrations

HMO	All		Secretors		Non-secretors	
	nmol/mL	%	nmol/mL	%	nmol/mL	%
2'FL	5929.5	36.9	7099.7	42.5	31.7	0.3
LNFP II	1411.1	8.8	1140.5	6.8	2884.1	32.1
LNT	1028.7	6.4	981.3	5.8	1330.7	14.8
DFLNT	865.9	5.4	1215.9	7.3	619.0	6.9
LNnT	805.2	5.0	789.1	4.7	870.1	9.7
3'SL	632.0	3.9	619.5	3.7	692.7	7.7
LNFP I	599.3	3.7	1019.8	6.1	91.5	1.0
DFlac	548.4	3.4	666.0	4.0	56.1	0.6
DSLNT	417.2	2.6	404.4	2.4	429.3	4.8
3FL	378.9	2.4	397.7	2.4	189.3	2.1
6'SL	157.8	1.0	150.9	0.9	185.9	2.1
FDSLNH	146.7	0.9	134.0	0.8	216.3	2.4
LSTb	123.5	0.8	119.2	0.7	183.8	2.0
LNH	71.2	0.4	68.6	0.4	80	0.9
LNFP III	70.0	0.4	63.5	0.4	93.8	1.0
LSTc	46.5	0.3	46.9	0.3	41.3	0.5
DSLNH	31.7	0.2	31.7	0.2	31.3	0.3
DFLNH	24.4	0.2	28.3	0.2	9.5	0.1
FLNH	24.2	0.2	25.8	0.2	21	0.2
Total HMO	16050.6		16699.2		8988.5	100
Fuc HMO	15962.3	99.5	14641.2	87.7	5611.3	62.4
Sial HMO	2395.0	14.9	2325.6	13.9	2984.1	33.2

HMOs = human milk oligosaccharides; nmol/mL = nanomoles/milliliter; % = percent

First Author: Tonny Jimmy Owalla

Supplementary Table 2: Median (IQR) HMO concentrations and sex of children

HMO nmol/mL	Total (N=114)	Females (n=60)	Males (n=54)	Unadjusted p-value
2'FL	5961.3 (2365.5- 9364.7)	5961.3 (3086.9- 9616.9)	6069.2 (1660.3- 9271.8)	0.97
3FL	355.5 (263.8- 593.5)	371.2 (240.5- 536.2)	353.8 (289.4- 660.3)	0.35
LNnT,	782.3 (578.2- 1042.6)	768.5 (582.9- 1037.4)	813.1 (567.9- 1093.1)	0.65
3'SL	614.1 (352.5- 964.9)	516.6 (309.8- 853.8)	701.8 (454.5- 1059.8)	0.035
DFLac	493.2 (208.5- 832.6)	457.6 (281.9- 744.6)	557.0 (194.8- 961.2)	0.57
6'SL	154.0 (100.3- 276.4)	148.9 (101.3- 230.4)	183.0 (97.8- 297.8)	0.39
LNT	1010.4 (702.1- 1618.0)	982.2 (730.7- 1513.2)	1076.7 (659.6- 1704.2)	0.88
LNFP I	623.9 (169.2- 1863.9)	729.9 (224.4- 2039.1)	455.8 (169.2- 1781.5)	0.64
LNFP II	1393.9 (721.0- 2582.7)	1359.6 (759.1- 2613.2)	1476.8 (599.7- 2529.6)	0.73
LNFP III	71.0 (55.0-88.8)	65.0 (54.2-84.6)	75.0 (58.9-93.2)	0.14
LSTb	123.3 (94.8- 182.2)	123.3 (101.3- 178.7)	124.7 (83.3- 182.2)	0.62
LSTc,	43.1 (22.2-92.4)	41.8 (23.4-72.3)	45.8 (21.4-104.3)	0.71
DFLNT	837.2 (106.0- 1609.2)	837.2 (129.8- 1666.7)	868.0 (80.4- 1544.0)	0.97
LNH	67.8 (37.2-107.4)	74.8 (41.6-125.0)	60.7 (30.7-82.9)	0.16

Title: Non-genetic mother-infant factors and Human milk oligossacharides composition/profile in breast milk of Ugandan mothers

First Author: Tonny Jimmy Owalla

DSLNT	425.8 (280.5-515.5)	377.7 (280.6-476.5)	443.6 (274.3-585.6)	0.12
FLNH	24.0 (13.2-52.0)	25.3 (13.1-50.9)	23.7 (13.2-58.5)	0.86
DFLNH	25.6 (10.1-66.1)	26.6 (10.5-66.1)	25.1 (9.3-66.1)	0.86
FDSLNT	144.7 (21.1-290.3)	171.5 (16.9-313.6)	102.2 (23.1-237.9)	0.54
DSLNT	26.1 (11.1-63.7)	24.3 (7.8-81.0)	30.9 (12.0-62.5)	0.95
Total HMO	16021.3 (12674.9-17668.9)	15804.3 (12947.1-17249.5)	16174.0 (12197.8-18096.5)	0.43
Sia HMO	13985.2 (10010.4-15618.4)	13756.7 (9913.9-15226.6)	14186.8 (10517.4-16252.6)	0.49
Fuc HMO	2388.4 (1965.7-3032.4)	2322.1 (1836.1-2861.0)	2403.0 (2026.4-3305.9)	0.17
Diversity	4.7 (3.0-5.9)	4.7 (3.0-6.0)	4.7 (3.0-5.9)	0.96

IQR = interquartile range; HMOs = human milk oligosaccharides; N = total number of mothers with available information about the sex of their children; data on sex was not available for 13 children; n = number of females or males. HMO with significant unadjusted p value is in bold.

Title: Non-genetic mother-infant factors and Human milk oligossacharides composition/profile in breast milk of Ugandan mothers

First Author: Tonny Jimmy Owalla

Supplementary Table 3: Correlations between HMO concentrations and lactation duration

HMOs nmol/mL	All, N= 127		Secretor, n=101		Non Secretor, n= 26	
	Coef. (rho)	Unadjusted p-value	Coef. (rho)	Unadjusted p-value	Coef. (rho)	Unadjusted p-value
2'FL	0.0479	0.5925	0.0729	0.4686	0.2652	0.1904
3FL	0.3551	<0.0001	0.3626	0.0002	0.3802	0.0554
LNnT,	-0.3680	<0.0001	-0.4188	<0.0001	-0.2440	0.2297
3'SL	0.4104	<0.0001	0.3378	0.0006	0.7062	0.0001
DFLac	0.2740	0.0018	0.4214	<0.0001	0.5195	0.0065
6'SL	-0.6915	<0.0001	-0.6586	<0.0001	-0.8010	<0.0001
LNT	-0.1993	0.0247	-0.1919	0.0545	-0.2785	0.1683
LNFP I	-0.2133	0.0160	-0.3105	0.0016	0.0544	0.7918
LNFP II	0.1097	0.2194	0.2002	0.0448	-0.0400	0.8460
LNFP III	-0.0596	0.5056	-0.0545	0.5881	-0.0127	0.9510
LSTb	0.0087	0.9226	-0.0250	0.8043	0.1417	0.4900
LSTc	-0.6630	<0.0001	-0.6429	<0.0001	-0.7343	<0.0001
DFLNT	-0.1125	0.2080	-0.1406	0.1607	0.1649	0.4207
LNH	-0.2756	0.0017	-0.2100	0.0350	-0.4660	0.0164
DSLNT	-0.0346	0.6994	-0.1501	0.1340	0.3904	0.0486
FLNH	-0.3090	0.0004	-0.3334	0.0007	-0.1393	0.4975
DFLNH	-0.2411	0.0063	-0.2860	0.0037	-0.0339	0.8694
FDSLNH	-0.0427	0.6333	0.0994	0.3226	-0.4722	0.0149
DSLNH	-0.5310	<0.0001	-0.5026	<0.0001	-0.6223	0.0007
Total HMO	-0.0183	0.8383	0.0070	0.9450	-0.1718	0.4014
Fuc HMO	0.0937	0.2948	0.1691	0.0909	0.0366	0.8591
Sia HMO	-0.0631	0.4806	-0.0709	0.4811	-0.0106	0.9590
Diversity	-0.0989	0.2687	-0.1030	0.3056	-0.1271	0.5362

Coef (rho) = Spearman's correlation coefficient. HMOs = human milk oligosaccharides; nmol/mL = nanomoles/milliliter; N = total number of mothers; n = number of secretors or non-secretors. HMOs with significant unadjusted p value are in bold.

Title: Non-genetic mother-infant factors and Human milk oligossacharides composition/profile in breast milk of Ugandan mothers

First Author: Tonny Jimmy Owalla

Supplementary Table 4: Correlations between HMO concentrations and maternal age

HMOs nmol/mL	All, N= 126		Secretor, N= 100		Non-Secretor, N= 26	
	Coef. (rho)	Unadjusted p-value	Coef. (rho)	Unadjusted p-value	Coef. (rho)	Unadjusted p-value
2'FL	0.0410	0.6488	0.1228	0.2235	0.4584	0.0185
3FL	0.0737	0.4124	0.1500	0.1363	-0.0646	0.7539
LNnT	-0.2213	0.0128	-0.3040	0.0021	0.0357	0.8624
3'SL	0.1753	0.0505	0.1302	0.1989	0.3591	0.0716
DFLac	0.0537	0.5505	0.1403	0.1639	0.4506	0.0209
6'SL	-0.0344	0.7024	-0.0409	0.6862	-0.0351	0.8650
LNT	-0.1152	0.1989	-0.1559	0.1214	-0.0213	0.9177
LNFP I	-0.0767	0.3933	-0.0103	0.9192	-0.3560	0.0742
LNFP II	-0.0080	0.9288	0.0090	0.9292	-0.1625	0.4276
LNFP III	0.0556	0.5362	0.0313	0.7569	-0.0653	0.7513
LSTb	-0.0153	0.8651	-0.0632	0.5324	0.2285	0.2615
LSTc	-0.0684	0.4468	-0.0925	0.3602	0.0502	0.8077
DFLNT	-0.1103	0.2188	-0.0844	0.4038	-0.0443	0.8297
LNH	-0.1639	0.0667	-0.1699	0.0911	-0.1966	0.3358
DSLNT	-0.0673	0.4542	-0.2181	0.0293	0.4612	0.0177
FLNH	-0.0372	0.6790	-0.0628	0.5348	0.1887	0.3560
DFLNH	0.0895	0.3190	0.1379	0.1713	0.0160	0.9382
FDSLNH	0.0531	0.5551	0.0798	0.4300	-0.1378	0.5020
DSLNH	-0.0254	0.7779	-0.0335	0.7411	0.0184	0.9290
Total HMO	0.0343	0.7030	0.1344	0.1826	0.0818	0.6912
Fuc HMO	0.0957	0.2863	0.2423	0.0151	-0.0704	0.7324
Sia HMO	-0.0093	0.9177	-0.1060	0.2941	0.2598	0.1999
Diversity	-0.1091	0.2240	-0.1425	0.1572	-0.0637	0.7571

Coef (rho) = Spearman's correlation coefficient. HMOs = human milk oligosaccharides; nmol/mL = nanomoles/milliliter; N = total number of mothers; n = number of secretors or non-secretors. HMOs with significant unadjusted p value are in bold.

Title: Non-genetic mother-infant factors and Human milk oligossacharides composition/profile in breast milk of Ugandan mothers

First Author: Tonny Jimmy Owalla

Supplementary Table 5: Median (IQR) HMO concentrations and maternal age

HMOs	Total (N=127)	>18 years (n=123)	≤ 18years (n=4)	Unadjusted p-value
2'FL	5912.6 (2715.8-9271.8)	5822.8 (2452.3-9279.6)	7181.2 (5115.1-8249.1)	0.73
3FL	375.4 (263.8-593.5)	384.4 (263.8-606.8)	324.9 (257.7-380.7)	0.43
LNnT	797.2 (578.2-1033.2)	775.1 (567.9-1033.2)	966.2 (872.0-1497.4)	0.12
3'SL	630.0 (353.4-985.9)	640.2 (364.9-987.8)	434.3 (303.9-545.7)	0.18
DFLac	546.5 (287.0-838.1)	546.5 (231.6-881.5)	582.7 (385.4-722.3)	0.92
6'SL	158.5 (102.5-270.9)	156.6 (102.5-263.3)	480.3 (272.6-909.2)	0.10
LNT	1039.5 (702.1-1618.0)	1039.5 (694.8-1531.9)	1267.2 (834.4-2033.7)	0.48
LNFP I	609.7 (172.6-1863.9)	579.4 (169.2-1846.9)	2279.7 (1137.6-3344.6)	0.050
LNFP II	1393.9 (721.0-2551.2)	1419.6 (735.3-2551.2)	360.8 (147.4-1569.8)	0.23
LNFP III	70.5 (55.4-88.8)	70.5 (57.1-89.8)	60.3 (45.2-74.1)	0.28
LSTb	123.3 (90.8-182.2)	123.2 (89.1-181.9)	202.0 (113.5-291.4)	0.19
LSTc	46.6 (24.8-93.3)	45.7 (24.8-87.9)	201.3 (70.3-690.2)	0.11
DFLNT	852.8 (122.3-1544.0)	852.8 (122.3-1526.3)	1085.4 (190.7-1916.3)	0.68
LNH	70.7 (37.7-105.2)	71.5 (38.2-106.4)	28.5 (24.6-61.9)	0.16

Title: Non-genetic mother-infant factors and Human milk oligossacharides composition/profile in breast milk of Ugandan mothers

First Author: Tonny Jimmy Owalla

	DSLNT	412.1 (274.3-512.6)	400.9 (271.0-504.0)	641.2 (600.7-922.3)	0.009
FLNH	24.2	(13.2-52.0)	24.0 (13.2-52.0)	28.3 (17.7-70.8)	0.80
DFLNH	24.5	(9.4-66.1)	23.5 (9.3-66.1)	34.3 (26.9-100.3)	0.24
FDSLNH	146.9	(21.1-310.1)	157.5 (21.2-311.1)	31.3 (14.9-106.7)	0.16
DSLNH	32.1	(11.6-64.5)	32.1 (11.6-64.5)	35.5 (15.3-219.8)	0.68
Total HMO	16021.3	(13219.4-17668.9)	15907.3 (13190.4-17550.4)	17502.3 (15383.9-18768.4)	0.22
Sia HMO	13925.6	(10903.9-15947.2)	13925.6 (10517.4-15947.2)	13607.5 (11691.6-15523.7)	0.97
Fuc HMO	2388.4	(1995.9-3149.7)	2381.0 (1977.8-3114.2)	3202.3 (2500.3-4456.0)	0.10
Diversity	4.7	(3.1-6.0)	4.7 (3.0-6.0)	4.8 (3.6-6.5)	0.52

IQR = interquartile range; HMOs = human milk oligosaccharides; N = total number of mothers; n = number of mothers in each age category. HMOs with significant unadjusted p values are in bold.

Title: Non-genetic mother-infant factors and Human milk oligossacharides composition/profile in breast milk of Ugandan mothers

First Author: Tonny Jimmy Owalla

Supplementary Table 6: Median (IQR) HMO concentrations and maternal parity

HMOs	Total (N=127)	Parity=1 (n=27)	Parity >1 (n=100)	Unadjusted p-value
nmol/mL				
2'FL	5929.5 (2715.8-9271.8)	6242.8 (624.9-8908.1)	5912.6 (3799.2-9275.7)	0.56
3FL	378.9 (263.8-606.8)	327.1 (221.0-406.1)	401.0 (265.2-632.7)	0.076
LNnT	805.2 (578.2-1033.5)	894.2 (594.1-1141.3)	790.0 (573.0-1032.7)	0.28
3'SL	632.0 (353.4-987.8)	413.1 (295.2-985.9)	700.6 (403.4-1016.9)	0.072
DFLac	548.4 (287.0-881.5)	399.3 (158.4-698.6)	594.8 (365.0-937.9)	0.018
6'SL	157.8 (102.2-270.9)	182.2 (86.3-569.7)	155.1 (102.8-253.6)	0.38
LNT	1028.7 (694.8-1618.0)	1106.3 (752.7-1747.0)	1026.5 (666.2-1473.4)	0.65
LNFP I	599.3 (172.6-1863.9)	1080.8 (117.9-2066.2)	529.4 (212.1-1730.5)	0.44
LNFP II	1411.1 (721.0-2551.2)	1580.5 (107.9-2766.3)	1393.9 (787.8-2495.3)	0.77
LNFP III	70.0 (55.0-88.8)	68.4 (44.9-82.5)	73.9 (58.8-93.2)	0.11
LSTb	123.5 (90.8-182.2)	134.5 (87.6-184.4)	122.3 (92.8-181.8)	0.51
LSTc	46.5 (24.3-93.3)	63.7 (18.5-140.8)	43.1 (25.2-75.1)	0.19
DFLNT	865.9 (122.3-1576.6)	802.1 (121.3-1705.9)	993.8 (126.1-1560.3)	0.91
LNH	71.2 (37.7-106.4)	68.6 (23.0-104.0)	71.8 (41.6-107.8)	0.40
DSLNT	417.2 (274.3-512.6)	454.3 (287.3-633.3)	412.1 (268.7-501.5)	0.30

Title: Non-genetic mother-infant factors and Human milk oligossacharides composition/profile in breast milk of Ugandan mothers

First Author: Tonny Jimmy Owalla

FLNH	24.2 (13.2-57.0)	20.5 (8.3-35.6)	24.5 (13.6-59.5)	0.23
DFLNH	24.4 (9.4-66.1)	16.9 (9.0-32.8)	26.1 (10.1-74.3)	0.21
FDSLNH	146.7 (21.1-310.1)	123.9 (15.8-337.3)	157.7 (25.5-302.0)	0.39
DSLNH	31.7 (11.6-64.5)	30.7 (6.5-65.8)	32.1 (12.7-64.1)	0.98
Total HMO	16050.6 (13219.4-17668.9)	15755.6 (10981.1-17187.8)	16125.7 (13379.3-17682.1)	0.73
Sia HMO	13962.3 (10903.9-15966.1)	13381.6 (7401.8-15258.1)	14117.4 (11166.4-16134.6)	0.22
Fuc HMO	2395.0 (1995.9-3149.7)	2398.6 (1795.0-3301.5)	2388.4 (1995.9-3022.9)	0.76
Diversity	4.7 (3.1-6.0)	4.3 (3.1-6.0)	4.7 (3.1-6.0)	0.91

IQR = interquartile range; HMOs = human milk oligosaccharides; N = total number of mothers; n = number of mothers in each parity category. HMO with a significant unadjusted p value is in bold.

Title: Non-genetic mother-infant factors and Human milk oligossacharides composition/profile in breast milk of Ugandan mothers

First Author: Tonny Jimmy Owalla

Supplementary Table 7: Correlations between HMOs and maternal post-partum BMI

HMO nmol/mL	All, N= 113		Secretor, n= 89		Non-Secretor, n= 24	
	Coef. (rho)	Unadjusted p-value	Coef. (rho)	Unadjusted p-value	Coef. (rho)	Unadjusted p-value
2'FL	-0.0547	0.5651	-0.0662	0.5376	-0.1283	0.5502
3FL	0.0184	0.8469	0.0502	0.6401	-0.1313	0.5407
LNnT	-0.0266	0.7794	-0.0219	0.8386	-0.0565	0.7930
3'SL	0.1361	0.1525	0.1700	0.1134	-0.0291	0.8925
DFLac	0.0255	0.7888	0.0580	0.5890	-0.0726	0.7359
6'SL	0.0088	0.9259	-0.0310	0.7733	0.2392	0.2603
LNT	0.0521	0.5837	0.0123	0.9092	0.1913	0.3704
LNFP I	-0.0848	0.3717	-0.1091	0.3090	-0.0939	0.6624
LNFP II	0.0703	0.4594	0.1374	0.1992	-0.1366	0.5246
LNFP III	0.0667	0.4824	0.0805	0.4534	-0.0992	0.6448
LSTb	-0.0196	0.8364	0.0308	0.7745	-0.1992	0.3508
LSTc	-0.0279	0.7695	-0.0249	0.8171	-0.0918	0.6698
DFLNT	-0.0965	0.3093	-0.1180	0.2709	-0.2231	0.2947
LNH	-0.1641	0.0824	-0.0917	0.3926	-0.4188	0.0417
DSLNT	0.0017	0.9855	-0.0297	0.7824	0.0952	0.6580
FLNH	-0.0383	0.6868	-0.0084	0.9380	-0.2092	0.3266
DFLNT	-0.0217	0.8194	-0.0066	0.9512	-0.1653	0.4401
FDSLNT	0.1107	0.2431	0.2126	0.0454	-0.2405	0.2577
DSLNT	-0.0712	0.4538	-0.0718	0.5035	-0.1131	0.5988
Total HMO	-0.0234	0.8059	-0.0108	0.9200	-0.2592	0.2213
Fuc HMO	-0.0649	0.4944	-0.0946	0.3777	-0.0822	0.7026
Sia HMO	0.0718	0.4500	0.1272	0.2348	-0.2283	0.2833
Diversity	0.0311	0.7439	0.0585	0.5860	-0.1864	0.3833

Coef (rho) = Spearman's correlation coefficient. HMO = human milk oligosaccharides; nmol/mL = nanomoles/milliliter; N = total number of mothers with BMI data available; height or weight data were unavailable for 14 mothers; n = number of secretors or non-secretors. HMOs with significant unadjusted p values are in bold.

Title: Non-genetic mother-infant factors and Human milk oligossacharides composition/profile in breast milk of Ugandan mothers

First Author: Tonny Jimmy Owalla

Supplementary Table 8: Median (IQR) HMO concentrations and Maternal post-partum BMI in Secretor Mothers

HMO nmol/mL	Total (N=101)	Normal (n=82)	Overweight (n=19)	Unadjusted p-value
2'FL	7059.5 (4978.3- 9808.0)	7215.0 (4991.9- 9774.8)	6242.8 (4468.4- 10037.1)	0.43
3FL	399.2 (294.5- 632.7)	400.8 (297.8- 633.6)	370.5 (212.5-543.2)	0.32
LNnT	797.2 (558.8- 1033.4)	805.2 (594.1- 1033.5)	774.7 (439.5- 1026.8)	0.46
3'SL	624.8 (346.2- 916.9)	630.0 (364.9- 916.3)	528.1 (309.0-958.8)	0.71
DFLac	658.6 (451.6- 1041.7)	698.6 (457.1- 1128.3)	480.8 (404.8-888.2)	0.28
6'SL	150.7 (99.0- 270.9)	150.5 (97.8- 263.3)	150.9 (107.7-320.8)	0.70
LNT	989.0 (649.9- 1473.4)	964.4 (653.7- 1395.3)	1106.3 (485.9- 1625.3)	0.78
LNFP I	974.5 (387.9- 2112.2)	845.0 (382.9- 2090.4)	1230.9 (537.4- 2133.9)	0.80
LNFP II	1154.3 (558.1- 2089.4)	1106.0 (535.5- 2085.7)	1580.5 (897.6- 2273.7)	0.22
LNFP III	63.3 (49.3-81.6)	63.5 (52.4-81.2)	61.7 (45.0-88.0)	0.73
LSTb	119.4 (85.6- 159.0)	119.2 (87.4- 142.8)	126.4 (65.7-216.6)	0.61
LSTc	46.8 (23.9-95.2)	46.7 (23.3-77.2)	58.3 (31.5-137.3)	0.31
DFLNT	1223.6 (128.0- 1739.9)	1313.1 (190.2- 1815.0)	144.8 (78.7-1421.5)	0.14
LNH	69.4 (37.6-97.9)	73.2 (37.2-103.1)	51.3 (41.2-83.7)	0.38

Title: Non-genetic mother-infant factors and Human milk oligossacharides composition/profile in breast milk of Ugandan mothers

First Author: Tonny Jimmy Owalla

DSLNT	405.7 (261.4-505.7)	426.3 (263.2-512.6)	320.0 (260.2-456.4)	0.21
FLNH	26.0 (11.8-58.5)	26.2 (11.9-58.5)	15.8 (9.1-42.0)	0.44
DFLNH	28.1 (12.1-74.3)	28.0 (11.5-67.4)	28.3 (16.7-76.6)	0.86
FDSLNH	140.3 (20.8-270.6)	102.2 (18.3-237.9)	213.6 (107.1-337.3)	0.028
DSLNH	31.4 (12.7-61.8)	29.1 (11.6-62.1)	33.7 (15.8-60.0)	0.62
Total HMO	16650.0 (15344.7-17878.3)	16725.9 (15434.2-18096.5)	16237.7 (14883.9-17695.3)	0.35
Sia HMO	2329.0 (1930.1-2836.4)	2335.1 (1934.2-2796.0)	2269.9 (1760.5-3169.8)	0.92
Fuc HMO	14638.9 (13461.9-16231.6)	14641.2 (13750.5-16277.2)	13528.5 (11826.2-16182.3)	0.086
Diversity	4.4 (2.9-6.1)	4.1 (2.9-6.2)	5.0 (2.8-6.0)	0.77

IQR = interquartile range; HMOs = human milk oligosaccharides; N = total number of mothers; n = number of mothers in each weight category. HMO with a significant unadjusted p value is in bold.

Title: Non-genetic mother-infant factors and Human milk oligossacharides composition/profile in breast milk of Ugandan mothers

First Author: Tonny Jimmy Owalla

Supplementary Table 9: Median (IQR) HMO concentrations and Maternal BMI in Non-secretor Mothers

HMO nmol/mL	Total (N=26)	Normal (n=20)	Overweight (n=6)	Unadjusted p-value
2'FL	31.7 (21.0-69.8)	35.7 (23.5-67.6)	27.2 (17.8-98.0)	0.81
3FL	189.3 (130.8- 515.8)	235.6 (104.6- 553.2)	185.1 (178.6- 378.9)	0.81
LNnT	870.1 (585.6- 1203.5)	870.1 (587.4- 1257.5)	760.0 (585.6- 914.7)	0.67
3'SL	692.7 (474.1- 1091.4)	692.7 (430.9- 1101.7)	850.0 (601.5- 1059.8)	0.86
DFLac	56.1 (31.1-65.1)	53.4 (30.0-69.0)	59.2 (41.3-61.4)	0.63
6'SL	185.9 (149.7- 297.8)	185.9 (149.8- 336.9)	200.9 (65.9-243.9)	0.67
LNT	1330.7 (905.1- 2293.9)	1330.7 (841.1- 1986.6)	1704.9 (964.6- 2426.7)	0.47
LNFP I	91.5 (54.4-105.9)	90.1 (49.2-108.0)	97.5 (82.7-102.1)	0.90
LNFP II	2884.1 (1836.4- 3417.2)	2884.1 (2356.9- 3360.9)	2626.8 (1836.4- 3507.9)	0.95
LNFP III	93.8 (76.6-111.6)	89.9 (74.3-114.0)	98.0 (86.5-101.5)	0.81
LSTb	183.8 (123.1- 230.0)	205.1 (150.6- 231.2)	131.2 (21.4-185.3)	0.11
LSTc	41.3 (26.7-67.1)	41.3 (26.8-77.5)	48.1 (20.0-57.5)	0.81
DFLNT	619.0 (121.3- 839.7)	663.6 (197.8- 852.8)	341.1 (80.4-834.7)	0.43
LNH	80.0 (37.7-155.1)	109.7 (60.5- 179.2)	36.4 (26.5-54.4)	0.024
DSLNT	429.3 (315.3- 593.0)	425.9 (284.1- 610.7)	432.2 (346.2- 593.0)	0.76
FLNH	21.0 (15.2-34.2)	21.0 (16.5-32.1)	19.5 (15.1-86.2)	0.86

Title: Non-genetic mother-infant factors and Human milk oligossacharides composition/profile in breast milk of Ugandan mothers

First Author: Tonny Jimmy Owalla

DFLNH	9.5 (7.7-22.8)	10.1 (7.8-21.1)	8.3 (6.7-144.1)	0.95
FDSLNH	216.3 (21.2-554.2)	338.5 (21.1-588.0)	90.5 (38.4-247.4)	0.36
DSLNH	31.3 (7.9-77.4)	31.6 (7.9-81.8)	29.4 (6.8-35.7)	0.50
Total HMO	8988.5 (8159.9-9535.5)	9246.0 (8373.8-9568.8)	8450.6 (7941.6-8965.8)	0.13
Sia HMO	2984.1 (2380.1-3986.7)	3132.0 (2631.3-4261.7)	2395.0 (2311.8-2901.2)	0.10
Fuc HMO	5611.3 (3099.0-5958.7)	5611.3 (4120.2-5922.5)	4422.9 (3009.3-5966.2)	0.76
Diversity	5.3 (4.6-5.9)	5.3 (4.9-5.9)	4.6 (4.6-5.4)	0.13

IQR = interquartile range; HMOs = human milk oligosaccharides; N = total number of mothers; n = number of mothers in each weight category. HMO with a significant unadjusted p value is in bold.