

Supplementary Materials

Supplementary Table 1. Epigenetic age estimates of the study cohort

Epigenetic clock	Tissue	Mean (SD) , years*
Horvath clock	Adipose	-0.2 (-8.7–7.7)
	Muscle	-0.3 (-7.6–6.2)
Hannum clock	Adipose	-0.2 (-8.0–6.5)
	Muscle	-0.1 (-5.3–5.0)
GrimAge	Adipose	-0.2 (-5.8–4.1)
	Muscle	-0.1 (-3.7–4.0)
PhenoAge	Adipose	-0.2 (-11.5–11.8)
	Muscle	-0.2 (-8.1–4.6)
DunedinPACE*	Adipose	1.7 (1.5–2.0)
	Muscle	1.4 (1.2–1.5)
MEAT	Muscle	0.2 (-8.5–7.3)

* unit: years/calendar year

Supplementary Table 2. Characteristics of the monozygotic twin pairs used in ICE FALCON analysis of SH3BP4 methylation, mtDNA quantity and obesity-related traits in adipose tissue.

Variable	Heavier co-twin (mean)	Leaner co-twin (mean)	Difference (Heavier-Leaner)	P-value**	N (MZ twin pairs)
cg19998400* (M-value)	0.82	0.79	0.03	8.0E-05	68
mtDNAq*	-0.06	0.02	-0.08	7.7E-05	71
Weight (kg)	91.60	77.84	13.76	5.4E-18	71
BMI (kg/m ²)	31.93	27.32	4.61	4.6E-19	71
Waist circumference (cm)	104.38	91.96	12.42	3.6E-17	64
WHR	0.93	0.90	0.04	8.9E-05	33
Fat (%)	40.65	35.33	5.32	1.4E-13	71
Fat (kg)	38.36	28.79	9.57	3.0E-14	71
Fat-free mass (kg)	51.14	47.60	3.55	1.4E-07	71
Intra-abdominal fat (cm ³)	1300.57	625.16	675.42	7.7E-04	21
Subcutaneous fat (cm ³)	5933.65	3739.47	2194.18	3.8E-05	21
Adipocyte volume (mm ³)	615	457	159	7.3E-08	60
Liver fat (%)	4.15	1.60	2.55	9.9E-04	42
Total cholesterol (mmol/l)	4.82	4.73	0.09	0.47	71
HDL (mmol/l)	1.47	1.64	-0.18	1.7E-04	71
LDL (mmol/l)	3.12	2.91	0.22	0.07	71
Triglycerides (mmol/l)	1.12	0.97	0.16	0.01	71
hsCRP (mg/l)	3.33	2.31	0.02	0.02	64
Adipsin (ug/l)	1.21	1.14	0.07	0.12	21
Adiponectin (ng/l)	2903.00	3694.00	-791.00	7.7E-04	21
ALAT (U/l)	29.20	24.93	4.27	0.03	71
ASAT (U/l)	28.94	27.27	1.68	0.40	71
BP (systolic) (mmHg)	134	134	0	0.74	69
BP (diastolic) (mmHg)	81	79	2	0.03	69
Fasting glucose (mmol/l)	5.82	5.55	0.27	0.11	69
Fasting insulin (mU/l)	9.50	7.28	2.22	3.0E-03	67
HOMA-IR	2.52	1.81	0.71	2.5E-03	64
Matsuda index	5.58	7.76	-2.18	7.8E-05	59
Leisure-time physical activity	2.91	2.92	-0.02	0.86	65
Sports activity	2.59	2.77	-0.18	0.12	66

Work activity	2.45	2.44	0.02	0.91	67
Total activity	7.98	8.17	-0.19	0.29	63
EAAHorvath*	0.14	-0.47	0.61	0.02	68
EAAHannum*	0.35	-0.49	0.84	5.5E-04	68
EAAGrim*	-0.43	-0.03	-0.39	0.05	68
EAAPheno*	0.26	-0.34	0.60	0.08	68
DunedinPACE*	1.70	1.69	0.00	0.80	68

* Estimates derived from adipose tissue

** Paired t-test (two-tailed)

ALAT = Alanine aminotransferase; ASAT = Aspartate aminotransferase; BMI = body mass index; DZ = dizygotic; EAA = Epigenetic Age Acceleration; HOMA-IR = Homeostatic model assessment for insulin resistance; hsCRP = high-sensitivity C-reactive protein; mtDNAq = mitochondrial DNA quantity; MZ = monozygotic; WHR = waist-to-hip ratio

Supplementary Table 3. Results from the causal inference using ICE FALCON between the expression and methylation of SH3BP4 and DHRS3 (n=40 monozygotic twin pairs).

Formula	Coefficient	Model 1			Model 2			Model 3			Change	
		Est	SE	p.val	Est	SE	p.val	Est	SE	p.val	Est	p.val
SH3BP4 ~ cg19998400	Bself	0.261	0.047	2.0E-08				0.267	0.047	1.3E-08	0.006	0.871
	Bcotwin				-0.112	0.064	0.078	0.014	0.066	0.833	0.126	0.134
cg19998400 ~ SH3BP4	Bself	0.652	0.186	4.7E-04				0.717	0.163	1.0E-05	0.065	0.494
	Bcotwin				0.031	0.162	0.846	-0.225	0.173	0.193	-0.256	0.299
DHRS3 ~ cg17468563	Bself	-0.643	0.156	3.7E-05				-0.657	0.146	6.7E-06	-0.014	0.822
	Bcotwin				-0.050	0.191	0.793	0.075	0.165	0.647	0.125	0.532
cg17468563 ~ DHRS3	Bself	-0.296	0.072	3.7E-05				-0.306	0.071	1.5E-05	-0.010	0.696
	Bcotwin				-0.008	0.078	0.916	-0.059	0.055	0.280	-0.051	0.581

Supplementary Table 4. Associations between adipose tissue SH3BP4 methylation and obesity-related traits.

Outcome	Coef	SE	P-value	FDR	N
Weight	0.246	0.070	4.1E-04	0.002	136
BMI	0.283	0.078	2.9E-04	0.001	136
Waist	0.300	0.065	4.0E-06	3.5E-05	122
WHR	0.123	0.062	0.050	0.079	64
Fat %	0.290	0.062	2.7E-06	4.7E-05	136
Fat (kg)	0.318	0.077	3.4E-05	2.3E-04	136
Fat-free mass	0.051	0.039	0.190	0.247	136
Ia. Fat ^a	0.458	0.099	3.6E-06	4.1E-05	42
Subcut. Fat ^a	0.352	0.086	4.5E-05	2.6E-04	42
Adipocyte volume	0.202	0.079	0.010	0.022	114
Liver fat % ^a	0.391	0.080	1.1E-06	3.8E-05	82
Total cholesterol	0.027	0.088	0.761	0.807	136
HDL	-0.189	0.068	0.006	0.015	136
LDL	0.059	0.092	0.523	0.631	136
Triglycerides ^a	0.222	0.090	0.013	0.028	136
hsCRP ^a	0.320	0.081	7.2E-05	3.6E-04	122
Adipsin	0.058	0.146	0.694	0.759	42
Adiponectin	-0.275	0.089	0.002	0.006	42
ASAT ^a	0.033	0.077	0.672	0.784	136
ALAT ^a	0.095	0.088	0.282	0.353	136
BP (systolic)	0.089	0.057	0.122	0.165	132
BP (diastolic)	0.206	0.063	0.001	0.003	132
Fasting glucose ^a	0.165	0.074	0.025	0.046	132
Fasting insulin ^a	0.233	0.097	0.016	0.032	128
HOMA-IR ^a	0.250	0.095	0.009	0.021	122
Matsuda ^a	-0.303	0.089	0.001	0.002	112
LTPA	-0.246	0.092	0.008	0.019	124
Sports activity	-0.178	0.088	0.042	0.074	126
Work activity	0.026	0.097	0.793	0.816	128
Total activity	-0.169	0.086	0.049	0.082	120
EAAHorvath	0.014	0.070	0.837	0.837	136
EAAHannum	0.134	0.070	0.055	0.080	136
EAAGrim	-0.116	0.069	0.090	0.126	136
EAAPheno	0.026	0.063	0.678	0.766	136
DunedinPACE	-0.156	0.080	0.053	0.081	136

^a Log10 transformed

Supplementary Table 5. Associations between adipose tissue mtDNA quantity and obesity-related traits.

Outcome	Coef	SE	P-value	FDR	N
Weight	-0.346	0.096	3.2E-04	0.001	142
BMI	-0.340	0.095	3.5E-04	0.001	142
Waist	-0.320	0.095	0.001	0.002	128
WHR	-0.175	0.095	0.066	0.096	66
Fat %	-0.239	0.073	0.001	0.002	142
Fat (kg)	-0.357	0.091	9.4E-05	0.001	142
Fat-free mass	-0.184	0.061	0.002	0.006	142
la. Fat ^a	-0.578	0.117	7.5E-07	2.6E-05	42
Subcut. Fat ^a	-0.464	0.105	1.1E-05	1.2E-04	42
Adipocyte volume	-0.322	0.085	1.6E-04	0.001	120
Liver fat % ^a	-0.366	0.123	0.003	0.006	84
Total cholesterol	0.076	0.087	0.382	0.432	142
HDL	0.324	0.089	2.7E-04	0.001	142
LDL	-0.062	0.092	0.505	0.535	142
Triglycerides ^a	-0.159	0.108	0.142	0.184	142
hsCRP ^a	-0.194	0.109	0.074	0.103	128
Adipsin	-0.229	0.100	0.023	0.040	42
Adiponectin	0.357	0.098	2.7E-04	0.001	42
ALAT ^a	-0.095	0.099	0.337	0.393	142
ASAT ^a	-0.091	0.092	0.322	0.389	142
BP (systolic)	0.025	0.074	0.739	0.761	138
BP (diastolic)	-0.163	0.085	0.057	0.086	138
Fasting glucose ^a	-0.097	0.085	0.258	0.322	138
Fasting insulin ^a	-0.296	0.103	0.004	0.009	134
HOMA-IR ^a	-0.355	0.093	1.4E-04	0.001	128
Matsuda ^a	0.345	0.095	2.8E-04	0.001	118
LTPA	0.183	0.094	0.051	0.080	130
Sports activity	0.260	0.101	0.010	0.020	132
Work activity	-0.068	0.092	0.458	0.501	134
Total activity	0.194	0.093	0.038	0.063	126
EAAHorvath	-0.176	0.073	0.016	0.030	136
EAAHannum	-0.273	0.072	1.4E-04	0.001	136
EAGrim	0.145	0.091	0.110	0.148	136
EAAPheno	-0.252	0.054	3.2E-06	5.5E-05	136
DunedinPACE	-0.004	0.094	0.964	0.964	136


^a Log₁₀ transformed

Supplementary Figure 1. Scatter plot representing the regression coefficients from EWAS on mtDNA quantity in adipose tissue (X) and skeletal muscle (Y).


Supplementary Figure 2. The amount of methylation probe quantile variation unexplained by the control probe principal components under 10-fold cross validation in (a) adipose tissue and (b) skeletal muscle. The number of principal components which the probe quantiles were normalized with is indicated in the plots.

M = methylated probes; U = unmethylated probes

Supplementary Figure 3. First four principal components of the adipose tissue DNA methylation data indicated by beadchip type (450K/EPIC), (a) before and (b) after adjustment with *ComBat*.



a



b

