

**Supplemental Table 1. Demographic Information**

	No history of suicide attempt (N=51)	History of suicide attempt (N=55)	Overall (N=106)	P-value
<b>Sex</b>				0.132 <sup>a</sup>
<b>Male</b>	22 (43%)	15 (27%)	37 (35%)	
<b>Female</b>	29 (57%)	40 (73%)	69 (65%)	
<b>Age (years)</b>				0.308 <sup>b</sup>
<b>Mean (SD)</b>	39.4 (13.7)	42.1 (13.6)	40.8 (13.6)	
<b>Median [Min, Max]</b>	37.0 [18.0, 66.0]	41.0 [18.0, 68.0]	41.0 [18.0, 68.0]	
<b>Race</b>				0.404 <sup>c</sup>
<b>Am. Indian/Alaskan Native</b>	1 (2%)	1 (2%)	2 (2%)	
<b>Asian</b>	1 (2%)	0 (0%)	1 (1%)	
<b>Black/African American</b>	1 (2%)	5 (9%)	6 (6%)	
<b>More than one race</b>	1 (2%)	2 (4%)	3 (3%)	
<b>Unknown</b>	0 (0%)	0 (0%)	0 (0%)	
<b>White</b>	47 (92%)	47 (86%)	94 (89%)	
<b>Smoking history</b>				<b>0.024<sup>c</sup></b>
<b>Current smoker</b>	5 (10%)	14 (26%)	19 (18%)	
<b>Missing</b>	0 (0%)	1 (2%)	1 (1%)	
<b>Never smoker</b>	28 (55%)	17 (31%)	45 (43%)	
<b>Past smoker</b>	18 (35%)	23 (42%)	41 (39%)	
<b>Body Mass Index (BMI)</b>				0.605 <sup>b</sup>
<b>Mean (SD)</b>	30.3 (7.61)	31.0 (7.39)	30.7 (7.47)	
<b>Median [Min, Max]</b>	29.2 [18.4, 59.0]	31.0 [18.9, 50.5]	29.9 [18.4, 59.0]	
<b>Number of lifetime suicide attempts</b>				<b>1.831x10<sup>-6b</sup></b>
<b>Mean (SD)</b>	0 (0)	3.56 (4.94)	1.85 (3.97)	
<b>Median [Min, Max]</b>	0 [0, 0]	2.00 [1.00, 30.0]	1.00 [0, 30.0]	

Descriptive statistics for the bipolar disorder cohort separated by history of suicide attempt.

<sup>a</sup>Pearson's Chi-squared test with Yates' continuity correction; <sup>b</sup>Welch's Two Sample T-Test;

<sup>c</sup>Fisher's Exact Test. Significant differences are indicated in bold. SD, standard deviation.

**Supplemental Table 2. Correlation between DNA methylation and number of lifetime suicide attempts**

Gene	CpG Genomic Location	Statistic	P-value	False Discovery Rate (FDR)
MIF	chr22:23894125	-3.664	2.49x10 <sup>-4</sup>	0.022
IMPA2	chr18:11994457	-3.136	1.71x10 <sup>-3</sup>	0.075
INPP1	chr2:190354545	-2.806	5.02x10 <sup>-3</sup>	0.147
IRX2*	chr5:2468211	-2.645	8.17x10 <sup>-3</sup>	0.177
NOS3	chr7:150998982	-2.573	0.010	0.177
CRHR1	chr17:45805706	-2.395	0.017	0.195
MIF	chr22:23894167	2.287	0.022	0.195
MIF	chr22:23894177	2.323	0.020	0.195
IRX2*	chr5:2468198	-2.341	0.019	0.195
TGFBR1	chr9:99146591	-2.430	0.015	0.195
ACP1	chr2:273567	2.198	0.028	0.224
INPP1	chr2:190354502	-2.145	0.032	0.235
OXTR	chr3:8762680	-2.055	0.040	0.270
ANK3	chr10:60124978	-1.948	0.051	0.283
BDNF	chr11:27658429	-1.984	0.047	0.283
INPP1	chr2:190354478	-1.966	0.049	0.283

Results with an uncorrected P-value less than 0.05 shown. GRCh38/hg38 used to annotate genomic locations. \*Closest gene if variant is intergenic.

**Supplemental Table 3. Correlation between DNA methylation and severity of most lethal attempt**

<b>Gene</b>	<b>CpG Genomic Location</b>	<b>Statistic</b>	<b>P-value</b>	<b>False Discovery Rate (FDR)</b>
CACNA1C	chr12:2613346	-3.459	5.43x10 <sup>-4</sup>	0.048
CACNA1C	chr12:2613360	-2.912	3.60x10 <sup>-3</sup>	0.158
BDNF	chr11:27658332	-2.612	9.01x10 <sup>-3</sup>	0.264
ACP1	chr2:273531	2.382	0.017	0.318
OXTR	chr3:8762680	-2.365	0.018	0.318
IMPA2	chr18:11994474	-2.118	0.034	0.502
IMPA2	chr18:11994457	-2.021	0.043	0.545

Results with an uncorrected P-value less than 0.05 are shown. GRCh38/hg38 used to annotate genomic locations.