






























# Supplementary Table 2. Vascular biomarkers

Biomarker	Description	Clinical significance
<b>Ang-1</b>	Vascular growth factor. Regulates angiogenesis, endothelial cell survival, and proliferation. Mediates blood vessel maturation and stability.	 A decline in Ang-1 plasma levels was associated with increasing <i>falciparum</i> and <i>vivax</i> malaria severity and widespread endothelial activation across African and Asian patient studies, irrespective of age (1–5). 
<b>Ang-2</b>	Vascular growth factor, biomarker of endothelial activation. Modulates Ang-1 signalling, and in the absence of angiogenic inducers such as VEGF, promotes vascular regression. In concert with VEGF, triggers a permissive angiogenic signal. Involved in lymphangiogenesis.	 An increase of Ang-2 plasma levels was associated with increasing <i>falciparum</i> and <i>vivax</i> malaria severity and widespread endothelial activation across African and Asian patient studies, irrespective of age (1,3,5).   Robust predictor of mortality in <i>falciparum</i> cerebral malaria, identified as a risk factor for blood-brain barrier dysfunction, neuroinflammation, and long-term cognitive injury in African children (4,6,7). 
<b>Ang2 Ang1</b>	Ratio between Ang-2 and Ang-1 plasma levels.	 Plasma ratio was higher in patients with <i>falciparum</i> and <i>vivax</i> severe malaria compared with uncomplicated malaria and healthy controls, across African, Asian, and Latin American patient studies and irrespective of age.  Fatal cerebral malaria cases showed the highest ratio (1,2,5,8–10).
<b>BMP-9</b>	Growth factor, member of the TGF- $\beta$ superfamily. Regulates angiogenesis by inhibiting VEGF-induced endothelial cell migration and proliferation.	To our knowledge, no studies have reported significant group differences in the context of human malaria.
<b>ICAM-1</b>	Intercellular adhesion molecule constitutively expressed on the vascular endothelium. Upon IL-1 and TNF- $\alpha$ stimulation, expression levels increase, so that leukocytes can bind to it and transmigrate into tissues. Increased plasma levels are suggestive of endothelial activation.	 Plasma levels were significantly higher in Ugandan children with <i>falciparum</i> severe malaria, compared with healthy controls (4).  Plasma levels were higher in Malawian children with <i>falciparum</i> cerebral malaria and retinopathy, compared with their retinopathy-negative counterparts (11).  In Ugandan children with <i>falciparum</i> malaria, plasma levels were elevated in severe malarial anaemia fatalities compared to survivors (12).  In Ghanaian children with <i>falciparum</i> cerebral malaria, plasma levels were higher compared with uncomplicated malaria cases (13).  In Pakistani patients with <i>vivax</i> malaria, plasma levels were significantly higher in uncomplicated cases than healthy controls, and higher again in complicated cases compared with uncomplicated ones. ICAM-1, VCAM-1, TNF- $\alpha$ and IL-10 were the best individual predictors of complicated <i>vivax</i> malaria (14).
<b>PDGF-AA</b>	Angiogenic promoter. Plays an important role in wound healing, and it is essential during embryonic development.	 Increased plasma levels predicted abnormal cerebral blood flow and stroke in children with sickle cell disease who presented with cerebrovascular disease (15).
<b>PDGF-BB</b>	Angiogenic promoter. Participates in wound healing, blood vessel development, and proliferation and recruitment of pericytes and vascular smooth muscle cells in the central nervous system.	 In Ugandan children with <i>falciparum</i> cerebral malaria, plasma levels were significantly higher in retinopathy-negative patients, compared to their retinopathy-positive peers (16). 
<b>Serpin E1</b>	Serine protease inhibitor. Plays a role in the controlled degradation of blood clots.	To our knowledge, no studies have reported significant group differences in the context of human malaria.
<b>VCAM-1</b>		 Plasma levels were significantly higher in Ugandan children with <i>falciparum</i> severe malaria, compared with healthy controls (4).

	Surface glycoprotein expressed on the vascular endothelium. Participates in immune surveillance and inflammation by regulating leukocyte adhesion to the endothelium and transendothelial migration.		In Pakistani patients with <i>vivax</i> malaria, plasma levels were significantly higher in uncomplicated cases than healthy controls, and higher again in complicated cases compared with uncomplicated ones. VCAM-1, ICAM-1, TNF- $\alpha$ and IL-10 were the best individual predictors of complicated <i>vivax</i> malaria (14).
<b>VEGF</b>	Vascular growth factor. Promotes angiogenesis, vasculogenesis, and endothelial cell growth. Induces endothelial cell proliferation, cell migration, and permeabilization of blood vessels.	 	Plasma levels were significantly higher in Indonesian patients with <i>falciparum</i> malaria compared with healthy controls (17). Serum levels were significantly lower in Indian patients with <i>vivax</i> severe malaria compared with uncomplicated malaria and healthy controls (5).
<b>vWF-A2</b>	Glycoprotein involved in haemostasis, biomarker of endothelial activation. Promotes adhesion of platelets to sites of vascular injury and acts as a chaperone for certain coagulation factors.	   	Plasma levels were significantly higher in Ugandan children with <i>falciparum</i> severe malaria, compared with healthy controls (4); increased plasma levels were associated with mortality (18). Plasma levels were significantly higher in Malawian children with cerebral malaria than in children with uncomplicated malaria, showing similar values in patients with and without retinopathy (19). Plasma levels were significantly increased in Malaysian patients with severe and uncomplicated <i>vivax</i> malaria, compared with controls, and correlated with parasitaemia in these patients (20).

**Ang-1:** angiopoietin-1; **Ang-2:** angiopoietin-2; **Ang-2/Ang-1:** ratio between Ang-2 and Ang-1; **BMP-9:** bone morphogenetic protein 9; **ICAM-1:** intercellular adhesion molecule 1; **PDGF-AA:** platelet-derived growth factor AA; **PDGF-BB:** platelet-derived growth factor BB; **VCAM-1:** vascular cell adhesion molecule; **Serpin E1:** Serine Proteinase Inhibitor-clade E1; **PEDF:** pigment epithelium derived factor; **VEGF:** vascular endothelial growth factor; **vWF-A2:** von Willebrand Factor (A2 domain).

**Legend:** Clinical findings related to...

 *Plasmodium falciparum* malaria infection;  *Plasmodium vivax* malaria infection;  *Plasmodium knowlesi* malaria infection;  
 Neurological conditions;  Other medical conditions.

## References:

1. De Jong GM, Slager JJ, Verbon A, *et al.* Systematic review of the role of angiotensin-1 and angiotensin-2 in *Plasmodium* species infections: biomarkers or therapeutic targets? *Malar J* (2016) 15(1):1–12.
2. Jain V, Lucchi NW, Wilson NO, *et al.* Plasma levels of angiotensin-1 and -2 predict cerebral malaria outcome in Central India. *Malar J* (2011) 10(1):1–7.
3. Conroy AL, Glover SJ, Hawkes M, *et al.* Angiotensin-2 levels are associated with retinopathy and predict mortality in Malawian children with cerebral malaria: A retrospective case-control study. *Crit Care Med* (2012) 40(3):952–9.
4. Ouma BJ, Ssenkusu JM, Shabani E, *et al.* Endothelial Activation, Acute Kidney Injury, and Cognitive Impairment in Pediatric Severe Malaria. *Crit Care Med* (2020) 48(9):E734–43.
5. Gowda S, Anghan H, Mishra H, *et al.* Serum Angiotensin-1 and -2 and VEGF are associated with severe disease in *vivax* malaria. *J Vector Borne Dis* (2020) 57(4):285–94.
6. Petersen JEV, Mkumbaye SI, Vaaben A V, *et al.* Plasma Ang2 and ADAM17 levels are elevated during clinical malaria; Ang2 level correlates with severity and expression of EPCR-binding PfEMP1. *Scientific Reports* (2016) 6(1):1–9.
7. Ouma BJ, Bangirana P, Ssenkusu JM, *et al.* Plasma angiotensin-2 is associated with age-related deficits in cognitive sub-scales in Ugandan children following severe malaria. *Malar J* (2021) 20(1):1–10.
8. Gomes LT, Alves- ER, Rodrigues-Jesus C, *et al.* Angiotensin-2 and Angiotensin-2/Angiotensin-1 Ratio as Indicators of Potential Severity of *Plasmodium vivax* Malaria in Patients with Thrombocytopenia. *PLoS One* (2014) 9(10):e109246.
9. Jain V, Thomas T, Basak S, *et al.* Sequential dysregulated plasma levels of angiotensins (ANG-2 and ratios of ANG-2/ANG-1) are associated with malaria severity and mortality among hospital admitted cases in South Bastar Region of Chhattisgarh, Central India. *Pathog Glob Health* (2022) 116(1):47–58.
10. Oluboyo AO, Chukwu SI, Oluboyo BO, Odewusi OO. Evaluation of Angiotensins 1 and 2 in Malaria-Infested Children. *J Environ Public Health*. 2020;2020.
11. Conroy AL, Phiri H, Hawkes M, Glover S, Mallewa M, Seydel KB, *et al.* Endothelium-Based Biomarkers Are Associated with Cerebral Malaria in Malawian Children: A Retrospective Case-Control Study. *PLoS One* [Internet]. 2010 [cited 2024 Jan 26];5(12):e15291. Available from: <https://journals.plos.org/plosone/article?id=10.1371/journal.pone.0015291>
12. Erdman LK, Dhabangi A, Musoke C, Conroy AL, Hawkes M, Higgins S, *et al.* Combinations of Host Biomarkers Predict Mortality among Ugandan Children with Severe Malaria: A Retrospective Case-Control Study. *PLoS One* [Internet]. 2011 [cited 2024 Jan 26];6(2):e17440. Available from: <https://journals.plos.org/plosone/article?id=10.1371/journal.pone.0017440>
13. Adukpo S, Kusi KA, Ofori MF, Tetteh JKA, Amoako-Sakyi D, Goka BQ, *et al.* High Plasma Levels of Soluble Intercellular Adhesion Molecule (ICAM)-1 Are Associated with Cerebral Malaria. *PLoS One* [Internet]. 2013 Dec 27 [cited 2024 Jan 26];8(12):e84181. Available from: <https://journals.plos.org/plosone/article?id=10.1371/journal.pone.0084181>
14. Raza A, Ghanchi NK, Zubairi ABS, *et al.* Tumor Necrosis Factor - $\alpha$ , Interleukin-10, Intercellular and Vascular Adhesion Molecules Are Possible Biomarkers of Disease Severity in Complicated *Plasmodium vivax* Isolates from Pakistan. *PLoS One* (2013) 8(12):e81363.
15. Hyacinth HI, Gee BE, Adamkiewicz T V, *et al.* Plasma BDNF and PDGF-AA levels are associated with high TCD velocity and stroke in children with sickle cell anemia. *Cytokine* (2012) 60(1):302–8.
16. Villaverde C, Namazzi R, Shabani E, *et al.* Retinopathy-Positive Cerebral Malaria Is Associated With Greater Inflammation, Blood-Brain Barrier Breakdown, and Neuronal Damage Than Retinopathy-Negative Cerebral Malaria. *J Pediatric Infect Dis Soc* (2020) 9(5):580–6.
17. Brouwers J, Noviyanti R, Fijnheer R, *et al.* Platelet Activation Determines Angiotensin-1 and VEGF Levels in Malaria: Implications for Their Use as Biomarkers. *PLoS One* (2013) 8(6):e64850.
18. Graham SM, Chen J, Chung DW, *et al.* Endothelial activation, haemostasis and thrombosis biomarkers in Ugandan children with severe malaria participating in a clinical trial. *Malar J* (2016) 15(1):1–9.
19. Phiri HT, Bridges DJ, Glover SJ, *et al.* Elevated Plasma Von Willebrand Factor and Propeptide Levels in Malawian Children with Malaria. *PLoS One* (2011) 6(11):e25626.
20. Barber BE, William T, Grigg MJ, *et al.* Parasite Biomass-Related Inflammation, Endothelial Activation, Microvascular Dysfunction and Disease Severity in *Vivax* Malaria. *PLoS Pathog* (2015) 11(1):e1004558.