

Supplementary Material

1 STABILITY OF NON-TRIVIAL STEADY STATE OF CC-ODE MODEL

The stability condition for the non-trivial steady state is computed with the help of the eigenvalues of the Jacobian at the non-trivial steady state:

$$\lambda_{1/2} = \frac{-\delta(\beta_1\beta_2 + \delta)}{(2\alpha\beta_2 + \gamma + \delta)} \pm \frac{\delta^{0.5}}{(2\alpha\beta_2 + \gamma + \delta)} \cdot \left(4\alpha^3\beta_2^3 - 4\alpha^2\beta_1\beta_2^3 + 12\alpha^2\gamma\beta_2^2 + \right. \quad (\text{S1})$$

$$\left. 8\alpha^2\delta\beta_2^2 - 8\alpha\beta_1\gamma\beta_2^2 - 8\alpha\beta_1\delta\beta_2^2 + 12\alpha\gamma^2\beta_2 + 16\alpha\gamma\delta\beta_2 + 4\alpha\delta^2\beta_2 + \beta_1^2\delta\beta_2^2 - \right. \quad (\text{S2})$$

$$\left. 4\beta_1\gamma^2\beta_2 - 8\beta_1\gamma\delta\beta_2 - 2\beta_1\delta^2\beta_2 + 4\gamma^3 + 8\gamma^2\delta + 4\gamma\delta^2 + \delta^3 \right)^{0.5}, \quad (\text{S3})$$

$$\lambda_3 = 0. \quad (\text{S4})$$

For the choice of $\delta = 0.1$, we obtain the following behaviour depending on α , β_1 , β_2 and γ .

Generally speaking, the non-trivial steady state is stable for low values of α and γ and high values of β_1 and β_2 . The latter parameters represent the rate of spread of the disease and the higher they are, the more likely it will be for the disease to remain present in the population for a longer time period. Hence, the




Figure S1. Stability of non-trivial steady state for the parameters $\beta_1 \in \{0.5, 0.7, 1.0\}$, $\gamma \in \{0.1, 0.2, 0.3\}$ and $\delta = 0.1$

non-trivial state that includes a constant proportion of infected individuals will be stable and there will always be infections. In contrast, the parameters α and γ generate the flux towards the non-infected state **R** and finally to **S**, if $\delta > 0$.