

12 **Kinesiophobia and Physical Activity: A Systematic Review and Meta-Analysis**

13

14

ABSTRACT

15 **Objective.** Physical activity contributes to the primary, secondary, and tertiary prevention of
16 multiple diseases. However, in some patients, an excessive, irrational, and debilitating fear of
17 movement (i.e., kinesiophobia) is thought to induce avoidance behaviors, contributing to decreased
18 engagement in physical activity. The aim of this study was to examine whether kinesiophobia is
19 negatively associated with physical activity in several health conditions and what factors may
20 influence this relationship.

21 **Methods.** Five databases were searched for studies including both a measure of kinesiophobia and
22 physical activity. Two reviewers screened articles for inclusion, assessed risk of bias, and extracted
23 data from each study. Pearson product-moment correlations were pooled from eligible studies
24 using the generic inverse pooling and random effects method to examine the relationship between
25 kinesiophobia and physical activity.

26 **Results.** Seventy-four studies were included in the systematic review and 63 studies (83 estimates,
27 12,278 participants) in the main meta-analysis. Results showed a small-to-moderate negative
28 correlation between kinesiophobia and physical activity ($r = -0.19$; 95% confidence interval: -0.26
29 to -0.13 ; $I^2 = 85.5\%$; $p < 0.0001$). Funnel plot analysis showed evidence of publication bias, but p-
30 curve analysis suggested that our results could not be caused by selective reporting. A subgroup
31 meta-analysis showed that the correlation was statistically significant in patients with cardiac,
32 rheumatologic, neurologic, or pulmonary conditions, but not in patients with chronic or acute pain.

33 **Conclusion.** Our results suggest that higher levels of kinesiophobia are associated with lower
34 levels of physical activity in several health conditions that are not necessarily painful.

35 **Impact.** Kinesiophobia should be dissociated from pain and considered in relation to specific
36 health conditions when implementing interventions to promote physical activity. Kinesiophobia
37 may have prognostic implications in patients for whom physical activity contributes to prevent
38 recurrence or worsening of their condition.

39

40 **Keywords.** Exercise, Health Status; Pain; Prevention; Psychology; Rehabilitation

41

42 **Disclosures:** None

43 **Kinesiophobia and Physical Activity: A Systematic Review and Meta-Analysis**

44

45

INTRODUCTION

46 Seven decades ago, the seminal work of Morris et al. (1953)¹ showed that conductors on London
47 double-decker buses, who were responsible for checking tickets, assisting passengers with
48 luggage, and supervising the loading and unloading of passengers, had a lower incidence and less
49 severe coronary heart disease than bus drivers. Since then, the scientific literature demonstrating
50 the health benefits of physical activity has grown exponentially and expanded to include multiple
51 health conditions². These benefits include reduced risk of disability, disease, and mortality^{2,3}.
52 Specifically, higher levels of physical activity have been shown to contribute to a reduced risk of
53 cardiovascular disease⁴, obesity⁵, depression⁶, hypertension⁷, cancer⁸, and dementia⁹. Yet, one in
54 four adults worldwide does not meet the recommendations for physical activity¹⁰. Physical activity
55 also plays an important role in secondary and tertiary prevention by reducing the impact, slowing
56 the progression, and preventing the recurrence of multiple health conditions, including
57 cardiovascular disease^{11,12}, osteoarthritis¹³, stroke^{14,15}, and cancer¹⁶.
58 Several factors may explain physical inactivity¹⁷, including environmental, interpersonal, and
59 intrapersonal factors¹⁸. Environmental factors include lack of access, weather conditions, and
60 safety concerns¹⁹. Interpersonal factors include family responsibilities, lack of support, and lack
61 of a gym partner²⁰. Intrapersonal factors include gender²¹, age²², cognitive function^{23,24}, and
62 socioeconomic circumstances²⁵. Another intrapersonal factor of interest is kinesiophobia, which
63 can be defined as an excessive, irrational, and debilitating fear of movement and activity resulting
64 from a sense of vulnerability to pain, injury, or a medical condition²⁶. Kinesiophobia is typically
65 measured using self-administered questionnaires, such as the Tampa Scale of Kinesiophobia

66 (TSK)^{27,28}, which assesses an individual's belief that physical activity can lead to injury or pain
67 and that the severity of their medical condition is underestimated. While kinesiophobia is often
68 observed in the context of pain or a clinical condition, its presence in otherwise healthy adults is
69 also possible²⁹ due to the irrational nature of this phobic condition. The irrational fear that
70 characterizes kinesiophobia is likely to influence the desires and impulses for movement and rest³⁰,
71 as well as affective determinants of physical activity in general³¹.

72 The relationship between kinesiophobia and physical activity can be explained by theories
73 suggesting that the perception of a cue related to physical activity automatically activates the
74 concept of physical activity as well as the unpleasant (or pleasant) affective memories associated
75 with this concept³²⁻³⁵. This activation results in an impulse that favors the tendency to avoid (or
76 approach) physical activity³⁶. Thus, negative affective associations are likely to hinder physical
77 activity. Accordingly, an aversive fear of pain, injury, or aggravation of a medical condition that
78 has been associated with the concept of movement may result in the development of automatic
79 avoidance behaviors that contribute to the maintenance and exacerbation of this fear, and
80 ultimately lead to a phobic state (i.e., kinesiophobia) that diminishes the ability to engage in regular
81 physical activity.

82 Previous systematic syntheses of the literature on this topic include a meta-analysis³⁷ and two
83 systematic reviews^{38,39}. The main results of these reviews suggest that exercise interventions may
84 reduce kinesiophobia in individuals with back pain. While back pain is one condition that may
85 contribute to kinesiophobia, it is not the only one. The relationship between physical activity and
86 kinesiophobia should be investigated in other conditions such as cardiac, neurological, and
87 rheumatologic conditions.

88 The main objective of this study was to systematically review and meta-analyze the direct
89 relationship between kinesiophobia and physical activity. We hypothesized that levels of
90 kinesiophobia would be negatively associated with levels of physical activity. In addition, we
91 examined the moderating effect of health status, physical activity measurement instruments (i.e.,
92 accelerometers, pedometers, questionnaires), physical activity outcomes (e.g., total physical
93 activity, moderate or vigorous physical activity, steps per day), and kinesiophobia measurement
94 instruments. Finally, because kinesiophobia and physical activity can vary with age, sex, and
95 pain^{40,41}, we explored the influence of these factors on the association between kinesiophobia and
96 physical activity.

97

98

METHODS

99 Search Strategy

100 This review was reported according to the Preferred Reporting Items for Systematic Reviews and
101 Meta-Analyses (PRISMA) guidelines⁴². Potential studies were identified by searching the
102 MEDLINE (via PubMed), PsychInfo, CINAHL, EMBASE, and SPORTDiscus databases. In
103 October 2023, two reviewers (MG and AF) searched for all available records using the following
104 combination of keywords in the title or abstract of the article: (“kinesiophobia” OR “fear
105 avoidance” OR “fear of movement” OR “movement phobia” OR “movement fear”) AND
106 (“physical activity” OR “exercise” OR “walking”). In PsychInfo the limits “clinical trial”,
107 “quantitative study”, “peer-reviewed journal”, “English”, and “human” were used. In PubMed the
108 limits “clinical trial”, “observational study”, “RCT”, “English” were used. In SPORTDiscus the
109 limits “peer-reviewed”, “English”, “academic journal”, and “article” were used. In CINAHL the

110 limits “peer-reviewed”, “English”, “research article”, “journal article”, and “humans” were used.
111 To reduce literature bias^{43,44}, this systematic review was pre-registered in PROSPERO⁴⁵.

112

113 **Eligibility Criteria and Study Selection**

114 *Inclusion Criteria*

115 To be included in this systematic review, articles had to be published in a peer-reviewed journal,
116 be written in English, report original data collected from human participants, include at least one
117 self-reported measure of kinesiophobia and one measure of physical activity, and formally test the
118 association between these two variables, be it a univariate or multivariate test. The physical activity
119 measure could be derived from a self-reported measure of the level of physical activity or from a
120 device (e.g., accelerometer, pedometer) worn while participants are engaged in their normal daily
121 activities.

122

123 *Exclusion Criteria*

124 Studies were excluded if they were published as a book chapter, study protocol, conference
125 abstract, or were based on laboratory-based measures of physical fitness (e.g., maximal muscle
126 force, $\dot{V}O_2$ max) and not on a measure of physical activity.

127

128 *Study Selection*

129 Article screening was performed in Covidence systematic review software (Veritas Health
130 Innovation, Melbourne, Australia; www.covidence.org), a web-based collaborative software
131 platform that streamlines the production of systematic reviews. After removing duplicates, titles
132 and abstracts were independently reviewed by two reviewers (MG, AF) according to the inclusion

133 and exclusion criteria using a systematic 5-step process. If there was any doubt at any step, the full
134 text was further reviewed. Step 1: Articles not written in English were excluded. Step 2: Articles
135 that did not report original empirical data were excluded (e.g., reviews, meta-analyses,
136 commentaries, technical reports, case studies). Step 3: Articles that did not involve human
137 participants were excluded. Step 4: Articles that did not assess both kinesiophobia and physical
138 activity were excluded. Step 5: Articles that did not formally test the association between
139 kinesiophobia and physical activity were excluded. In addition, we performed reference screening
140 and forward citation tracking on the articles remaining after step 5. Any disagreements between
141 the two reviewers were resolved by consensus among three reviewers (MG, AF, MPB).

142

143 **Data Extraction**

144 Data extracted from selected articles included first author's name, article title, publication year,
145 digital object identifier (DOI), number of participants, number of men and women, age range,
146 mean age, mean weight, mean height, mean body mass index, health status, mean pain intensity,
147 type of kinesiophobia measure, level of kinesiophobia, type of physical activity measure, type of
148 physical activity outcome, level of physical activity (continuous or categorical), as well as
149 statistical estimates and significance of the association between kinesiophobia and physical
150 activity.

151

152 **Methodological Quality and Risk of Bias Assessment**

153 The risk of bias of the studies included in the systematic review was estimated using the National
154 Institutes of Health (NIH) Quality Assessment Tool for Observational Cohort and Cross-Sectional
155 Studies⁴⁶, the Transparent Reporting of Evaluations with Non-Randomized Designs (TREND)

156 reporting checklist⁴⁷, and the Consolidated Standards of Reporting Trials (CONSORT) reporting
157 checklist for randomized trials⁴⁸. All scores were normalized to a 0-10 scale to make them
158 comparable across assessment instruments (Table 1).

159

160 **Meta-Analysis**

161 All analyses were performed in R Studio integrated development environment (IDE)
162 (2023.06.1+524, “Mountain Hydrangea” release) for R software environment⁴⁹ using the {meta}⁵⁰
163 and {metafor}^{51,52} R packages⁵³.

164

165 *Main Meta-Analysis*

166 We pooled Pearson product-moment correlations from eligible studies to examine the relationship
167 between kinesiophobia and physical activity. Correlations were pooled using the generic inverse
168 pooling method via the ‘metacor’ function in the {meta} R package⁵⁰. This function automatically
169 performs a necessary Fisher’s z-transformation on the original, untransformed correlations prior
170 to pooling. The ‘metacor’ function also reconverts the pooled association back to its original form
171 for ease of interpretation. Correlation estimates were nested within studies using the ‘cluster’
172 argument to account for the dependencies between these estimates, resulting in a three-level meta-
173 analysis (level 1: participants, level 2: correlation estimates, level 3: studies). The distribution of
174 variance across levels was assessed using the multilevel version of I^2 ⁵⁴. The performance of the 2-
175 level and 3-level meta-analyses was assessed and compared using the {metafor} R package^{51,52}.
176 We anticipated considerable between-study heterogeneity, and therefore used a random-effects
177 model to pool correlations. The restricted maximum likelihood (RML) estimator⁵⁵ was used to
178 calculate the heterogeneity variance τ^2 . In addition to τ^2 , to quantify between-study

179 heterogeneity, we report the I^2 statistic, which provides the percentage of variability in the
180 correlations that is not caused by sampling error⁵⁶. The I^2 statistic was interpreted as follows: 0-
181 40%, may not be important; 30-60%, may represent moderate heterogeneity; 50-90%, may
182 represent substantial heterogeneity; and 75-100%, may represent considerable heterogeneity. To
183 reduce the risk of false positives, we used a Knapp-Hartung adjustment⁵⁷ to calculate the
184 confidence interval around the pooled association. We also report the prediction interval, which
185 provides a range within which we can expect the associations of future studies to fall based on the
186 current evidence. The pooled correlation was interpreted using Cohen's conventions⁵⁸: $r \approx -0.10$,
187 small negative correlation; $r \approx -0.30$, moderate negative correlation; $r \approx -0.50$, large negative
188 correlation.

189

190 *Publication Bias Assessment*

191 Publication bias was assessed using a funnel plot, which is a scatter plot of the studies' effect size
192 expressed as the Fisher's z transformed correlation on the x-axis against a measure of their standard
193 error (which is indicative of precision of the study's effect size) on the y-axis. When there is no
194 publication bias, the data points in a funnel plot should form a roughly symmetrical, upside-down
195 funnel. Studies in the top part of the plot, which have lower standard errors, are expected to lie
196 closely together, and not far away from the pooled effect size. In the lower part of the plot, studies
197 have higher standard errors, the funnel "opens up", and effect sizes are expected to scatter more
198 heavily to the left and right of the pooled effect. Egger's regression⁵⁹ can be used to formally test
199 funnel plot's asymmetry. However, since there is no direct function to conduct Egger's test for 3-
200 level models, we calculated it by using the standard errors of the effect size estimates as a predictor
201 in the meta-regression⁶⁰.

202 P-curve analysis⁶¹ was conducted to assess whether the distribution of the statistically significant
203 results was consistent with what would be expected if only true effects were present. When the
204 null hypothesis is true (i.e., there is no true effect), p-values are assumed to follow a uniform
205 distribution: highly significant effects (e.g., $p = 0.01$) are as likely as barely significant effects
206 (e.g., $p = 0.049$). However, when the null hypothesis is false (i.e., there is a true effect in our data),
207 p-values are assumed to follow a right-skewed distribution: highly significant effects are more
208 likely than barely significant effects. A left-skewed distribution would suggest that some studies
209 used statistical tests to find significant results in ways that may not be reproducible or generalizable
210 (i.e., p-hacking).

211

212 *Secondary Meta-Analysis*

213 A secondary meta-analysis was conducted using the same approach, but based on Spearman's rho
214 values, to further test the relationship between kinesiphobia and physical activity.

215

216 *Subgroup analyses and meta-regressions*

217 Subgroup analyses were conducted to examine the differences in correlations between studies
218 including participants with different health conditions and using different types of physical activity
219 measures (i.e., device-based versus self-reported), physical activity measurement instruments (i.e.,
220 type of questionnaires, type of devices), physical activity outcomes, and kinesiphobia measures.
221 Exploratory meta-regressions were conducted to examine if the average age of participants, the
222 proportion of women, and pain in a study predicted the reported correlation between kinesiphobia
223 and physical activity. Pain was normalized to a 0-100 scale to make the data comparable across

224 pain scales. A sensitivity analysis was conducted to examine whether the quality of the studies
225 affected the results.

226

227

RESULTS

228 Literature Search

229 The primary search identified 3,015 potentially relevant articles from the five databases (Figure
230 1), including 912 duplicates. Of the 2,103 articles screened, disagreement occurred in 210 cases
231 (10%), all of which were resolved by consensus. All articles remained after step 1 as they were all
232 written in English. 1,133 articles were excluded in step 2 because they were irrelevant (n = 710)
233 or did not report original data (n = 423). No articles were excluded in step 3 because they all
234 involved human participants. Eight hundred and fifty-two articles were excluded in step 4 because
235 they did not assess kinesiophobia (n = 117) or physical activity (n = 735). Seventy-seven articles
236 were initially excluded at step 5 because they did not formally test the correlation between
237 kinesiophobia and physical activity or did not report the estimate of this correlation. However, the
238 corresponding authors of these articles were contacted by email to request the Pearson correlation
239 estimate of this association and the sample size used to calculate it. Nineteen authors replied to our
240 email: Eight authors provided raw data for 10 studies⁶²⁻⁷¹ and 11 authors provided the Pearson's
241 correlation estimate^{29,72-81}. In addition, the Pearson's correlation estimate of two articles were
242 calculated based on information reported in the article^{82,83}. This process reduced the number of
243 studies excluded at step 5 to 54, resulting in a total of 64 articles included from the databases.

244 Using reference screening and forward citation tracking, the authors identified 27 studies that
245 assessed both physical activity and kinesiophobia, of which 8 reported an estimate of their
246 relationship⁸⁴⁻⁹¹ and 19 did not⁹²⁻¹¹⁰. The corresponding authors of these 19 studies were asked by

247 email to provide this estimate or their data. Two authors sent the estimate^{104,107}. Seventeen emails
248 remained unanswered^{92-103,105,106,108-110}.

249

250 **Descriptive Results**

251 *Participants*

252 The 74 articles identified by the systematic review included a total of 13388 participants aged 11
253 to 85 years, including 7308 women, 4729 men, and 1351 participants whose gender and sex was
254 not reported. The studies investigated populations with pain (n = 37)^{62,64-71,73,76-81,83,88,89,111-128}
255 cardiac conditions (n = 6)^{86,87,104,129-131}, surgery (n = 8)^{63,131-137}, arthritis (n = 10)^{75,82,84,85,91,138-142},
256 neurologic conditions (n = 3)^{90,143,144}, pulmonary conditions (n = 3)¹⁴⁵⁻¹⁴⁷, cancer (n = 1)⁷⁴, women
257 health conditions (n = 2)^{72,148}, as well as healthy adults (n = 6)^{29,85,107,149-151} (Table 1).

258

259 *Kinesiophobia*

260 In 54 of the 74 studies, kinesiophobia was assessed using the 17-item TSK (TSK-17; n =
261 38)^{29,62,64,65,68,72,73,77,84,85,90,91,111-114,117-120,122-125,130,133-135,138-144,146,148,150} shorter versions of the TSK
262 [(TSK-11¹⁵²; n = 10)^{63,81,88,116,121,122,128,131,136,151}, (TSK-14; n = 1)⁷⁴, (TSK-13¹⁵³; n = 2)^{115,137}, (TSK-
263 7; n = 1)¹⁴⁵], or its adaptation for patients with coronary artery disease (TSK-Heart¹⁵⁴; n = 2)^{86,87}.
264 The TSK is a questionnaire that assesses the belief that movements can lead to (re)injury, pain, or
265 aggravation of an underlying and serious medical condition²⁸. Each item is rated on a Likert scale
266 ranging from 1 (strongly disagree) to 4 (strongly agree). On the TSK-17, a score of 37 is used to
267 distinguish between low (≤ 37) and high (> 37) levels of kinesiophobia²⁷. On the TSK-13, scores
268 inferior to 23 are considered sub-clinical¹⁵⁵. The other measures that were used are the Fear-
269 Avoidance Belief Questionnaire¹⁵⁶ (FABQ; n = 15)^{66,67,69-71,76,78-80,82,83,104,126,127,132}, Kinesiophobia

270 Causes Scale¹⁵⁷ (KCS; n = 2)^{107,149}, the Fear of Activities in Situations scale (FActS; n = 1)¹²⁹, the
271 Brief Fear of Movement Scale for Osteoarthritis¹⁵⁸ (BFMSO; n = 1)⁷⁵, and Breathlessness Beliefs
272 Questionnaire (BBQ; n = 1)¹⁴⁷.

273 Sixty-four studies reported mean levels of kinesiophobia (Table 1). The studies based on the TSK-
274 17 or TSK-Heart (mean range: 17 to 68) reporting the highest levels of kinesiophobia were those
275 involving participants with a cardiovascular condition (41.4 to 49.7), followed by studies testing
276 participants with arthritis (31.8 to 45.27) or chronic pain (30.5 to 44.6). Levels of kinesiophobia
277 were lower in participants with a neurological (36.6 to 41), pulmonary (20.7 to 39.6), women
278 health (36), or surgical condition (32.9 to 35.9), and in healthy adults (18.9 to 39.0).

279

280 *Physical Activity*

281 Fifty-one studies assessed physical activity using a self-reported measure (Table 1). Most of these
282 questionnaire-based studies used the short form of the International Physical Activity
283 Questionnaire (IPAQ-SF; n = 20)^{29,73,76,78,79,82,85,86,112,114,116,130,131,133,138-141,146,147} which consists of
284 6 items assessing time spent in light (i.e., walking), moderate (e.g., carrying light loads, cycling at
285 moderate speed, doubles tennis), and vigorous physical activity (e.g., digging, fast cycling, heavy
286 lifting, aerobics) over the last 7 days¹⁵⁹. Other questionnaires were used to assess physical activity,
287 such as the Baecke Habitual Physical Activity Questionnaire¹⁶⁰ (BHPAQ; n = 5)^{64,89,90,107,113},
288 Saltin-Grimby Physical Activity Level Scale¹⁶¹ (SGPALS; n = 5)^{80,104,115,125,135}, Godin-Shephard
289 Leisure-Time Exercise Questionnaire¹⁶² (GLTEQ; n = 2)^{121,134}, Minnesota Leisure Time Physical
290 Activity Questionnaire¹⁶³ (MLTPAQ; n = 1)¹³⁶, Physical Activity Scale for the Elderly¹⁶⁴ (PASE;
291 n = 2)^{123,149}, Physical Activity Questionnaire for the Elderly¹⁶⁴ (PAQE; n = 1)⁷², Short
292 Questionnaire to Assess Health Enhancing Physical Activity¹⁶⁵ (SQUASH; n = 1)¹⁴², the Tegner

293 Assessment Scale¹⁶⁶ (TAS; n = 1)⁶³, University of California Los Angeles (UCLA) activity score¹⁶⁷
294 (n = 1)⁸⁴, Leisure Time Physical Activity Index¹⁶⁸ (LTPAI; n = 1)⁶⁷, Global Physical Activity
295 Questionnaire¹⁶⁹ (GPAQ; n = 2)^{77,83}, Freiburger Questionnaire on Physical Activity¹⁷⁰ (FQPA; n
296 = 1)¹²⁷, Jurka Physical Activity Scale¹⁷¹ (JPAS; n = 1)⁸¹, Rapid Assessment of Physical Activity
297 questionnaire¹⁷² (RAPAQ; n = 1)⁶⁸, Tecumseh Occupational Activity Questionnaire¹⁷³ (TOAQ; n
298 = 1)¹³⁶, and Australian Health Survey-derived questions (AHS; n = 1)⁷¹.

299 Physical activity was also assessed with devices such as accelerometers measuring accelerations
300 in 3 dimensions (n = 23)^{65,66,74,75,78,91,110,113,117,118,120,122,124,126,129,132,137,143-145,148,150,151} and
301 pedometers measuring the number of steps (n = 3)^{63,88,128} (Table 1). In most studies, the device
302 was worn at the hip (n = 10)^{63,91,113,117,124,129,137,148,150,151}. Other positions included wrist (n =
303 5)^{65,111,126,132,143}, arm (n = 3)^{74,144,145}, trunk (n = 2)^{118,122}, and thigh (n = 1)⁷⁵, with five studies not
304 reporting where the device was worn^{66,78,88,120,128}. Most studies that employed accelerometer-based
305 measures used the ActiGraph (Actigraph, LLC, Pensacola, FL, USA) GT3X+ (n = 4)^{120,137,150,151},
306 wGT3X-BT (n = 2)^{138,148} or GT9X Link (n = 2)^{78,132}. The other accelerometers were the RT3
307 (Stayhealthy Inc., Monrovia, CA, USA; n = 3)^{117,118,124}, the SenseWear Pro3 Armband
308 (BodyMedia, Pittsburgh, PA, USA; n = 3)^{74,144,146}, the Activity Sensory Move II (movisens GmbH,
309 Karlsruhe, Germany; n = 1)¹²⁹, the LifeShirt (Vivometrics, Inc., Ventura, CA, USA; n = 1)¹²², the
310 ActiWatch (Mini Mitter Co., Inc., Bend, OR, USA; n = 1)¹¹¹, AX3 (Axtivity, Newcastle upon
311 Tyne, UK; n = 1)⁶⁵, FitBit (FitBit Inc., San Francisco, CA) Charge HR (n = 1)¹²⁶, Charge 3 (n =
312 1)¹⁴³, and the Activ8 (2M Engineering, North Brabant, Netherland, n = 1)⁶⁶. The type of
313 accelerometer was not reported in one study¹¹³. The pedometers were the Digi-Walker SW-200
314 (New Lifestyles Inc., Lees Summit, MO, USA; n = 1)⁶³, the Active Style Pro HJA-350IT (Omron
315 Healthcare, Kyoto, Japan; n = 1)⁸⁸ and Yamax Power-Walker EX-510 3D (Pedometer Express,

316 Minnesota, USA; $n = 1$)¹²⁸. These devices were worn for 5 days ($n = 1$)¹¹¹, 6 days ($n = 1$)¹²⁹, 7 days
317 ($n = 16$)^{63,65,75,88,91,113,117,118,120,124,132,137,144,148,150,151}, or 14 days ($n = 1$)¹²⁶. The remaining 7 studies
318 did not specify the number of days the device was worn^{66,74,78,122,128,143,145}. All studies provided the
319 accelerometer or pedometer on the day kinesiophobia was assessed ($n =$
320 18)^{63,65,66,75,78,88,89,111,113,120,122,126,132,137,143,148,150}. The remaining studies did not specify whether
321 kinesiophobia was measured the day the device was provide or the last day of physical activity
322 assessment ($n = 7$)^{74,118,124,128,144,146,151}.

323 To assess physical activity, the studies used the following outcomes: Score from a questionnaire
324 (e.g., TAS, PAQE, BHPAQ, SGPALS, LTPAQ, $n =$
325 24)^{63,64,67,68,72,80,81,84,87,89,90,104,107,113,115,119,121,123,125,127,134,137,149}, MET-min/week ($n = 23$)^{31,73,76,77-}
326 79,83,85,86,112,114,116,130,131,133,136,138-142,146,147, steps per day ($n = 14$)^{63,65,75,78,88,113,120,126,128,132,143-145,150},
327 hours per day or week ($n = 12$)^{62,65,69-71,74,91,113,120,138,143,145}, counts per minute ($n = 4$)^{111,113,117,137},
328 kilocalories per day ($n = 2$)^{129,144}, or percentage of active time ($n = 1$)¹²². Nine studies used multiple
329 physical activity outcomes^{63,65,78,91,113,120,143-145}.

330

331 *Association Between Physical Activity and Kinesiophobia*

332 Among the 74 articles included in the systematic review, 42 reported correlation coefficients of
333 the association between physical activity and kinesiophobia. Specifically, 32 articles reported at
334 least one Pearson's r correlation coefficient and 12 articles reported at least one Spearman's
335 rho^{87,89,91,113,116,124,127,132,135,143,149,151}. When a correlation coefficient was not reported, but the exact
336 p-value (or t value) and sample size were available and it was possible to know the sign of the
337 correlation, which was the case for 7 studies^{83,111,115,120,125,133,144}, the Pearson's r estimate was
338 computed using an ad-hoc R code (Supplemental Code 1A). For the studies that reported a relative

339 p-value < 0.001 instead of an exact p-value, we used a p-value of 0.0009 to estimate an
340 approximate r value⁸².

341 Through email correspondence with the authors, we obtained 23 additional Pearson's r
342 estimates^{29,62-81,104,107}. In total, 83 Pearson's r estimates from 63 studies and 21 Spearman's rho
343 estimates from 12 studies were used in the meta-analysis (Table 1). The remaining study did not
344 report a correlation coefficient and was therefore not included in the meta-analysis¹¹⁷. This study
345 reported a non-statistically significant positive association between physical activity and
346 kinesiophobia based on a standardized beta coefficient.

347

348 *Pain*

349 Mean pain intensity at rest was reported in 45 out of the 74 articles included in the systematic
350 review. Most studies used the Visual Analog Scale¹⁷⁴ (VAS; n = 21)<sup>65,69-
351 71,76,78,82,85,89,90,112,114,117,119,120,122,124,126,138,139,141</sup> or the Numeric Rating Scale¹⁷⁵ (NRS; n =
352 15)^{29,62,66,68,75,79,80,83,88,111,113,116,127,128,137}. Other studies used the Knee Injury Osteoarthritis
353 Outcome Score pain subscale¹⁷⁶ (KOOS-P; n = 3)^{63,84,134}, Brief Pain Inventory¹⁷⁷ (n = 1)⁶⁴, Oxford
354 Knee Score¹⁷⁸ (OKS; n = 1)¹⁴⁰, the Quality of Well-Being Scale – Self-administered Pain Scale¹⁷⁹
355 (QWBS-P; n = 1)¹⁵⁰, the Short Form 36 bodily pain¹⁸⁰ (SF-36; n = 1)¹³⁰, the Graphic Rating
356 Scale¹⁸¹ (GRS; n = 1)¹³³, Fibromyalgia Impact Questionnaire-Pain¹⁸² (FIQ-Pain, n = 1)⁶⁷ and the
357 Verbal Rating Scale¹⁸³ (VRS; n = 1)¹⁰⁴. In the meta-analysis, scores that were not on a 0-100 scale
358 in the initial measure were scaled to that range.

359

360 **Meta-Analysis**

361 *Main Meta-Analysis*

362 Our main meta-analysis of 63 studies, 83 Pearson's r correlation estimates, and 12278 participants
363 revealed a statistically significant small-to-moderate negative correlation between kinesiophobia
364 and physical activity ($r = -0.19$; 95% confidence interval [95CI]: -0.26 to -0.13 ; $p < 0.0001$) (Table
365 2; Figure 2). However, we observed substantial-to-considerable between-study statistical
366 heterogeneity ($\text{Tau}^2 = 0.06$, 95CI: 0.02 to 0.09 ; $I^2 = 85.5\%$, 95CI: 82.6 to 87.9%), and the
367 prediction interval ranged from $r = -0.61$ to 0.30 , indicating that a moderate positive correlation
368 cannot be ruled out for future studies.

369 The sampling error variance on level 1 and the value of I^2 on level 2, i.e., the amount of
370 heterogeneity variance within studies, were small (10.3% and 8.2% , respectively). The largest
371 share of heterogeneity variance was from level 3, with between-study heterogeneity making up
372 81.5% of the total variation in our data (Supplemental Figure 1). Overall, this indicates that there
373 is considerable between-study heterogeneity, and less than one tenth of the variance can be
374 explained by differences within studies.

375 The 3-level model showed a better fit than the 2-level model with lower Akaike's information
376 criterion (AIC) (28.4 vs. 39.0) and Bayesian information criterion (BIC) (35.6 vs. 43.8), indicating
377 better performance. These lower AIC and BIC are consistent with the significant likelihood ratio
378 test (LRT) comparing the two models ($\chi^2 = 12.67$, $p = 0.0004$). Therefore, although the 3-level
379 model introduces an additional parameter, this added complexity has improved our estimate of the
380 pooled effect.

381

382 *Publication bias assessment*

383 Egger's regression test using the standard errors of the effect size estimates as a predictor in the
384 meta-regression showed that the coefficient of the standard error was significant ($b = -1.497$, 95CI:

385 -2.618 to -0.3754, $p = 0.0095$), suggesting that the data in the funnel plot was asymmetrical (Figure
386 3A). This asymmetry may be explained by publication bias, but also by other potential causes,
387 such as different study procedures and between-study heterogeneity¹⁸⁴, which was substantial-to-
388 considerable here.

389 The 83 Pearson's r correlation estimates were provided to the p-curve analysis. The observed p-
390 curve included 35 statistically significant results ($p < 0.05$), 27 of which were highly significant
391 ($p < 0.025$), and was visually right-skewed (Figure 3B). The other results were excluded because
392 they had a $p > 0.05$. The p-value of the right-skewness test was < 0.001 for both the half curve
393 (curve of p-values ≤ 0.025) and the full curve (curve of p-values < 0.05), confirming that the p-
394 curve was right-skewed and suggesting that the effect of our meta-analysis is true, i.e., that the
395 effect we estimated is not an artifact caused by selective reporting (e.g., p-hacking) in the
396 literature¹⁸⁵. In addition, the statistical power of the studies that were included in the p-curve
397 analysis was 97% (90CI: 93 to 98%), suggesting that approximately 90% of the significant
398 results are expected to be replicable.

399

400 *Secondary Meta-Analyses*

401 Results of the secondary meta-analysis of 12 studies, 21 Spearman's rho correlation estimates, and
402 2084 participants was consistent with the main meta-analysis as it showed a statistically significant
403 small-to-moderate negative correlation between kinesophobia and physical activity ($r = -0.20$;
404 95CI: -0.38 to -0.01; $p = 0.049$) (Table 2; Supplemental Figure 2). However, we observed
405 substantial-to-considerable between-study statistical heterogeneity ($\text{Tau}^2 = 0.10$, 95CI: 0.04 to
406 0.28; $I^2 = 86.3\%$) and the prediction interval ranged from $r = -0.71$ to 0.45, indicating that a
407 moderate positive correlation cannot be ruled out for future studies.

408

409 *Subgroup Meta-Analyses*

410 The test of subgroup differences between health status was conducted on studies comprising
411 people with chronic (k = 35) or acute pain (k = 2), arthritis (k = 11), a cardiovascular condition (k
412 = 10), a neurological condition (k = 8), surgery (k = 5), older age (k = 3), obstructive sleep apnea
413 (k = 2), a pulmonary condition (k = 2), fibromyalgia (k = 2), cancer (k = 1), as well as in post-
414 partum women (k = 1) and healthy young adults (k = 1) (Table 2; Supplemental Figure 3). We
415 found a statistical moderating effect of health status (p = 0.0014). The relationship between
416 kinesiophobia and physical activity was statistically significant only in studies that included
417 participants with cardiac condition (r = -0.30; 95CI: -0.47 to -0.11), arthritis (r = -0.25; 95CI: -0.39
418 to -0.10), a neurologic condition (r = -0.53; 95CI: -0.69 to -0.32), a pulmonary condition (r = -
419 0.68; 95CI: -0.82 to -0.46), or older adults (r = -0.40; 95CI: -0.60 to -0.14). We found no evidence
420 of an association between kinesiophobia and physical activity in studies that included participants
421 with chronic pain (r = -0.07; 95CI: -0.16 to 0.01) or acute pain (r = -0.13; 95CI: -0.45 to 0.23).
422 Statistical heterogeneity was higher in the studies comprising people with a pulmonary condition
423 (I² = 98.1%), arthritis (I² = 93.4%), or older adults (I² = 91.2%) than in the studies comprising
424 people with a cardiac (I² = 28.7%) or neurologic condition (I² = 55.9%).

425 The test of subgroup differences between self-reported (k = 54) and device-based (k = 29)
426 measures of physical activity showed no evidence of a moderating effect of the type of physical
427 activity measure (p = 0.171; Table 2). Both self-reported measures (r = -0.22; 95CI: -0.29 to -0.14;
428 I² = 89.3%) and device-based measures (r = -0.13; 95CI: -0.24 to -0.02; I² = 57.2%) (Supplemental
429 Figure 4) showed a negative association between kinesiophobia and physical activity.

430 We also found no evidence of a moderating effect of physical activity instruments ($p = 0.209$)
431 (Supplemental Figure 5), physical activity outcome ($p = 0.685$) (Supplemental Figure 6), or
432 kinesiophobia instrument ($p = 0.452$) (Supplemental Figure 7).

433

434 *Meta-Regressions*

435 Age did not statistically influence the correlation estimates of the meta-analysis studies ($k = 72$; p
436 $= 0.349$). Similarly, the proportion of women ($k = 72$; $p = 0.555$) and the mean level of pain in the
437 studies ($k = 49$; $p = 0.481$) did not influence correlation estimates.

438

439 *Sensitivity Analysis*

440 The meta-regression by quality score showed that a study's quality did not influence correlation
441 estimates ($k = 83$; $p = 0.373$).

442

443

DISCUSSION

444 The main objective of this study was to systematically review and meta-analyze the direct
445 relationship between kinesiophobia and physical activity. In addition, we examined the influence
446 of potential moderators, such as health status. To our knowledge, this is the first review of its kind
447 on this research topic.

448

449 **Kinesiophobia and Physical Activity**

450 Both the main meta-analysis based on Pearson's r correlation estimates and the secondary meta-
451 analysis based on Spearman's ρ correlation estimates showed a small-to-moderate negative
452 correlation between kinesiophobia and physical activity. Importantly, this correlation was

453 observed both in studies using both self-report (e.g., IPAQ) and device-based measures (i.e.,
454 accelerometers or pedometers). These results are consistent with our hypothesis and the dual
455 models of physical activity³²⁻³⁵. According to these theoretical models, our findings suggest that
456 the fear of movement characteristic of kinesiophobia triggers an impulse to avoid physical activity
457 behaviors, which contributes to the maintenance or exacerbation of the initial fear. Accordingly,
458 kinesiophobia and physical inactivity would be self-perpetuating or even self-reinforcing.

459

460 **Health Status**

461 Our results suggest that patients with a cardiac, neurologic, arthritic, or pulmonary condition, as
462 well as older adults, may be at greater risk for this negative relationship between kinesiophobia
463 and physical activity than those with other conditions. In individuals with a cardiac condition,
464 kinesiophobia and its impact on physical activity may be explained by a fear of inducing a new
465 cardiac event¹⁸⁶ or “causing more damage to the heart”¹⁸⁷, but also by breathlessness (i.e.,
466 dyspnea), which reduces the ability to be physically active and damages confidence, leading to
467 persistent anticipation of negative outcomes from physical activity¹⁸⁸. Dyspnea is also a major
468 barrier to physical activity in people with a pulmonary condition, such as chronic obstructive
469 pulmonary disease (COPD)¹⁴⁷. Patients with asthma may have additional disease-related barriers
470 to physical activity, such as the fear of provoking respiratory symptoms and exacerbations¹⁸⁹.
471 Regarding neurologic conditions, chest tightness reported in patients with Parkinson’s disease as
472 a barrier to exercise may be a factor contributing to the association between kinesiophobia and
473 physical activity¹⁹⁰. Another potential factor in these patients¹⁹⁰, as well as in stroke survivors¹⁹¹
474 and healthy older adults¹⁹², is fear of falling. In patients with osteoarthritis, the belief that physical

475 activity will “damage the joints”¹⁹³ and the perceived fragility of their physical status¹⁹⁴ may be
476 factors contributing to the relationship between kinesiophobia and physical activity.

477 Although our results showed no evidence of an association between kinesiophobia and physical
478 activity in other health conditions such as cancer, post-surgery, post-partum, or obstructive sleep
479 apnea, these effects cannot be fully ruled out, as the lack of statistical significance could be
480 explained by a lack of statistical power in these subgroup meta-analyses including fewer estimates
481 (k = 1 to 5).

482

483 **Pain versus Fear**

484 Our results showed no evidence of an association between kinesiophobia and physical activity in
485 people with fibromyalgia, acute pain, or chronic pain. This finding was surprising because fear of
486 pain is a key component of kinesiophobia, appearing in 10 of the 17 items on the TSK-17 and
487 TSK-Heart scales, and reinforces the importance of considering the multidimensional nature of
488 kinesiophobia, which not only relates to pain but also reflects fear of injury and fear of worsening
489 a health condition.

490 In addition, contrary to our expectations, we found no statistical evidence showing that pain
491 intensity at rest influenced the effect of kinesiophobia on physical activity, despite the substantial
492 number of estimates included in this analysis (k = 23). This result is consistent with the weak
493 relationship that has been shown between kinesiophobia and pain¹⁹⁵, further suggesting that it is
494 not the actual pain that prevents physical activity, but the fear of triggering pain, injury, or
495 aggravating an underlying condition. However, this absence of evidence might be related to the
496 methods used to assess pain, which may be better assessed by pain history (e.g., pain duration in
497 months) or pain intensity during exercise.

498

499 **Limitations**

500 The results of this systematic review and meta-analysis should be interpreted with consideration
501 of several limitations. (1) We report considerable heterogeneity across the included studies, which
502 may be explained by the diversity of the methods used to assess physical activity (questionnaires
503 vs. accelerometers vs. pedometers), the instruments used in these methods (14 different
504 questionnaires, 14 different accelerometers and pedometers), and the physical activity outcomes
505 ($n = 6$), but also by the different questionnaires used to assess kinesiophobia ($n = 11$). This
506 heterogeneity suggests that the measures of kinesiophobia and physical activity used in the
507 literature reflect different dimensions of these two constructs. For example, self-reported measures
508 of physical activity do not accurately reflect actual levels of physical activity¹⁹⁶. (2) Because
509 kinesiophobia is a state, i.e., a dynamic psychological variable, the time difference between the
510 physical activity and kinesiophobia assessments, as well as the context of assessment, may have
511 influenced the results. (3) While a subgroup meta-analysis showed no evidence of an effect of the
512 type of TSK scale, inconsistencies have been noted in the purported dimensions assessed by
513 different TSK scales or across populations¹⁹⁷, which may have influenced our results. (4) The
514 meta-analysis was not pre-registered because the author with this expertise (IML) joined only after
515 the systematic review was conducted. However, since the pre-registered systematic review defines
516 the studies included in the meta-analysis and since the {meta} R package⁵⁰ provides very few
517 degrees of freedom, there was little room for bias in the main and secondary meta-analyses. This
518 reasoning also applies for the subgroup analyses as the health conditions, types of physical activity
519 measures, measurement instruments, and outcomes, as well as the kinesiophobia measurement
520 instruments were imposed by the studies that were included in the systematic review. (5) Only 21

521 of the 98 authors we contacted (21%) shared their estimates (n = 13) or raw data (n = 8) with us,
522 which is more than reported in previous literature¹⁹⁸. Including these missing data may have had
523 an impact on the results.

524

525 **Conclusions**

526 Higher levels of kinesiophobia were associated with lower levels of physical activity, especially
527 in people with a cardiac, neurologic, arthritic, and pulmonary condition. According to theoretical
528 models, this relationship between kinesiophobia and physical activity results from automatic
529 processes that may be self-reinforcing and should therefore not be overlooked. However,
530 heterogeneity between studies was substantial-to-considerable for some results, and the evidence
531 for publication bias calls for cautious conclusions about this potential relationship. In sum, our
532 results suggest that kinesiophobic individuals have a small-to-moderate risk of being more
533 physically inactive than other individuals. More evidence is required to determine the impact
534 kinesiophobia should have on therapeutic decisions when aiming to maintain or increase physical
535 activity. Particularly, prospective studies are needed to better understand the factors and
536 mechanisms that influence the relationship between kinesiophobia and physical activity.

537

5. ARTICLE INFORMATION

538 **Funding**

539 Matthieu P. Boisgontier is supported by the Natural Sciences and Engineering Research Council
540 of Canada (NSERC; RGPIN-2021-03153), the Canada Foundation for Innovation (CFI), Mitacs,
541 and the Banting Research Foundation. The funders had no role in the data collection, management,
542 analysis and interpretation, writing of the report, or the decision to submit the report for
543 publication. Ata Farajzadeh is supported by an Admission Scholarship, a Doctoral International
544 Scholarship, and a Special Merit Scholarship from the University of Ottawa.

545

546 **Data and Code Sharing**

547 According to good research practices⁴⁴, the dataset, R Markdown script, and supplemental material
548 are freely available in Zenodo: <https://doi.org/10.5281/zenodo.8216112125>¹⁹⁹

549

550 **Acknowledgements**

551 Based on the Contributor Roles Taxonomy (CRediT)²⁰⁰ individual author contributions to this
552 work are as follows: Miriam Goubran: Conceptualization, Methodology (Systematic Review),
553 Investigation, Data Curation, Writing (Original Draft), Writing (Review and Editing); Ata
554 Farajzadeh: Methodology (Systematic Review), Investigation, Data Curation, Writing (Original
555 Draft), Writing (Review and Editing); Ian M. Lahart: Methodology (Meta-Analysis), Formal
556 Analysis, Visualization, Data Curation, Writing (Original Draft), Writing (Review and Editing);
557 Martin Bilodeau: Conceptualization, Methodology (Systematic Review), Writing (Review and
558 Editing), Supervision (MG); Matthieu P. Boisgontier: Conceptualization, Methodology
559 (Systematic Review and Meta Analysis), Investigation, Formal Analysis, Data Curation,

560 Visualization; Writing (Original Draft), Writing (Review and Editing), Supervision (MG and AF),
561 Project Administration, Funding Acquisition.

562

563 **Conflict of Interest Disclosure**

564 The authors declare they have no conflict of interest relating to the content of this article

REFERENCES

1. Morris JN, Heady JA, Raffle PA, Roberts CG, Parks JW. Coronary heart-disease and physical activity of work. *Lancet*. 1953;262(6795):1053-1057.
[https://doi.org/10.1016/s0140-6736\(53\)90665-5](https://doi.org/10.1016/s0140-6736(53)90665-5)
2. Lee IM, Shiroma EJ, Lobelo F, et al. Effect of physical inactivity on major non-communicable diseases worldwide: an analysis of burden of disease and life expectancy. *Lancet*. 2012;380(9838):219-229.
[https://doi.org/10.1016/S0140-6736\(12\)61031-9](https://doi.org/10.1016/S0140-6736(12)61031-9)
3. Salinas-Rodríguez A, Manrique-Espinoza B, Palazuelos-González R, Rivera-Almaraz A, Jáuregui A. Physical activity and sedentary behavior trajectories and their associations with quality of life, disability, and all-cause mortality. *Eur Rev Aging Phys Act*. 2022;19(1):13.
<https://doi.org/10.1186/s11556-022-00291-3>
4. Wahid A, Manek N, Nichols M, et al. Quantifying the association between physical activity and cardiovascular disease and diabetes: a systematic review and meta-analysis. *J Am Heart Assoc*. 2016;5(9):e002495. <https://doi.org/10.1161/JAHA.115.002495>
5. Bleich SN, Vercammen KA, Zatz LY. Interventions to prevent global childhood overweight and obesity: a systematic review. *Lancet Diabetes Endocrinol*. 2018;6:332-346.
[https://doi.org/10.1016/S2213-8587\(17\)30358-3](https://doi.org/10.1016/S2213-8587(17)30358-3)
6. Schuch F, Vancampfort D, Firth J, et al. Physical activity and sedentary behavior in people with major depressive disorder: a systematic review and meta-analysis. *J Affect Disord*. 2017;210:139-150. <https://doi.org/10.1016/j.jad.2016.10.050>

7. Liu X, Zhang D, Liu Y, et al. Dose-response association between physical activity and incident hypertension: a systematic review and meta-analysis of cohort studies. *Hypertension*. 2017;69(5):813-820.
<https://doi.org/10.1161/HYPERTENSIONAHA.116.08994>
8. Moore SC, Lee IM, Weiderpass E, et al. Association of leisure-time physical activity with risk of 26 types of cancer in 1.44 million adults. *JAMA Intern Med*. 2016;176(6):816-825.
<https://doi.org/10.1001/jamainternmed.2016.1548>
9. Najjar J, Östling S, Gudmundsson P, et al. Cognitive and physical activity and dementia: a 44-year longitudinal population study of women. *Neurology*. 2019;92(12):E1322-E1330.
<https://doi.org/10.1212/WNL.0000000000007021>
10. World Health Organization. *Physical activity - Key facts*. October 2022. Accessed May 1, 2024. <https://www.who.int/news-room/fact-sheets/detail/physical-activity>
11. Taylor RS, Walker S, Smart NA, et al. Impact of exercise rehabilitation on exercise capacity and quality-of-life in heart failure: individual participant meta-analysis. *J Am Coll Cardiol*. 2019;73(12):1430-1443. <https://doi:10.1016/j.jacc.2018.12.072>
12. Molloy C, Long L, Mordi IR, et al. Exercise-based cardiac rehabilitation for adults with heart failure. *Cochrane Database Syst Rev*. 2024;3(3):CD003331.
<https://doi:10.1002/14651858.CD003331.pub6>
13. Daste C, Kirren Q, Akoum J, Lefèvre-Colau MM, Rannou F, Nguyen C. Physical activity for osteoarthritis: Efficiency and review of recommendations. *Joint Bone Spine*. 2021;88(6):105207. <http://doi:10.1016/j.jbspin.2021.105207>

14. van Allen Z, Orsholits D, Boisgontier MP. Pre-stroke physical activity matters for functional limitations: A longitudinal case-control study of 12,860 participants. *Phys Ther.* 2024. <http://doi.org/10.1101/2023.09.14.23295576>
15. Turan TN, Nizam A, Lynn MJ, et al. Relationship between risk factor control and vascular events in the SAMMPRIS trial. *Neurology.* 2017;88(4):379-385. <http://doi:10.1212/WNL.0000000000003534>
16. Brenner H, Chen C. The colorectal cancer epidemic: challenges and opportunities for primary, secondary and tertiary prevention. *Br J Cancer.* 2018;119(7):785-792. <http://doi:10.1038/s41416-018-0264-x>
17. Boisgontier MP, Iversen MD. Physical inactivity: a behavioral disorder in the physical therapist's scope of practice. *Phys Ther.* 2020;100(5):743-746. <http://doi.org/10.1093/PTJ/PZAA011>
18. O'Donoghue G, Perchoux C, Mensah K, et al. A systematic review of correlates of sedentary behaviour in adults aged 18-65 years: a socio-ecological approach. *BMC Public Health.* 2016;16(1):163. <https://doi.org/10.1186/s12889-016-2841-3>
19. Garcia L, Mendonça G, Benedetti TRB, et al. Barriers and facilitators of domain-specific physical activity: a systematic review of reviews. *BMC Public Health.* 2022;22(1):1964. <https://doi.org/10.1186/s12889-022-14385-1>
20. Biddle SJH, Gorely T, Faulkner G, Mutrie N. Psychology of physical activity: a 30-year reflection on correlates, barriers, and theory. *Int J Sport Exerc Psychol.* 2023;21(1):1-14. <https://doi.org/10.1080/1612197X.2022.2147261>

21. Azevedo MR, Araújo CLP, Reichert FF, Siqueira FV, da Silva MC, Hallal PC. Gender differences in leisure-time physical activity. *Int J Public Health*. 2007;52(1):8-15.
<https://doi.org/10.1007/s00038-006-5062-1>
22. Caspersen CJ, Pereira MA, Curran KM. Changes in physical activity patterns in the United States, by sex and cross-sectional age. *Med Sci Sports Exerc*. 2000;32(9):1601-1609.
<https://doi.org/10.1097/00005768-200009000-00013>
23. Cheval B, Rebar AL, Miller MW, et al. Cognitive resources moderate the adverse impact of poor perceived neighborhood conditions on self-reported physical activity of older adults. *Prev Med*. 2019;126:105741. <https://doi.org/10.1016/j.ypmed.2019.05.029>
24. Cheval B, Orsholits D, Sieber S, Courvoisier D, Cullati S, Boisgontier MP. Relationship between decline in cognitive resources and physical activity. *Health Psychol*. 2020;39(6):519-528. <https://doi.org/10.1037/hea0000857>
25. Cheval B, Sieber S, Guessous I, et al. Effect of early- and adult-life socioeconomic circumstances on physical inactivity. *Med Sci Sports Exerc*. 2018;50(3):476-485.
<https://doi.org/10.1249/MSS.0000000000001472>
26. Kori SH, Miller RP, Todd DD. Kinesiophobia: a new view of chronic pain behavior. *Pain Manag*. 1990;3:35-43.
27. Vlaeyen JWS, Kole-Snijders AMJ, Boeren RGB, van Eek H. Fear of movement/(re)injury in chronic low back pain and its relation to behavioral performance. *Pain*. 1995;62(3):363-372. [https://doi.org/10.1016/0304-3959\(94\)00279-N](https://doi.org/10.1016/0304-3959(94)00279-N)
28. Miller R, Kori S, Todd D. The Tampa Scale: a measure of kinesiophobia. *Clin J Pain*. 1991;7(1):51-52.

29. Bahar Özdemir Y. Investigation of low back pain in the white-collar population working from home due to the COVID-19 pandemic. *J Phys Med Rehabil Sci*. 2021;24(2):135-142. <https://doi.org/10.31609/jpmrs.2021-81527>
30. Stults-Kolehmainen MA, Blacutt M, Bartholomew JB, et al. Motivation states for physical activity and sedentary behavior: desire, urge, wanting, and craving. *Front Psychol*. 2020;11:568390. <https://doi.org/10.3389/fpsyg.2020.568390>
31. Stevens CJ, Baldwin AS, Bryan AD, Conner M, Rhodes RE, Williams DM. Affective determinants of physical activity: a conceptual framework and narrative review. *Front Psychol*. 2020;11:568331. <https://doi.org/10.3389/fpsyg.2020.568331>
32. Brand R, Ekkekakis P. Affective–reflective theory of physical inactivity and exercise. *Ger J Exerc Sport Res*. 2018;48:48-58. <https://doi.org/10.1007/s12662-017-0477-9>
33. Cheval B, Boisgontier MP. Promouvoir une activité physique régulière chez les patients : l'importance de la perception de l'effort. *STAPS*. 2024. <https://doi.org/10.3917/sta.pr1.0091>
34. Cheval B, Boisgontier MP. The theory of effort minimization in physical activity. *Exerc Sport Sci Rev*. 2021;49(3):168-178. <https://doi.org/10.1249/JES.0000000000000252>
35. Conroy DE, Berry TR. Automatic affective evaluations of physical activity. *Exerc Sport Sci Rev*. 2017;45(4):230-237. <http://doi.org/10.1249/JES.0000000000000120>
36. Farajzadeh A, Goubran M, Beehler A, et al. Relationship between age and physical and sedentary stimuli. *Peer Community J*. 2023;3:e21. <https://doi.org/10.24072/pcjournal.246>
37. Hanel J, Owen PJ, Held S, et al. Effects of exercise training on fear-avoidance in pain and pain-free populations: systematic review and meta-analysis. *Sports Med*. 2020;50(1):2193-2207. <https://doi.org/10.1007/s40279-020-01345-1>

38. Jadhakhan F, Sobeih R, Falla D. Effects of exercise/physical activity on fear of movement in people with spine-related pain: a systematic review. *Front Psychol.* 2023;14:1213199.
<https://doi.org/10.3389/fpsyg.2023.1213199>
39. Martinez-Calderon J, Flores-Cortes M, Morales-Asencio JM, Luque-Suarez A. Conservative interventions reduce fear in individuals with chronic low back pain: a systematic review. *Arch Phys Med.* 2020;101(2):329-358.
<https://doi.org/10.1016/j.apmr.2019.08.470>
40. Rovner GS, Sunnerhagen KS, Björkdahl A, et al. Chronic pain and sex-differences; women accept and move, while men feel blue. *PLOS One.* 2017;12(4):e0175737.
<https://doi.org/10.1371/journal.pone.0175737>
41. John JN, Ugwu EC, Okezue OC, et al. Kinesiophobia and associated factors among patients with chronic non-specific low back pain. *Disabil Rehabil.* 2023;45(16):2651-2659.
<https://doi.org/10.1080/09638288.2022.2103747>
42. Page MJ, Moher D, Bossuyt PM, et al. PRISMA 2020 explanation and elaboration: updated guidance and exemplars for reporting systematic reviews. *BMJ.* 2021;372:n160.
<https://doi.org/10.1136/bmj.n160>
43. Caldwell AR, Vigotsky AD, Tenan MS, et al. Moving sport and exercise science forward: a call for the adoption of more transparent research practices. *Sports Med.* 2020;50(3):449-459. <https://doi.org/10.1007/s40279-019-01227-1>
44. Boisgontier MP. Research integrity requires to be aware of good and questionable research practices. *Eur Rehabil J.* 2021;2(1):1-3. <https://doi.org/10.52057/erj.v2i1.24>

45. Goubran M, Boisgontier MP, Bilodeau M. Kinesiophobia and physical activity: a systematic review. *PROSPERO* (CRD42022364063). October 17, 2022. Accessed May 1, 2024. https://www.crd.york.ac.uk/prospero/display_record.php?ID=CRD42022364063
46. National Institutes of Health. Quality assessment tool for observational cohort and cross-sectional studies. Bethesda, MD: National Institutes of Health; 2013. Accessed May 1, 2024. <https://www.nhlbi.nih.gov/health-topics/study-quality-assessment-tools>
47. Des Jarlais DC, Lyles C, Crepaz N; TREND Group. Improving the reporting quality of nonrandomized evaluations of behavioral and public health interventions: the TREND statement. *Am J Public Health*. 2004;94(3):361-366. <http://doi:10.2105/ajph.94.3.361>
48. Schulz KF, Altman DG, Moher D. CONSORT 2010 statement: updated guidelines for reporting parallel group randomised trials. *J Pharmacol Pharmacother*. 2010;1(2):100-107. <http://doi:10.4103/0976-500X.72352>
49. R Core Team. R: *a language and environment for statistical computing* [Computer software]. Version 4.3.1. Vienna, Austria: Foundation for Statistical Computing; 2023. <https://www.r-project.org>
50. Schwarzer G. *meta: general package for meta-analysis* [R package]. Version 6.5-0; 2023. <https://cran.r-project.org/web/packages/meta/meta.pdf>
51. Viechtbauer W. *metafor: meta-analysis package for R* [R package]. Version 4.2-0; 2023. <https://cran.r-project.org/web/packages/metafor/metafor.pdf>
52. Viechtbauer W. Conducting meta-analyses in R with the metafor package. *J Stat Softw*. 2010;36(3):1-48. <https://doi.org/10.18637/jss.v036.i03>

53. Harrer M, Cuijpers P, Furukawa TA, Ebert DD. *Doing Meta-Analysis with R: A Hands-On Guide*. Boca Raton, FL and London: Chapman & Hall/CRC Press. ISBN. 2021;978-0-367-61007. <https://doi.org/10.1201/9781003107347>
54. Cheung M. Modeling dependent effect sizes with three-level meta-analyses: a structural equation modeling approach. *Psychol Methods*. 2014;19(2):211-229. <https://doi.org/10.1037/a0032968>
55. Viechtbauer W. Bias and efficiency of meta-analytic variance estimators in the random-effects model. *J Educ Behav Stat*. 2005;30(3):261-293. <https://doi.org/10.3102/10769986030003261>
56. Higgins JPT, Thompson SG. Quantifying heterogeneity in a meta-analysis. *Stat Med*. 2002;21(11):1539-1558. <https://doi.org/10.1002/sim.1186>
57. Knapp G, Hartung J. Improved tests for a random effects meta-regression with a single covariate. *Stat Med*. 2003;22(17):2693-2710. <https://doi.org/10.1002/sim.1482>
58. Cohen J. *Statistical power analysis for the behavioral sciences*. 2nd ed.; 1988.
59. Egger M, Davey Smith G, Schneider M, Minder C. Bias in meta-analysis detected by a simple, graphical test. *BMJ*. 1997;315(7109):629-634. <https://doi.org/10.1136/bmj.315.7109.629>
60. Marey A. Multilevel meta analysis. RPubS, 2021. Accessed May 1, 2024. <https://rpubs.com/amany33/814435>
61. Simonsohn U, Nelson LD, Simmons JP. P-curve: A key to the file-drawer. *J Exp Psychol Gen*. 2014;143(2):534-547. <http://doi.org/10.1037/a0033242>

62. Assadourian M, Bailly F, Letellier P, et al. Criteria for inclusion in programs of functional restoration for chronic low back pain: pragmatic study. *Ann Phys Rehabil Med*. 2020;63(3):189-194. <https://doi.org/10.1016/j.rehab.2019.06.019>
63. Baez SE, Hoch MC, Hoch JM. Psychological factors are associated with return to pre-injury levels of sport and physical activity after ACL reconstruction. *Knee Surg Sports Traumatol Arthrosc*. 2020;28(2):495-501. <https://doi.org/10.1007/s00167-019-05696-9>
64. Luthi F, Vuistiner P, Favre C, Hilfiker R, Léger B. Avoidance, pacing, or persistence in multidisciplinary functional rehabilitation for chronic musculoskeletal pain: an observational study with cross-sectional and longitudinal analyses. *PLOS One*. 2018;13(9):e0203329. <https://doi.org/10.1371/journal.pone.0203329>
65. Alzahrani H, Mackey M, Stamatakis E, Shirley D. Wearables-based walking program in addition to usual physiotherapy care for the management of patients with low back pain at medium or high risk of chronicity: a pilot randomized controlled trial. *PLOS One*. 2021;16(8):e0256459. <https://doi.org/10.1371/journal.pone.0256459>
66. Koppelaar T, van Dongen JM, Kloek CJ, et al. Effectiveness and cost-effectiveness of a stratified blended physiotherapy intervention compared with face-to-face physiotherapy in patients with nonspecific low back pain: cluster randomized controlled trial. *J Med Internet Res*. 2023;25:e43034. <https://doi.org/10.2196/43034>
67. Palstam A, Larsson A, Bjersing J, et al. Perceived exertion at work in women with fibromyalgia: explanatory factors and comparison with healthy women. *J Rehabil Med*. 2014;46(8):773-780. <https://doi.org/10.2340/16501977-1843>
68. Zadro JR, Shirley D, Nilsen TI, Mork PJ, Ferreira PH. Family history influences the effectiveness of home exercise in older people with chronic low back pain: a secondary

- analysis of a randomized controlled trial. *Arch Phys Med Rehabil*. 2020;101(8):1322-1331.
<https://doi.org/10.1016/j.apmr.2020.03.019>
69. Marshall PW, Schabrun S, Knox MF. Physical activity and the mediating effect of fear, depression, anxiety, and catastrophizing on pain related disability in people with chronic low back pain. *PLOS One*. 2017;12(7). <https://doi.org/10.1371/journal.pone.0180788>
70. Marshall PW, Morrison, NM V, Mifsud A, Gibbs M, Khan N, Meade T. The moderating effect of treatment engagement on fear-avoidance beliefs in people with chronic low back pain. *Clin J Pain*. 2021;37(12):872-880. <https://doi.org/10.1097/AJP.0000000000000991>
71. Marshal PW, Morrison NMV, Gibbs M, Schabrun SM. The effect of exercise engagement on low back disability at 12-months is mediated by pain and catastrophizing in a community sample of people with chronic low back pain. *Behav Res Ther*. 2022;159:104205.
<https://doi.org/10.1016/j.brat.2022.104205>
72. Bernard P, Ninot G, Bernard PL, et al. Effects of a six-month walking intervention on depression in inactive post-menopausal women: a randomized controlled trial. *Aging Ment Health*. 2015;19(6):485-492. <https://doi.org/10.1080/13607863.2014.948806>
73. Priore LB, Lack S, Garcia C, Azevedo FM, de Oliveira Silva D. Two weeks of wearing a knee brace compared with minimal intervention on kinesiophobia at 2 and 6 weeks in people with patellofemoral pain: a randomized controlled trial. *Arch Phys Med Rehabil*. 2020;101(4):613-623. <https://doi.org/10.1016/j.apmr.2019.10.190>
74. Strandberg E, Bean C, Vassbakk-Svindland K, et al. Who makes it all the way? Participants vs. decliners, and completers vs. drop-outs, in a 6-month exercise trial during cancer treatment. Results from the Phys-Can RCT. *Support Care Cancer*. 2022;30(2):1739-1748.
<https://doi.org/10.1007/s00520-021-06576-0>

75. Uritani D, Kasza J, Campbell PK, Metcalf B, Egerton T. The association between psychological characteristics and physical activity levels in people with knee osteoarthritis: a cross-sectional analysis. *BMC Musculoskelet Disord.* 2020;21(1):269.
<https://doi.org/10.1186/s12891-020-03305-2>
76. Alamam DM, Moloney N, Leaver A, Alsobayel HI, Mackey MG. Multidimensional prognostic factors for chronic low back pain-related disability: a longitudinal study in a Saudi population. *Spine J.* 2019;19(9),1548-1558.
<https://doi.org/10.1016/j.spinee.2019.05.010>
77. Massé-Alarie H, Beaulieu LD, Preuss R, Schneider C. Influence of paravertebral muscles training on brain plasticity and postural control in chronic low back pain. *Scand J Pain.* 2016;12(1):74-83. <https://doi.org/10.1016/j.sjpain.2016.03.005>
78. Polaski AM, Phelps AL, Smith TJ, et al. Integrated meditation and exercise therapy: a randomized controlled pilot of a combined nonpharmacological intervention focused on reducing disability and pain in patients with chronic low back pain. *Pain Med.* 2021;22(2):444-458. <https://doi.org/10.1093/pm/pnaa403>
79. Rabey M, Smith A, Beales D, Slater H, O'Sullivan P. Multidimensional prognostic modelling in people with chronic axial low back pain. *Clin J Pain.* 2017;33(10):877-891.
<https://doi.org/10.1097/AJP.0000000000000478>
80. Sandal LF, Bach K, Øverås CK, et al. Effectiveness of app-delivered, tailored self-management support for adults with lower back pain-related disability: a SELFBACK randomized clinical trial. *JAMA Intern Med.* 2021;181(10):1288-1296.
<https://doi:10.1001/jamainternmed.2021.4097>

81. Suttmiller AMB, Cavallario JM, Baez SE, Martinez JC, McCann RS. Perceived instability, pain, and psychological factors for prediction of function and disability in individuals with chronic ankle instability. *J Athl Train*. 2022;57(11-12):1048-1054.
<https://doi.org/10.4085/1062-6050-0605.21>
82. Demmelmaier I, Björk A, Dufour AB, Nordgren B, Opava, CH. Trajectories of fear-avoidance beliefs on physical activity over two years in people with rheumatoid arthritis. *Arthritis Care Res*. 2018;70(5):695-702. <http://doi.org/10.1002/acr.23419>
83. Ho-A-Tham N, Struyf N, Ting-A-Kee B, de Almeida Mello J, Vanlandewijck Y, Dankaerts W. Physical activity, fear avoidance beliefs and level of disability in a multi-ethnic female population with chronic low back pain in Suriname: a population-based study. *PLOS One*. 2022;17(10):e0276974. <https://doi.org/10.1371/journal.pone.0276974>
84. Aydemir B, Huang CH, Foucher KC. Strength and physical activity in osteoarthritis: the mediating role of kinesiophobia. *J Orthop Res*. 2022;40(5):1135-1142.
<http://doi.org/10.1002/jor.25151>
85. Baday-Keskin D, Ekinçi B. The relationship between kinesiophobia and health-related quality of life in patients with rheumatoid arthritis: a controlled cross-sectional study. *Joint Bone Spine*. 2022;89(2):105275. <http://doi.org/10.1016/j.jbspin.2021.105275>
86. Dąbek J, Knapik A, Gallert-Kopyto W, Brzęk AM, Piotrkowicz J, Gąsior Z. Fear of movement (kinesiophobia) - an underestimated problem in Polish patients at various stages of coronary artery disease. *Ann Agric Environ Med*. 2020;27(1):56-60.
<http://doi.org/10.26444/aaem/106143>

87. Knapik A, Dąbek J, Brzęk A. Kinesiophobia as a problem in adherence to physical activity recommendations in elderly Polish patients with coronary artery disease. *Patient Preference Adherence*. 2019;13:2129-2135. <http://doi.org/10.2147/PPA.S216196>
88. Minetama M, Kawakami M, Teraguchi M, et al. Associations between psychological factors and daily step count in patients with lumbar spinal stenosis. *Physiother Theory Pract*. 2022;38(10):1519-1527. <http://doi.org/10.1080/09593985.2020.1855685>
89. Pazzinatto MF, Silva DO, Willy RW, Azevedo FM, Barton CJ. Fear of movement and (re)injury is associated with condition specific outcomes and health-related quality of life in women with patellofemoral pain. *Physiother Theory Pract*. 2022;38(9):1254-1263. <http://doi.org/10.1080/09593985.2020.1830323>
90. Wasiuk-Zowada D, Brzęk A, Krzystanek E, Knapik A. Kinesiophobia in people with multiple sclerosis and its relationship with physical activity, pain and acceptance of disease. *Medicina*. 2022;58(3):414. <http://doi.org/10.3390/medicina58030414>
91. Yuksel Karsli T, Bayraktar D, Ozer Kaya D, et al. Comparison of physical activity levels among different sub-types of axial spondyloarthritis patients and healthy controls. *Mod Rheumatol*. 2021;31(6):1202-1207. <http://doi.org/10.1080/14397595.2021.1891676>
92. Birimoglu Okuyan C, Deveci E. The effectiveness of Tai Chi Chuan on fear of movement, prevention of falls, physical activity, and cognitive status in older adults with mild cognitive impairment: a randomized controlled trial. *Perspect Psychiatr Care*. 2021;57(3):1273-1281. <https://doi.org/10.1111/ppc.12684>
93. de Vries HJ, Reneman MF, Groothoff JW, Geertzen JHB, Brouwer S. Workers who stay at work despite chronic nonspecific musculoskeletal pain: do they differ from workers with

- sick leave? *J Occup Rehabil.* 2012;22(4):489-502. <https://doi.org/10.1007/s10926-012-9360-6>
94. Duarte N, Santos C, Hughes SL, Paúl C. Feasibility and impact of Fit & Strong! Program in Portuguese older adults with osteoarthritis: a pilot randomized controlled trial. *Geriatr Nurs.* 2020;41(6):804-811. <https://doi.org/10.1016/j.gerinurse.2020.04.005>
95. Gutke A, Lundberg M, Östgaard HC, Öberg B. Impact of postpartum lumbopelvic pain on disability, pain intensity, health-related quality of life, activity level, kinesiophobia, and depressive symptoms. *Eur Spine J.* 2011;20(3):440-448. <https://doi.org/10.1007/s00586-010-1487-6>
96. Ilves O, Häkkinen A, Dekker J, et al. Effectiveness of postoperative home-exercise compared with usual care on kinesiophobia and physical activity in spondylolisthesis: a randomized controlled trial. *J Rehab Med.* 2017;49(9):751-757. <https://doi.org/10.2340/16501977-2268>
97. Jakobsson M, Brisby H, Gutke A, et al. Prediction of objectively measured physical activity and self-reported disability following lumbar fusion surgery. *World Neurosurg.* 2019;121:e77-e88. <https://doi.org/10.1016/j.wneu.2018.08.229>
98. Larsson C, Ekvall Hansson E, Sundquist K, Jakobsson U. Kinesiophobia and its relation to pain characteristics and cognitive affective variables in older adults with chronic pain. *BMC Geriatr.* 2016;16:128. <https://doi.org/10.1186/s12877-016-0302-6>
99. Ledoux E, Dubois JD, Descarreaux M. Physical and psychosocial predictors of functional trunk capacity in older adults with and without low back pain. *J Manipulative Physiol Ther.* 2012;35(5):338-345. <https://doi.org/10.1016/j.jmpt.2012.04.007>

100. Lindbäck Y, Tropp H, Enthoven P, Abbott A, Öberg B. PREPARE: presurgery physiotherapy for patients with degenerative lumbar spine disorder: a randomized controlled trial. *Spine J.* 2018;18(8):1347-1355. <https://doi.org/10.1016/j.spinee.2017.12.009>
101. Lundgren S, Olausson A, Bergström G, Stenström CH. Physical activity and pain among patients with rheumatoid arthritis - a cognitive approach. *Adv Physiother.* 2005;7(2):77-83. <https://doi.org/10.1080/14038190510010322>
102. Nijs J, Meeus M, Heins M, Knoop H, Moorkens G, Bleijenberg G. Kinesiophobia, catastrophizing and anticipated symptoms before stair climbing in chronic fatigue syndrome: an experimental study. *Disabil Rehabil.* 2012;34(15):1299-1305. <https://doi.org/10.3109/09638288.2011.641661>
103. Okan F, Özsoy AZ, Sandalcı T, Yılar Erkek Z, Zincir H. The relationship between kinesiophobia and physical activity levels, gestational weight gain, and musculoskeletal pain in pregnant women. *Women Health.* 2023;63(7):551-561. <https://doi.org/10.1080/03630242.2023.2238843>
104. Roaldsen KS, Elfving B, Stanghelle JK, Talme T, Mattsson E. Fear-avoidance beliefs and pain as predictors for low physical activity in patients with leg ulcer. *Physiother Res Int.* 2009;14(3):167-180. <https://doi.org/https://doi.org/10.1002/pri.433>
105. Román-Veas J, Gutiérrez-Monclus R, López-Gil JF, et al. Baseline predictors related to functional outcomes in patients older than sixty years with complex regional pain syndrome type 1 after distal radius fracture treated conservatively: a prospective observational study. *Int Orthop.* 2023;47(9):2275-2284. <https://doi.org/10.1007/s00264-023-05880-0>

106. Roussel NA, De Kooning M, Nijs J, Wouters K, Cras P, Daenen L. The role of sensorimotor incongruence in pain in professional dancers. *Motor Control*. 2015;19(4):271-288. <https://doi.org/10.1123/ijsnem.2013-0074>
107. Saulicz M, Saulicz E, Knapik A, et al. Impact of physical activity and fitness on the level of kinesiophobia in women of perimenopausal age. *Prz Menopauzalny*. 2016;15(2):104-111. <https://doi.org/10.5114/pm.2016.61193>
108. Smith TO, Parsons S, Ooms A, et al. Randomised controlled trial of a behaviour change physiotherapy intervention to increase physical activity following hip and knee replacement: the PEP-TALK trial. *BMJ Open*. 2022;12(5):e061373. <https://doi.org/10.1136/bmjopen-2022-061373>
109. Tore NG, Oskay D, Haznedaroglu S. The quality of physiotherapy and rehabilitation program and the effect of telerehabilitation on patients with knee osteoarthritis. *Clin Rheumatol*. 2023;42(3):903-915. <https://doi.org/10.1007/s10067-022-06417-3>
110. Zhang S, Wang Z, Lin X, et al. Kinesiophobia and self-management behaviour related to physical activity in Chinese patients with coronary heart disease: the mediating role of self-efficacy. *Nurs Open*. 2023;10(1):105-114. <https://doi.org/10.1002/nop2.1283>
111. Alschuler KN, Hoodin F, Murphy SL, Rice J, Geisser ME. Factors contributing to physical activity in a chronic low back pain clinical sample: a comprehensive analysis using continuous ambulatory monitoring. *Pain*. 2011;152(11):2521-2527. <https://doi.org/10.1016/j.pain.2011.07.017>
112. Altuğ F, Ünal A, Kilavuz G, Kavlak E, Çitişli V, Cavlak U. Investigation of the relationship between kinesiophobia, physical activity level and quality of life in patients

- with chronic low back pain. *J Back Musculoskeletal Rehabil.* 2016;29(3):527-531.
<https://doi.org/10.3233/BMR-150653>
113. Carvalho FA, Maher CG, Franco MR, et al. Fear of movement is not associated with objective and subjective physical activity levels in chronic nonspecific low back pain. *Arch Phys Med Rehabil.* 2017;98(1):96-104. <http://doi.org/10.1016/j.apmr.2016.09.115>
114. Demirbüken İ, Özgül B, Kuru Çolak T, Aydoğdu O, Sarı Z, Yurdalan SU. Kinesiophobia in relation to physical activity in chronic neck pain. *J Back Musculoskeletal Rehabil.* 2016;29(1):41-47. <http://doi.org/10.3233/BMR-150594>
115. Elfving B, Andersson T, Grooten WJ. Low levels of physical activity in back pain patients are associated with high levels of fear-avoidance beliefs and pain catastrophizing. *Physiother Res Int.* 2007;12(1):14-24. <http://doi.org/10.1002/pri.355>
116. González de la Flor Á, García Pérez de Sevilla G, Domínguez Balmaseda D, Martín Vera D, Montero Martínez M, Del Blanco Muñoz JÁ. Relationship between self-efficacy and headache impact, anxiety, and physical activity levels in patients with chronic tension-type headache: an observational study. *Behav Neurol.* 2022;2022:8387249.
<http://doi.org/10.1155/2022/8387249>
117. Helmus M, Schiphorst Preuper HR, Hof AL, Geertzen JH, Reneman MF. Psychological factors unrelated to activity level in patients with chronic musculoskeletal pain. *Eur J Pain.* 2012;16(8):1158-1165. <http://doi.org/10.1002/j.1532-2149.2011.00109.x>
118. Huijnen IP, Verbunt JA, Peters ML, Seelen HA. Is physical functioning influenced by activity-related pain prediction and fear of movement in patients with subacute low back pain? *Eur J Pain.* 2010;14(6):661-666. <http://doi.org/10.1016/j.ejpain.2009.10.014>

119. Koho P, Orenius T, Kautiainen H, Haanpää M, Pohjolainen T, Hurri H. Association of fear of movement and leisure-time physical activity among patients with chronic pain. *J Rehabil Med.* 2011;43(9):794-799. <http://doi.org/10.2340/16501977-0850>
120. Lotzke H, Jakobsson M, Gutke A, et al. Patients with severe low back pain exhibit a low level of physical activity before lumbar fusion surgery: a cross-sectional study. *BMC Musculoskelet Disord.* 2018;19(1):365. <http://doi.org/10.1186/s12891-018-2274-5>
121. Navarro-Ledesma S, Pruijboom L, Lluch E, Dueñas L, Mena-Del Horno S, Gonzalez-Muñoz A. The relationship between daily physical activity, psychological factors, and vegetative symptoms in women with fibromyalgia: a cross-sectional observational study. *Int J Environ Res Public Health.* 2022;19(18):11610. <http://doi.org/10.3390/ijerph191811610>
122. Pedler A, Kamper SJ, Maujean A, Sterling M. Investigating the fear avoidance model in people with whiplash: the association between fear of movement and in vivo activity. *Clin J Pain.* 2018;34(2):130-137. <http://doi.org/10.1097/AJP.0000000000000524>
123. Sertel M, Aydoğan Arslan S, Tütün Yümin E, Demirci CS, Tarsuslu Şimşek T. Investigation of the relationship between physical activity, kinesiophobia and fear of falling in older adults with chronic pain. *Somatosens Mot Res.* 2021;38(3):241-247. <http://doi.org/10.1080/08990220.2021.1958774>
124. Verbunt JA, Sieben JM, Seelen HA, et al. Decline in physical activity, disability and pain-related fear in sub-acute low back pain. *Eur J Pain.* 2005;9(4):417-425. <http://doi.org/10.1016/j.ejpain.2004.09.011>
125. Corrigan P, Cortes DH, Pontiggia L, Silbernagel KG. The degree of tendonitis is related to symptom severity and physical activity levels in patients with midportion Achilles

- tendinopathy. *Int J Sports Phys Ther*. 2018;13(2):196-207.
<http://doi.org/10.26603/ijspt20180196>
126. Glaviano NR, Baellow A, Saliba S. Physical activity levels in individuals with and without patellofemoral pain. *Phys Ther Sport*. 2017;27:12-16.
<https://doi.org/10.1016/j.ptsp.2017.07.002>
127. Leonhardt C, Lehr D, Chenot JF, et al. Are fear-avoidance beliefs in low back pain patients a risk factor for low physical activity or vice versa? A cross-lagged panel analysis. *Psychosoc Med*. 2009;6:Doc01. <https://doi.org/10.3205/psm000057>
128. Pastor-Mira, MA, López-Roig S, Peñacoba C, Sanz-Baños Y, Lledó A, Velasco L. Predicting walking as exercise in women with fibromyalgia from the perspective of the theory of planned behavior. *Women Health*. 2020;60(4):412-425.
<https://doi.org/10.1080/03630242.2019.1662869>
129. Spaderna H, Hoffman JM, Hellwig S, Brandenburg VM. Fear of physical activity, anxiety, and depression: barriers to physical activity in outpatients with heart failure? *Eur J Health Psychol*. 2020;27(1):3-13. <http://doi.org/10.1027/2512-8442/a000042>
130. Baykal Şahin H, Kalaycıoğlu E, Şahin M. The effect of cardiac rehabilitation on kinesiophobia in patients with coronary artery disease. *Turk J Phys Med Rehabil*. 2021;67(2):203-210. <http://doi.org/10.5606/TFTRD.2021.5164>
131. Marques-Sule E, Söderlund A, Almenar L, Espí-López GV, López-Vilella R, Bäck M. Influence on kinesiophobia by disability, physical, and behavioural variables after a heart transplantation. *Eur J Cardiovasc Nurs*. 2022;21(6):537-543.
<http://doi.org/10.1093/eurjcn/zvab134>

132. Barchek AR, Dlugonski D, Baez SE, Hoch MC, Hoch J. The relationship between injury-related fear and physical activity in people with a history of anterior cruciate ligament reconstruction. *Phys Ther Sport*. 2021;50:201-205.
<https://doi.org/10.1016/j.ptsp.2021.05.010>
133. Donnarumma P, Presaghi F, Tarantino R, Fragale M, Rullo M, Delfini R. The impact of pelvic balance, physical activity, and fear-avoidance on the outcome after decompression and instrumented fusion for degenerative lumbar stenosis. *Eur Spine J*. 2017;26(2):428-433.
<http://doi.org/10.1007/s00586-016-4644-8>
134. Norte GE, Solaas H, Saliba SA, Goetschius J, Slater LV, Hart JM. The relationships between kinesiophobia and clinical outcomes after ACL reconstruction differ by self-reported physical activity engagement. *Phys Ther Sport*. 2019;40:1-9.
<http://doi.org/10.1016/j.ptsp.2019.08.002>
135. Olsson N, Karlsson J, Eriksson BI, Brorsson A, Lundberg M, Silbernagel KG. Ability to perform a single heel-rise is significantly related to patient-reported outcome after Achilles tendon rupture. *Scand J Med Sci Sports*. 2014;24(1):152-158. <http://doi.org/10.1111/j.1600-0838.2012.01497.x>
136. Zelle DM, Corpeleijn E, Klaassen G, Schutte E, Navis G, Bakker SJL. Fear of movement and low self-efficacy are important barriers in physical activity after renal transplantation. *PLOS One*. 2016;11(2):e0147609. <http://doi.org/10.1371/journal.pone.0147609>
137. Coronado RA, Robinette PE, Henry AL, et al. Bouncing back after lumbar spine surgery: early postoperative resilience is associated with 12-month physical function, pain interference, social participation, and disability. *Spine J*. 2021;21(1):55-63.
<http://doi.org/10.1016/j.spinee.2020.07.013>

138. Aykut Selçuk M, Karakoyun A. Is there a relationship between kinesiophobia and physical activity level in patients with knee osteoarthritis? *Pain Med.* 2020;21(12):3458-3469.
<http://doi.org/10.1093/pm/pnaa180>
139. Doğan N, Taşci S. Pain, physical activity, and kinesiophobia levels in individuals with knee osteoarthritis: a cross-sectional study. *Turk Klin J Nurs Sci.* 2022;14(4):1144-1154.
<https://doi:10.5336/nurses.2022-89904>
140. Kiliñç H, Karahan S, Atilla B, Kinikli Gİ. Can fear of movement, depression and functional performance be a predictor of physical activity level in patients with knee osteoarthritis? *Arch Rheumatol.* 2019;34(3):274-280.
<http://doi.org/10.5606/ArchRheumatol.2019.7160>
141. Özlü A, Akdeniz Leblebicier M. Does remission in rheumatoid arthritis bring kinesiophobia, quality of life, fatigue, and physical activity closer to normal? *Arch Rheumatol.* 2022;37(4):603-612. <http://doi.org/10.46497/ArchRheumatol.2022.9552>
142. Peres D, Tordi N, Demartino AM, Cheng JL, Sagawa Jr Y, Prati C. Relationships between physical activity levels and disease activity, functional disability and kinesiophobia in chronic rheumatic diseases. *Sci Sports.* 2023;38(5-6):607-615.
<https://doi.org/10.1016/j.scispo.2022.07.013>
143. Smulligan KL, Wingerson MJ, Seehusen CN, Little CC, Wilson JC, Howell DR. More physical activity is correlated with reduction in kinesiophobia for adolescents with persistent symptoms after concussion. *J Sport Rehabil.* 2022;32(2):196-202.
<https://doi.org/10.1123/jsr.2022-0193>

144. Sütçü G, Ayvat E, Kiliñç M. Effects of fatigue and kinesiophobia on functional capacity, physical activity and quality of life in Parkinson's disease. *Int J Rehabil Res.* 2021;44(1):65-68. <http://doi.org/10.1097/MRR.0000000000000449>
145. Igelström H, Emtner M, Lindberg E, Åsenlöf P. Physical activity and sedentary time in persons with obstructive sleep apnea and overweight enrolled in a randomized controlled trial for enhanced physical activity and healthy eating. *Sleep Breath.* 2013;17(4):1257-1266. <https://doi.org/10.1007/s11325-013-0831-6>
146. Ozer AY, Karaca S, Senocak E, Oguz S, Polat MG. Does kinesiophobia limit physical activity and quality of life in asthmatic patients? *Int J Rehabil Res.* 2022;45(3):230-236. <http://doi.org/10.1097/MRR.0000000000000534>
147. Wang J, Bai C, Zhang Z, Chen O. The relationship between dyspnea-related kinesiophobia and physical activity in people with COPD: cross-sectional survey and mediated moderation analysis. *Heart Lung.* 2023;59:95-101. <https://doi.org/10.1016/j.hrtlng.2023.02.007>
148. Crommert ME, Flink I, Gustavsson C. Predictors of disability attributed to symptoms of increased interrecti distance in women after childbirth: an observational study. *Phys Ther.* 2021;101(6):pzab064. <https://doi.org/10.1093/ptj/pzab064>
149. Atıcı E, Girgin N, Çevik Saldıran T. The effects of social isolation due to COVID-19 on the fear of movement, falling, and physical activity in older people. *Australas J Ageing.* 2022;41(3):407-413. <http://doi.org/10.1111/ajag.1306390>
150. Miller L, Ohlman T, Naugle KM. Sensitivity to physical activity predicts daily activity among pain-free older adults. *Pain Med.* 2018;19(8):1683-1692. <http://doi.org/10.1093/pm/pnx251>

151. Ohlman T, Miller L, Naugle KE, Naugle KM. Physical activity levels predict exercise-induced hypoalgesia in older adults. *Med Sci Sports Exerc.* 2018;50(10):2101-2109.
<http://doi.org/10.1249/MSS.0000000000001661>
152. Woby SR, Roach NK, Urmston M, Watson PJ. Psychometric properties of the TSK-11: a shortened version of the Tampa Scale for Kinesiophobia. *Pain.* 2005;117(1-2):137-144.
<http://doi.org/10.1016/j.pain.2005.05.029>
153. Jørgensen MB, Damsgård E, Holtermann A, Anke A, Sjøgaard K, Røe C. Properties of the Tampa Scale for Kinesiophobia across workers with different pain experiences and cultural backgrounds: a Rasch analysis. *J Appl Meas.* 2015;16(2):218-227.
<http://jampress.org/abst2015.htm>
154. Bäck M, Jansson B, Cider A, Herlitz J, Lundberg M. Validation of a questionnaire to detect kinesiophobia (fear of movement) in patients with coronary artery disease. *J Rehabil Med.* 2012;44(4):363-369. <http://doi.org/10.2340/16501977-0942>
155. Neblett R, Hartzell MM, Mayer TG, Bradford EM, Gatchel RJ. Establishing clinically meaningful severity levels for the Tampa Scale for Kinesiophobia (TSK-13). *Eur J Pain.* 2016;20(5):701-710. <http://doi.org/10.1002/ejp.795>
156. Waddell G, Newton M, Henderson I, Somerville D, Main CJ. A Fear-Avoidance Beliefs Questionnaire (FABQ) and the role of fear-avoidance beliefs in chronic low back pain and disability. *Pain.* 1993;52(2):157-168. [http://doi.org/10.1016/0304-3959\(93\)90127-B](http://doi.org/10.1016/0304-3959(93)90127-B)
157. Knapik A, Saulicz E, Gnat R. Kinesiophobia - introducing a new diagnostic tool. *J Hum Kinet.* 2011;28:25-31. <http://doi.org/10.2478/v10078-011-0019-8>
158. Shelby RA, Somers TJ, Keefe FJ, et al. Brief Fear of Movement Scale for osteoarthritis. *Arthritis Care Res.* 2012;64(6):862-871. <http://doi.org/10.1002/acr.21626>

159. Craig CL, Marshall AL, Sjöström M, et al. International physical activity questionnaire: 12-country reliability and validity. *Med Sci Sports Exerc.* 2003;35(8):1381-1395.
<http://doi.org/10.1249/01.MSS.0000078924.61453.FB>
160. Baecke JA, Burema J, Frijters JE. A short questionnaire for the measurement of habitual physical activity in epidemiological studies. *Am J Clin Nutr.* 1982;36(5):936-942.
<http://doi.org/10.1093/ajcn/36.5.936>
161. Grimby G, Börjesson M, Jonsdottir IH, Schnohr P, Thelle DS, Saltin B. The "Saltin-Grimby Physical Activity Level Scale" and its application to health research. *Scand J Med Sci Sports.* 2015;25 Suppl 4:119-125. <http://doi.org/10.1111/sms.12611>
162. Godin G, Shephard RJ. A simple method to assess exercise behavior in the community. *Can J Appl Sport Sci.* 1985;10(3):141-146.
163. Taylor HL, Jacobs DR Jr, Schucker B, Knudsen J, Leon AS, Debacker G. A questionnaire for the assessment of leisure time physical activities. *J Chronic Dis.* 1978;31(12):741-755.
[http://doi.org/10.1016/0021-9681\(78\)90058-9](http://doi.org/10.1016/0021-9681(78)90058-9)
164. Voorrips LE, Ravelli AC, Dongelmans PC, Deurenberg P, Van Staveren WA. A physical activity questionnaire for the elderly. *Med Sci Sports Exerc.* 1991;23(8):974-979.
165. Wendel-Vos GC, Schuit AJ, Saris WH, Kromhout D. Reproducibility and relative validity of the short questionnaire to assess health-enhancing physical activity. *J Clin Epidemiol.* 2003;56(12):1163-1169. [http://doi.org/10.1016/S0895-4356\(03\)00220-8](http://doi.org/10.1016/S0895-4356(03)00220-8)
166. Tegner Y, Lysholm J. Rating systems in the evaluation of knee ligament injuries. *Clin Orthop Relat Res.* 1985;198:43-49. <http://doi.org/10.1097/00003086-198509000-00007>

167. Zahiri CA, Schmalzried TP, Szuszczewicz ES, Amstutz HC. Assessing activity in joint replacement patients. *J Arthroplasty*. 1998;13(8):890-895. [http://doi.org/10.1016/s0883-5403\(98\)90195-4](http://doi.org/10.1016/s0883-5403(98)90195-4)
168. Mannerkorpi K, Hernelid C. Leisure time physical activity instrument and Physical Activity at Home and Work instrument. Development, face validity, construct validity and test-retest reliability for subjects with fibromyalgia. *Disabil Rehabil*. 2005;27:695-701. <https://doi.org/10.1080/09638280400009063>
169. Armstrong T, Bull F. Development of the World Health Organization Global Physical Activity Questionnaire (GPAQ). *J Pub Health*. 2006;14:66-70. <https://doi.org/10.1007/s10389-006-0024-x>
170. Frey I, Berg A, Grathwohl D, Keul J. Freiburg Questionnaire of physical activity--development, evaluation and application. *Soz Praventivmed*. 1999;44(2):55-64. <http://doi.org/10.1007/BF01667127>.
171. Jurca R, Jackson AS, LaMonte MJ, et al. Assessing cardiorespiratory fitness without performing exercise testing. *Am J Prev Med*. 2005;29(3):185-193. <https://doi.org/10.1016/j.amepre.2005.06.004>
172. Topolski TD, LoGerfo J, Patrick DL, Williams B, Walwick J, Patrick MMB. Peer reviewed: the Rapid Assessment of Physical Activity (RAPA) among older adults. *Prev Chronic Dis*. 2006;3(4):A118.
173. Montoye HJ, Kemper HCG, Saris WHM, Washburn RA. *Measuring physical activity and energy expenditure*. Champaign, IL:Human Kinetics. 1996.

174. Melzack R, Katz J. The McGill Pain Questionnaire: appraisal and current status. In: Turk DC, Melzack R, eds. *Handbook of Pain Assessment*. 2nd ed. The Guilford Press; 2001:35-52.
175. Ross RLP. Clinical assessment of pain. In: van Dieen JH, ed. *Assessment in Occupational Therapy and Physical Therapy*. WB Saunders; 1997:123-133.
176. Roos EM, Roos HP, Lohmander LS, Ekdahl C, Beynnon BD. Knee Injury and Osteoarthritis Outcome Score (KOOS)—development of a self-administered outcome measure. *J Orthop Sports Phys Ther*. 1998;28(2):88-96.
<http://doi.org/10.2519/jospt.1998.28.2.88>
177. Cleeland CS, Ryan KM. Pain assessment: global use of the Brief Pain Inventory. *Ann Acad Med Singap*. 1994;23(2):129-138.
178. Murray DW, Fitzpatrick R, Rogers K, et al. The use of the Oxford hip and knee scores. *J Bone Joint Surg Br*. 2007;89(8):1010-1014. <http://doi.org/10.1302/0301-620X.89B8.19424>
179. Bush JW, Kaplan RM. The quality of well-being scale (formerly the index of well-being). In: McDowell I, ed. *Measuring Health: A Guide to Rating Scales and Questionnaires*. 3rd ed. Oxford University Press; 2006:675-683.
180. Ware JE Jr. SF-36 health survey update. *Spine*. 2000;25(24):3130-3139.
<http://doi.org/10.1097/00007632-200012150-00008>
181. Freyd M. The Graphic Rating Scale. *J Educ Psychol*. 1923;14:83-102.
<https://doi.org/10.1037/h0074329>
182. Burckhardt CS, Clark SR, Bennett RM. The Fibromyalgia Impact Questionnaire: development and validation. *J Rheumatol*. 1991;18(5):728-733.

183. Melzack R, Torgerson WS. On the language of pain. *Anesthesiology*. 1971;34(1):50-59. <http://doi.org/10.1097/00000542-197101000-00017>
184. Page MJ, Sterne JAC, Higgins JPT, Egger M. Investigating and dealing with publication bias and other reporting biases in meta-analyses of health research: a review. *Res Synth Methods*. 2021;12(2):248-259. <http://doi.org/10.1002/jrsm.1468>
185. Simonsohn U, Simmons JP, Nelson LD. Better P-curves: making P-curve analysis more robust to errors, fraud, and ambitious P-hacking, a reply to Ulrich and Miller (2015). *J Exp Psychol Gen*. 2015;144(6):1146-1152. <http://doi.org/10.1037/xge0000104>
186. Coull A, Pugh G. Maintaining physical activity following myocardial infarction: a qualitative study. *BMC Cardiovasc Disord*. 2021;21(1):105. <http://doi.org/10.1186/s12872-021-01898-7>
187. Rogerson MC, Murphy BM, Bird S, Morris T. "I don't have the heart": a qualitative study of barriers to and facilitators of physical activity for people with coronary heart disease and depressive symptoms. *Int J Behav Nutr Phys Act*. 2012;9:140. <http://doi.org/10.1186/1479-5868-9-140>
188. Amirova A, Lucas R, Cowie MR, Haddad M. Perceived barriers and enablers influencing physical activity in heart failure: a qualitative one-to-one interview study. *PLOS One*. 2022;17(8):e0271743. <http://doi.org/10.1371/journal.pone.0271743>
189. Freeman AT, Hill D, Newell C, et al. Patient perceived barriers to exercise and their clinical associations in difficult asthma. *Asthma Res Pract*. 2020;6:5. <http://doi.org/10.1186/s40733-020-00058-6>
190. Ellis T, Boudreau JK, DeAngelis TR, et al. Barriers to exercise in people with Parkinson disease. *Phys Ther*. 2013;93(5):628-636. <http://doi.org/10.2522/ptj.20120279>

191. Simpson LA, Eng JJ, Tawashy AE. Exercise perceptions among people with stroke: Barriers and facilitators to participation. *Int J Ther Rehabil*. 2011;18(9):520-530.
<http://doi.org/10.12968/ijtr.2011.18.9.520>
192. Ramsey KA, Zhou W, Rojer AGM, Reijnierse EM, Maier AB. Associations of objectively measured physical activity and sedentary behaviour with fall-related outcomes in older adults: A systematic review. *Ann Phys Rehabil Med*. 2022;65(2):101571.
<http://doi.org/10.1016/j.rehab.2021.101571>
193. Gay C, Eschalier B, Levyckyj C, Bonnin A, Coudeyre E. Motivators for and barriers to physical activity in people with knee osteoarthritis: A qualitative study. *Joint Bone Spine*. 2018;85(4):481-486. <http://doi.org/10.1016/j.jbspin.2017.07.007>
194. Lööf H, Johansson UB. "A body in transformation"-An empirical phenomenological study about fear-avoidance beliefs towards physical activity among persons experiencing moderate-to-severe rheumatic pain. *J Clin Nurs*. 2019;28(1-2):321-329.
<http://doi.org/10.1111/jocn.14606>
195. Rethman KK, Mansfield C, Moeller J, et al. Kinesiophobia is associated with poor function and modifiable through interventions in people with patellofemoral pain: a systematic review with individual participant data correlation meta-analysis. *Phys Ther*. 2023;103(9):pzad074. <http://doi.org/10.1093/ptj/pzad074>
196. Prince SA, Adamo KB, Hamel ME, Hardt J, Connor Gorber S, Tremblay M. A comparison of direct versus self-report measures for assessing physical activity in adults: a systematic review. *Int J Behav Nutr Phys Act*. 2008;5:56. <http://doi.org/10.1186/1479-5868-5-56>

197. Lundberg M, Styf J, Jansson B. On what patients does the Tampa Scale for Kinesiophobia fit? *Physiother Theory Pract.* 2009;25(7):495-506.
<http://doi.org/10.3109/09593980802662160>
198. Gabelica M, Bojčić R, Puljak L. Many researchers were not compliant with their published data sharing statement: a mixed-methods study. *J Clin Epidemiol.* 2022;150:33-41. <http://doi.org/10.1016/j.jclinepi.2022.05.019>
199. Lahart IM, Boisgontier MP. Kinesiophobia and physical activity: data and Rmd script for meta-analysis. [Data set, Rmd scripts, supplementary material]. Version 1.1; 2024.
<https://doi.org/10.5281/zenodo.8216112>
200. Allen L, O'Connell A, Kiermer V. How can we ensure visibility and diversity in research contributions? How the contributor role taxonomy (CRediT) is helping the shift from authorship to contributorship. *Learn Publ.* 2019;32(1):71-74.
<https://doi.org/http://doi.org/10.1002/leap.1210>

TABLES AND FIGURES

Table 1. Sample characteristics of studies included in the systematic review.

Study (Year)	N (# women)	Mean age (SD or range)	Health status	Mean kinesiophobia (SD or range; measure)	Mean physical activity (SD or range; tool)	Level of pain (tool)	Study Design	Context of Assessment	Normalized Quality Score	Corr.	p- value
Alamam (2019)	100 (62)	40 (13.6)	Chronic low back pain	n.a. (FABQ)	n.a. (IPAQ)	4.8 (VAS)	Cohort	Clinical	7.0	r = -0.17	0.116
Alzahrani (2021)	26 (11)	43.6 (14.3)	Chronic low back pain	40.19 (9.20; TKS-17)	MPA: 76.07 min/day (34.45, wrist accelero.) VPA: 0.29 min/day (1.22; wrist accelero.) 13302 steps/day (5141; wrist accelero.)	4(3.VAS)	RCT	Clinical	8.8	r = 0.22 r = -0.05 r = -0.21	0.28 [‡] 0.003 [‡] 0.30 [‡]
Alschuler (2011)	20 (9)	46.1 (9.35)	Chronic low back pain	30.55 (TSK-17)	228 counts/min (wrist accelero.)	4.87 (NRS)	Cross-Sectional	Clinical	6.0	r = -0.48 [‡]	0.03
Altuğ (2016)	112 (73)	45.0 (14.6)	Chronic low back pain	44.30 (TSK-17)	5495 MET-min/week (IPAQ)	3.45 (VAS)	Cross-Sectional	Clinical	8.0	r = -0.096	0.313
Assadourian (2020)	147 (88)	49 (12)	Chronic low back pain	n.a. (TSK-17)	n.a. (diary; < vs. ≥ 1h/week)	6.7 (NRS)	Cross-Sectional	n.a.	8.0	r = -0.022 [†]	0.813 [†]
Atici (2022)	254 (171)	n.a. (>65)	Older adults	54.55 (KCS)	182.8 (PASE)	n.a.	Cross-Sectional	Online	9.0	ρ = -0.345	<0.001
Ayedemir (2022)	37 (25)	58.8 (8.6)	Knee osteoarthritis	40.3 (TSK-17)	4.8 (UCLA)	52.0 (KOOS-P)	Cross-Sectional	Clinical	7.0	r = -0.773	<0.05
Aykut Selcuk (2020)	67 (67) 29 (0)	60.6 (8.0) 61.6 (8.1)	Knee osteoarthritis Knee osteoarthritis	44.8 (TSK-17) 42.0 (TSK-17)	n.a. (IPAQ; low vs. moderate vs. high) n.a. (IPAQ; low vs. moderate vs. high)	4.6 (VAS) 2.9 (VAS)	Cross-Sectional	Clinical	7.0	r = -0.247 r = -0.309	0.019 0.116
Baday-Keskin (2022)	88 (67) 93 (67)	52 (n.a.) 45 (n.a.)	Rheumatoid arthritis Healthy adults	45 (TSK-17) 39 (TSK-17)	594 MET-min/week (IPAQ) 971 MET-min/week (IPAQ)	4.8 (VAS) n.a.	Cross-Sectional	Clinical	7.0	r = -0.12 n.a.	>0.05
Baez (2020)	40 (24)	24.3 (4.1)	Surgery (ACLR)	18.2 (TSK-11)	8 657 steps/day (hip pedo.) 7.7 (TAS)	81.5 (KOOS-P)	Cross-Sectional	Clinical	8.0	r = 0.181 [†] n.a.	0.265 [†] n.a.
Bahar Özdemir (2021)	101 (59)	33.9 (6.0)	Healthy	36.4 (5.8; TSK-17)	756 MET-min/week (1090; IPAQ)	3.1 (3.3; NRS)	Cross-Sectional	Online	8.0	r = -0.007 [†]	0.944 [†]
Barchek (2021)	19 (13)	22.9 (3.2)	Anterior cruciate ligament reconstruction	6 (8; FABQ)	11 237.7 steps/day (5667.7; wrist accelero.)	n.a.	Cross-Sectional	Clinical	7.0	ρ _{steps/day} = 0.12 ρ _{counts/min} = -0.13	0.63 0.59
Baykal Şahin (2021)	98 (35)	58.1 (10.4)	Coronary artery disease	41.4 (6.2; TSK-17)	839 MET-min/week (1212; IPAQ)	60.1 (27.2; SF-36)	Cross-Sectional	Clinical	8.0	r = -0.315	0.002
Bernard (2015)	121 (121)	65.5 (57-75)	Post-menopausal women	36 (TSK-17)	n.a. (PAQE)	n.a.	RCT	Clinical	8.8	r = -0.05 [†]	0.55 [†]
Carvalho (2017)	119 (82)	39.1 (11.2)	Chronic low back pain	41 (TSK-17)	6844 steps/day (hip accelero.) 296 counts/min (hip accelero.) 22 min MVPA/day (hip accelero.) 333 min LPA/day (hip accelero.) 6.7 (BHPAQ)	6.7 (NRS)	Cross-Sectional	Clinical	9.0	ρ = -0.15 [†] ρ = -0.02 ρ = -0.13 r = 0.09 r = -0.18	>0.05 >0.05 >0.05 >0.05 <0.05
Coronado (2021)	248 (126)	62.2 (11.9)	Surgery (laminectomy)	28.4 (TSK-13)	427 counts/min (hip accelerometer)	3.1 (NRS)	Cohort	n.a.	8.0	r = -0.05	>0.05
Corrigan (2018)	53 (18)	54.8 (34-65)	Achilles tendinopathy	35.4 (TSK-17)	n.a. (SGPALS)	n.a.	Cross-Sectional	Clinical	8.0	r = -0.005 [‡]	0.969
Crommert (2021)	139 (139)	37 (4.9)	Post-partum disability	n.a. (TSK-17)	n.a. (hip accelero.)	n.a.	Cross-Sectional	Online	7.0	r = -0.13	>0.05
Dı̇bek (2020)	130 (n.a) 119 (n.a)	n.a.	Coronary disease Hypertension	44.3 (TSK-Heart) 44.4 (TSK-Heart)	1545 MET-min/week (IPAQ) 1509 MET-min/week (IPAQ)	n.a.	Cross-Sectional	Clinical	8.0	r = -0.523 r = -0.410	<0.001 <0.001

	27 (n.a)		Heart valve defect	44.4 (TSK-Heart)	1308 MET-min/week (IPAQ)					r = -0.201	>0.05
	72 (n.a)		Myocardial infarction	46.7 (TSK-Heart)	1369 MET-min/week (IPAQ)					r = -0.428	<0.001
	86 (n.a)		Rhythm disorder	43.3 (TSK-Heart)	1660 MET-min/week (IPAQ)					r = -0.563	<0.001
	18 (n.a)		Stroke	49.7 (TSK-Heart)	1135 MET-min/week (IPAQ)					r = -0.868	<0.001
	15 (n.a)		Other CVD	44.2 (TSK-Heart)	2207 MET-min/week (IPAQ)					r = -0.663	0.01
Demmelmaier (2018)	2569 (1875)	60 (11)	Rheumatoid arthritis	n.a. (FABQ)	n.a. (IPAQ)	3.2 (VAS)	Cross-Sectional	Online	9.0	r = -0.07 [‡]	<0.0001
Demirbüken (2016)	99 (65)	43.5 (12.8)	Chronic neck pain	41.82 (TSK-17)	3749 MET-min/week (IPAQ)	6.47 (VAS)	Cross-Sectional	Clinical	8.0	r = -0.153	0.13 [‡]
Doğan (2022)	290 (178)	59.86 (15.64)	Knee Osteoarthritis	45.3 (8.6; TSK-17)	n.a. (IPAQ)	5.23 (1.52, VAS)	Cross-Sectional	Clinical	8.0	r = -0.061	0.299
Donnarumma (2017)	51 (12)	61.9 (13.9)	Surgery (laminectomy)	n.a. (TSK-17)	n.a. (IPAQ)	3.5 (GRS)	Cohort	Clinical	7.0	r = -0.35 [‡]	0.01
Elfving (2007)	64 (39)	47 (19-64)	Chronic low back pain	n.a. (TSK-13)	n.a. (SGPALS)	n.a.	Cross-Sectional	Online	9.0	r = -0.31 [‡]	0.010
Glaviano (2017)	20 (15)	22.2 (2.6)	Patellofemoral pain	13.6 (4.4; FABQ)	8629.7 steps/day (1665.3; wrist accelero.)	4.4 (1.9; VAS)	Cross-Sectional	Clinical	8.0	r = -0.481	0.02
González de La Flor (2022)	42 (32)	36.7 (13.2)	Chronic headache	9 (TSK-11)	n.a. (IPAQ)	7.14 (NRS)	Cross-Sectional	n.a.	8.0	ρ = 0.204	0.20 [‡]
Helmus (2012)	53 (37)	39.9 (11.3)	Chronic MSK pain	35.4 (TSK-17)	138 counts/min (hip accelero.)	5.8 (VAS)	Cross-Sectional	Clinical	8.0	r = -0.05	0.75
Ho-A-Tham (2022)	210 (210)	48.7 (16.9)	Chronic low back pain	n.a. (FABQ)	2,240 MET-min/week (GPAQ)	n.a.(NRS)	Cross-Sectional	Home Visit	8.0	r = -0.04 [‡]	0.49
Huijinen (2010)	111 (52)	44.1 (10.3)	Subacute low back pain	36.0 (TSK-17)	n.a. (trunk accelero.)	n.a. (NRS)	Cohort	Clinical	7.0	β = 0.12	>0.05
Igelström (2013)	63 (15)	55 (12)	Obstructive sleep apnea syndrome	12.4 (3.1, TSK-7)	7734 steps (3528; armband accelero.) 77 min (54)(MVPA; armband accelero.)	n.a.	Cross-Sectional	Clinical	8.0	r = -0.251	0.049
Kiliç (2019)	200 (120)	53.2 (6.0)	Knee osteoarthritis	31.8. (TSK-17)	1947 MET-min/week (IPAQ)	24.1 (OKS)	Cross-Sectional	Clinical	8.0	r = -0.693	<0.001
Knapik (2019)	135 (59)	71.9 (4.8)	Coronary artery disease	43 (TSK-Heart)	2.60 (ad-hoc questionnaire)	n.a.	Cross-Sectional	Clinical	8.0	ρ = -0.8	2.6x10 ⁻³⁵
Koppelaar (2022)	204(102)	47.68 (49.05)	Chronic low back pain	26.5 (16.1; FABQ)	77.58 minutes/day (38.85; accelero.)	5.3 (2; NRS)	RCT	Clinical	9.2	r _{MVA} =-0.06 r _{VPA} =-0.03	0.39 0.69
Koho (2011)	93 (60)	44.0 (17-68)	Chronic pain	n.a. (TSK-17)	n.a. (LTPAQ)	6.4 (VAS)	Cohort	Clinical	7.0	r = 0.10	>0.05
Leonhardt (2009)	449 (224)	45.4 (12.3)	Acute back pain	17 (FABQ)	34.7 (FQPA)	4.9 (1.7; NRS)	Cohort	Clinical	7.0	ρ = -0.93	<0.05
	338 (17)	50.4 (13.2)	Chronic back pain	18.4 (FABQ)	44.1 (FQPA)					ρ = -0.95	>0.05
Lotzke (2018)	118 (63)	46 (8)	Chronic low back pain	38.1 (TSK-17)	198 min MVPA/week (accelero.) 7493.5 steps/day (accelero.)	6.1 (VAS)	Cross-Sectional	Clinical	9.0	n.a.	n.a.
										r = -0.19 [‡]	0.034
Luthi (2018)	433 (n.a.)	n.a.	Chronic MSK pain	44.6 (TSK-17)	4.45 (BHPAQ)	4.45 (BPI-S)	Cross-Sectional	Clinical	8.0	r = 0.067 [†]	0.759 [†]
Marques-Sule (2022)	117 (51)	56 (12.1)	Heart transplantation	32.5 (TSK-11)	219 MET-min/week (IPAQ)	n.a.	Cross-Sectional	Clinical	9.0	r = -0.32	0.001
Marshall (2017)	218 (130)	36.3 (6.6)	Chronic low back pain	13.8 (5.6, FABQ))	n.a. (open-ended questions < vs. ≥ 30 min/week)	3.6 (2.3; VAS)	Cross-Sectional	Clinical	8.0	r = -0.24	0.0003 [‡]
Marshall (2021)	508 (259)	38.2 (11.8)	Chronic low back pain	13.2 (5.7, FABQ)	n.a. (open-ended questions, < vs. ≥ 30 min/week)	4.5 (2.5, VAS)	Cross-Sectional	Clinical	8.0	r = -0.28	1.3x10 ⁻¹⁰
Marshall (2022)	393 (193)	39.2 (12.2)	Chronic low back pain	13.3 (5.2, FABQ)	n.a. (AHS-derived questions, < vs. ≥ 30 min/week)	4.3 (2.3; VAS)	Cohort	Laboratory and Online	7.0	r = -0.01	0.84 [‡]
Massé-Alarie (2016)	22 (8)	n.a	Chronic low back pain	n.a. (TSK-17)	n.a MET-min/week (GPAQ)	n.a (VAS)	RCT	Clinical	7.8	r = 0.09	0.69 [‡]
Miller (2018)	52 (32)	67.4 (5.1)	Older Adults	18.9 (TSK-17)	6743 steps/day (hip accelero.)	1.4 (QWBS-P)	Cross-Sectional	Clinical	7.0	r = -0.54	<0.001
Minetama (2022)	71 (36)	71.6 (5.6)	Lumbar spinal stenosis	24.8 (TSK-11)	3601 steps/day (pedo.)	6.2 (NRS)	Cross-Sectional	Clinical	7.0	r = -0.229	0.055
Navarro-Ledesma (2022)	41 (41)	52.6 (8.0)	Fibromyalgia	27.5 (6.9; TSK-11)	29.1 (18.2; GLTEQ)	n.a.	Cross-Sectional	n.a.	7.0	r = -0.059	>0.05

Norte (2019)	77 (35)	21.6 (7.8)	Surgery (ACLR)	32.9 (6.0; TSK-17)	72.7 (34.9; GLTEQ)	91.4 (9.2; KOOS-P)	Cross-Sectional	Laboratory	7.0	$r = -0.312$	<0.05
Ohlman (2018)	52 (33)	67.4 (5.1)	Older adults	18.8 (4.5; TSK-11)	n.a. (hip accelero.)	n.a.	Cohort	Laboratory	8.0	$\rho = -0.29$	<0.05
Olsson (2014)	81 (12)	40.0 (9.6)	Achilles tendon rupture	35.9 (7.5; TSK-17)	2.9 (1.0; SGPALS)	n.a.	Cross-Sectional	Clinical and Laboratory	7.0	$\rho = -0.275$	0.013
Ozer (2022)	62 (30)	36.8 (6.1)	Asthma	39.6 (5.8; TSK-17)	2249 MET-min/week (1333; IPAQ)	n.a.	Cross-Sectional	n.a.	8.0	$r = -0.889$	0.001
Özlu (2022)	45 (45)	54.22 (8.2)	Rheumatoid arthritis	38.5 (15.7; TSK-17)	1980.7 MET-min/week (1104.8; IPAQ)	2.4 (1.7; VAS)	Controlled Trial	Clinical	7.3	$r = 0.152$	0.32
Palstam (2014)	73 (73)	50.4 (9.3)	Fibromyalgia	9.7 (6.08; FABQ)	4.47(3.64; LTPAI)	58.07 (20.29; FIQpain)	Cross-Sectional	Clinical	7.0	$\Gamma_{MPA} = 0.03$ $\Gamma_{VPA} = -0.14$	0.80 [‡] 0.23 [‡]
Pastor-Mira (2020)	274 (274)	51.8 (9.1)	Fibromyalgia	27.5 (7.1; TSK-11)	3922.40 steps/week (n.a.; pedo.)	6.51 (1.62; NRS)	Cohort	Clinical	8.0	$r = -0.01$	>0.05
Pazzinatto (2022)	92 (92)	n.a. (18-35)	Patellofemoral pain	35.3 (6.8; TSK-17)	7.8 (1.5; BHPAQ)	5.1 (2.1; VAS)	Cross-Sectional	Clinical	8.0	$\rho = -0.14$	0.18 [‡]
Pedler (2018)	103 (74)	39.7 (13.9)	Whiplash injury	26 (TSK-11)	9.9% of active time (8.2; trunk accelero.)	4.0 (2.4; VAS)	Cross-Sectional	n.a.	7.0	$r = 0.140$	>0.05
Peres (2023)	50 (38) 50 (31)	62.8 (10.9) 46.8 (12.3)	Rheumatoid arthritis Spondyloarthritis	42.7 (7.8; TSK-17) 40.8 (7.6; TSK-17)	3706.9 MET-min/week (2958.3; SQUASH) 5614.9 MET-min/week (3681.7; SQUASH)	n.a.	Cross-Sectional	n.a.	8.0	$r = -0.251$ $r = 0.170$	0.055 0.188
Polaski (2021)	38 (26)	37.6 (13.4)	Chronic low back pain	19.5 (12.6; FABQ)	2821 MET-min/week (IPAQ) 11 441 steps/day (accelero.)	3.1 (VAS)	RCT	Clinical	8.4	n.a. $r = -0.033$	n.a. 0.42
Priore (2020)	50 (37)	22.4 (3.9)	Patellofemoral pain	36.7 (TSK-17)	3088 MET-min/week (IPAQ)	n.a. (VAS)	RCT	Laboratory	9.2	$r = -0.251^{\dagger}$	0.072 [‡]
Rabey (2017)	266 (157)	51 (n.a.)	Chronic low back pain	14.3 (FABQ)	114 (0, 302.5) MET-min/week (IPAQ)	5.8 (NRS)	Cohort	Clinical	8.0	$r = 0.077$	0.25 [‡]
Roaldsen (2009)	98 (62)	76 (60-86)	Leg ulcer	12 (FABQ)	2.6 (SGPALS)	1.3 (VRS)	Cross-Sectional	Online	8.0	$r = -0.39^{\dagger}$	7.1x10 ^{-5*}
Sandal (2021)	461 (255)	45.7 (14.7)	Chronic low back pain	10.3 (5.4; FABQ)	n.a. (SGPALS)	4.9 (1.9; NRS)	RCT	Online	8.8	$r = -0.024$	0.66 [‡]
Saulicz (2016)	105 (105)	n.a.	Older adults	45.2 (15.6; KCS)	n.a. (BHPAQ)	n.a.	Cross-Sectional	Online	7.0	$r = -0.577^{\dagger}$	$<0.001^{\dagger}$
Sertel (2021)	163 (76)	71.4 (6.0)	Chronic pain	44.4 (7.7; TSK-17)	171.3 (76.2; PASE)	n.a. (VAS)	Cross-Sectional	Home Visit	7.0	$r = -0.021$	>0.05
Smulligan (2023)	23 (11)	14.9 (1.8)	Concussion (no persistant symptoms)	37 (TSK-17)	10545 steps/day (3405; wrist accelero.) 4.4 sessions/week (1.9; wrist accelero.)	n.a.	Cohort	Clinical	8.0	$r_{steps} = -0.18$ $r_{frequency} = -0.34$	0.41 0.12 0.67
	18 (9)	14.5 (2)	Concussion (persistant symptoms)	41 (TSK-17)	46.8 min/session (26.2; wrist accelero.) 7047 steps/day (2499; wrist accelero.) 2.3 sessions/week (2.1; wrist accelero.) 38 min/session (12; wrist accelero.)	n.a.				$\rho_{duration} = 0.10$ $r_{steps} = -0.60$ $r_{frequency} = -0.63$ $\rho_{duration} = -0.12$	0.008 0.05 0.65
Spaderna (2020)	61 (13)	67.5 (10.7)	Heart failure	1.5 (FActS)	2332 kcal/day (hip accelero.)	n.a.	Cross-Sectional	Laboratory	9.0	$r = -0.28$	<0.05
Strandberg (2022)	451 ^{††} (n.a.)	n.a.	Cancer	n.a. (TSK-14)	n. 1.3 h MVPA/week (arm accelero.)	n.a.	RCT	Laboratory	8.8	$r = -0.084^{\dagger}$	0.074 [‡]
Sütçü (2021)	20 (10)	69.8 (9.4)	Parkinson's disease	39.8 (7.4; TSK-17)	3078 steps/day (arm accelero.) 2055 kcal/day (475; arm accelero.)	n.a.	Cross-Sectional	n.a.	6.0	$r = -0.32^{\dagger}$ $r = -0.54^{\dagger}$	0.157 0.013
Suttmiller (2022)	126 (107)	32.69 (4.38)	Chronic pain	21.36 (5.53; TSK-11)	n.a. (JPAS)	n.a. (ad-hoc questionnaire, Yes and No)	Cross-Sectional	Online	8.0	$r = 0.219$	0.014
Uritani (2020)	167 (105)	62.2 (7.5)	Knee osteoarthritis	12.5 (BFMSO)	7998 steps/day (thigh accelero.)	5.7 (NRS)	Cross-Sectional	n.a.	7.0	$r = -0.163^{\dagger}$	0.035 [†]

Verbunt (2005)	123 (57)	44.1 (10.3)	Subacute low back pain	36.0 (TSK-17)	n.a. (hip accelero.)	4.2 (VAS)	Cross-Sectional	n.a.	7.0	$\rho = 0.06$	>0.05
Wang (2023)	223 (37)	72.35 (8.96)	Pulmonary (COPD)	20.72 (3.79; BBQ)	n.a. (IPAQ)	n.a.	Cross-Sectional	n.a.	7.0	$r = -0.35$	<0.001
Wasiuk-Zowada (2022)	80 (60)	45.5 (8.6)	Multiple sclerosis	36.6 (TSK-17)	5.1 (BHPAQ)	3.5 (VAS)	Cross-Sectional	Clinical	7.0	$r = -0.363$	0.001
Yuksel Karsli (2021)	34 (12)	41 (n.a.)	Radiographic SpA	42 (TSK-17)	2203 min LPA/day (hip accelero.)	n.a.	Cross-Sectional	Clinical	8.0	$\rho = -0.16$	0.929
	33 (10)	33 (n.a.)	Non-radiographic SpA	36 (TSK-17)	210 min MPA/day (hip accelero.) 0 min VPA/day (hip accelero.) 2576 min LPA/day (hip accelero.) 265 min MPA/day (hip accelero.) 2 min VPA/day (hip accelero.)	n.a.				$\rho = -0.158$	0.373
										$\rho = -0.394$	0.021
										$\rho = -0.001$	0.997
										$\rho = 0.013$	0.947
										$\rho = -0.240$	0.209
Zadro (2019)	60 (31)	68.3 (5.7)	Chronic low back pain	34.2 (5.9; TSK-17)	n.a.	5 (1.7; NRS)	RCT	Laboratory	3.2	$r_{MPA} = -0.18$	0.16 [†]
										$r_{VPA} = -0.12$	0.36 [†]
Zelle (2016)	487 (209)	51.6 (12.5)	Renal transplantation	n.a. (TSK-11)	165 METs (MLTPAQ and TOAQ)	n.a.	Cross-Sectional	n.a.	7.0	$r = -0.22$	<0.001

Notes. Accelerometer, ACLR = anterior cruciate ligament reconstruction, BBQ = Breathlessness Beliefs Questionnaire (17-85), BFOMSO = Brief Fear of Movement Scale for Osteoarthritis (6 – 24), BHPAQ = Baecke Habitual Physical Activity Questionnaire (3 – 15), BPI-S = Brief Pain Inventory-Severity (1 – 10), COPD = Chronic Obstructive Pulmonary Disease, CVD = cardiovascular disease, FABQ = Fear-Avoidance Belief Questionnaire (0 – 24), FActS = Fear of Activity in Situations (0 – 30), GLTEQ = Godin Leisure Time Exercise Questionnaire (0 – 119), GPAQ = Global Physical Activity Questionnaire, GRS = Graphic Rating Scale (0 – 10), IPAQ = short form of the International Physical Activity Questionnaire, JPAS = Jurka Physical Activity Scale, KCS = Kinesiophobia Causes Scale (0 – 100), KOOS-P = Knee Injury and Osteoarthritis Outcome Score – Pain (0 – 100), LPA = light physical activity, LTPAI = Leisure Time Physical Activity Index, n.a. = not available, MLTPAQ = Minnesota Leisure Time Physical Activity Questionnaire, MPA = moderate physical activity, MSK = Musculoskeletal, MVPA = moderate-to-vigorous physical activity, NRS = Numeric Rating Scale (maximum score = 0 – 10), OKS = Oxford Knee Score – Pain (0-28), PA = Physical Activity, PAQE = Physical Activity Questionnaire for the Elderly (0-3), PASE = Physical Activity Scale for the Elderly, Pedom. = Pedometer, r = Pearson's correlation coefficient, RAPAQ = Rapid Assessment of Physical Activity questionnaire (0-3), ρ = Spearman's correlation coefficient, SF-36 = 36-Item Short Form Survey (0 – 100), QWBS-P = Quality of Well-Being Scale-Pain (0-5), SGPALS = Saltin-Grimby Physical Activity Level Scale (1 – 4), SpA = axial spondyloarthritis, SQUASH = Short Questionnaire to Assess Health Enhancing Questionnaires, TAS = Tegner Activity Scale (0 – 10), TOAQ = Tecumseh Occupational Activity Questionnaire, TSK = Tampa Scale of Kinesiophobia [TSK-17: 17 – 68, TSK-Heart = 17 – 68, TSK-14 = 14 – 56, TSK-13 = 13 – 52, TSK-11 = 11 – 44], UCLA = University of California Los Angeles activity score (1 – 10), VAS = Visual Analog Scale (0 – 10), VPA = vigorous physical activity, VRS = Verbal Rating Scale for pain assessment (0 – 5), [†]Obtained by email from authors, ^{††}Number of participants used to calculate the correlation, according to the email sent by the authors, [‡]When Pearson's correlation coefficient (r) was not reported in an article, but the exact p -value and sample size (n) were available and it was possible to know the sign of the correlation based on the information provided in the article, the r value was computed using an ad-hoc R code (Supplemental Code 1A). [‡]When exact p -values were not reported in an article, but the sample size (n) and Pearson's correlation coefficient were available, the exact p -value was computed using an ad-hoc R code (Supplemental Code 1B).

Table 2. Results of the main, secondary, and subgroup meta-analyses

	n	k	o	cor.	95% CI	I² (%)	p
Main: Pearson's r estimates							<.0001
Kinesiophobia and physical activity	63	83	12278	-.19	[-.26; -.13]	85	
Secondary: Spearman's rho estimates							.0486
Kinesiophobia and device-based physical activity	12	21	2084	-.20	[-.38; .001]	86	
Subgroup: Health status							<.0001
Chronic pain	29	35	5091	-.07	[-.16; .01]	62	
Arthritis	9	11	3592	-.25	[-.39; -.10]	93	
Cardiovascular condition	5	10	823	-.30	[-.47; -.11]	29	
Neurological condition	4	8	220	-.53	[-.69; -.32]	56	
Surgery	5	5	903	-.16	[-.36; .05]	69	
Older adults	3	3	278	-.40	[-.60; -.14]	91	
Acute pain	2	2	103	-.13	[-.29; .04]	34	
Pulmonary condition	2	2	285	-.68	[-.82; -.46]	98	
Obstructive sleep apnea	1	2	146	-.18	[-.56; .26]	0	
Fibromyalgia	1	2	146	-.06	[-.46; .37]	2	
Cancer	1	1	451	-.19	[-.26; -.13]		
Post-partum women	1	1	139	-.13	[-.29; -.04]		
Young adults	1	1	101	-.01	[-.20; .19]		
Subgroup: Physical activity measure							.1714
Self-reported	44	54	9882	-.22	[-.29; -.14]	89	
Device-based	20	29	2396	-.13	[-.24; -.02]	57	
Subgroup: Physical activity instrument							.2092
Accelerometer	18	27	2462	-.17	[-.29; -.05]	55	
IPAQ	18	25	5034	-.28	[-.39; -.16]	93	
BHPAQ	4	4	737	-.34	[-.50; -.15]	94	
SGPALS	4	4	675	-.19	[-.43; .08]	80	
Ad-hoc questionnaire	4	4	1240	-.15	[-.39; .11]	85	
Pedometer	3	3	385	-.02	[-.33; .29]	59	
GPAQ	2	2	232	.01	[-.38; .40]	0	
MLTPAQ	2	2	580	-.07	[-.42; .29]	87	
GLTEQ	2	2	118	-.20	[-.54; .20]	43	
LTPAI	1	2	146	-.06	[-.52; .44]	2	
SQUASH	1	2	100	-.04	[-.52; .46]	77	
RAPAQ	1	2	120	-.15	[-.59; .30]	0	
JPAS	1	1	126	.22	[.05; .38]		
PASE	1	1	163	-.02	[-.17; .13]		
UCLA	1	1	37	-.77	[-.88; -.60]		
Diary	1	1	123	-.02	[-.20; .16]		
Subgroup: Physical activity outcome							.6098
MET-min/week	25	35	6439	-.20	[-.30; -.09]	91	
Score	15	16	2090	-.24	[-.36; -.11]	85	
Steps/day	12	13	945	-.18	[-.33; -.03]	56	
Active time	10	12	2144	-.11	[-.25; .03]	77	
Counts/min	5	5	579	-.14	[-.33; .05]	8	
Kcal/day	2	2	81	-.38	[-.67; .02]	24	
Subgroup: Kinesiophobia instrument							.4520
TSK-17	33	44	3679	-.23	[-.32; -.14]	86	
FABQ	13	15	5434	-.13	[-.27; .02]	74	
TSK-11	8	8	1259	-.04	[-.23; .15]	82	
TSK-Heart	1	7	467	-.51	[-.78; -.08]	61	
TSK-13	3	3	372	-.16	[-.44; .16]	44	
TSK-14	1	1	451	-.08	[-.17; .01]		
TSK-12	1	1	60	-.18	[-.41; .08]		
BBQ	1	1	223	-.35	[-.46; -.23]		
BFOMSO	1	1	167	-.16	[-.31; -.01]		
FActS	1	1	61	-.28	[-.50; -.03]		
KCS	1	1	105	-.57	[-.69; -.42]		

Notes. 95% CI = 95% confidence interval, BBQ = Breathlessness Beliefs Questionnaire, BFOMSO = Brief Fear of Movement Scale for Osteoarthritis, BHPAQ = Baecke Habitual

Physical Activity Questionnaire, Cor. = Correlation estimate, FABQ = Fear-Avoidance Belief Questionnaire, FActS = Fear of Activity in Situations, GLTEQ = Godin Leisure Time Exercise Questionnaire, GPAQ = Global Physical Activity Questionnaire, IPAQ = short form of the International Physical Activity Questionnaire, JPAS = Jurka Physical Activity Scale, k = number of estimates, KCS = Kinesiophobia Causes Scale, LTPAI = Leisure Time Physical Activity Index, MLTPAQ = Minnesota Leisure Time Physical Activity Questionnaire, n = number of studies, o = number of observations (participants), p = p-value for between-group difference, PAQE = Physical Activity Questionnaire for the Elderly, PASE = Physical Activity Scale for the Elderly, RAPAQ = Rapid Assessment of Physical Activity questionnaire, SGPALS = Saltin-Grimby Physical Activity Level Scale, SQUASH = Short Questionnaire to Assess Health Enhancing Questionnaires, TSK = Tampa Scale of Kinesiophobia, UCLA = University of California Los Angeles activity score.

Figure 1. PRISMA 2020 flow diagram




Figure 2. Main meta-analysis: Correlation between kinesophobia and physical activity

Notes: 95% CI = 95% confidence interval, IV = Inverse variance method, Random = Random effects method.





Figure 3. Publication bias assessment. Contour-enhanced funnel plot of the main meta-analysis (A). The vertical dashed line represents the average effect size. The two other dashed lines represent the idealized funnel-shape that studies are expected to follow. P-curve analysis (B). The blue line indicates the distribution of the analyzed p-values. The red dotted line illustrates a uniform distribution of the p-values, indicating the absence of a true effect.



SUPPLEMENTAL MATERIAL

Physical Activity and Kinesiophobia: A Systematic Review and Meta-Analysis

Miriam Goubran¹, Ata Farajzadeh¹, Ian M. Lahart², Martin Bilodeau¹,
& Matthieu P. Boissongontier^{1,3,*}

¹School of Rehabilitation Sciences, Faculty of Health Sciences, University of Ottawa, Canada; ²Faculty of Education, Health and Wellbeing, Institute of Human Sciences, University of Wolverhampton, Wolverhampton, UK; ³Perley Health Centre of Excellence in Frailty-Informed Care, Ottawa, Canada. *Corresponding author: matthieu.boissongontier@uottawa.ca

Supplemental Code 1. R scripts.

Supplemental Figure 1. Heterogeneity variance.

Supplemental Figure 2. Secondary meta-analysis based on Pearson's rho estimates.

Supplemental Figure 3. Subgroup meta-analysis: Differences by health status.

Supplemental Figure 4. Subgroup meta-analysis: Differences by physical activity measure (self-reported vs. device-based).

Supplemental Figure 5. Subgroup meta-analysis: Differences by physical activity measurement instrument.

Supplemental Figure 6. Subgroup meta-analysis: Differences by physical activity outcome

Supplemental Figure 7. Subgroup meta-analysis: Differences by kinesiophobia measurement instrument.

Supplemental Code 1. R scripts.

A. R script for the calculation of Pearson's r estimate based on the degrees of freedom (sample size - 2) and exact p-value, when the direction of the relationship is known.

```
# Sample size
n <- 50

# Pearson correlation coefficient
r <- -0.57

# Compute the t-statistic
t_statistic <- r * sqrt((n - 2) / (1 - r^2))

# Degrees of freedom
df <- n - 2

# Compute the p-value
p_value <- 2 * pt(abs(t_statistic), df = df, lower.tail = FALSE)

# Print the p-value
print(p_value)
```

B. R script for the calculation of exact p-values based on sample size (n) and Pearson's correlation coefficient (r).

```
# Sample size
n <- 45

# Pearson correlation coefficient
r <- -0.25


# Compute the t-statistic
t_statistic <- r * sqrt((n - 2) / (1 - r^2))

# Degrees of freedom
df <- n - 2


# Compute the p-value
p_value <- 2 * pt(abs(t_statistic), df = df, lower.tail = FALSE)

# Print the p-value
print(p_value)
```


Supplemental Figure 1. Heterogeneity variance.




Supplemental Figure 2. Secondary meta-analysis based on Pearson’s rho estimates.




Supplemental Figure 3. Subgroup meta-analysis: Differences according to health status




Supplemental Figure 4. Subgroup meta-analysis: Differences according to physical activity measure (self-reported vs. device-based).



Supplemental Figure 5. Subgroup meta-analysis: Differences by physical activity measurement instrument.



Supplemental Figure 6. Subgroup meta-analysis: Differences by physical activity outcome



Supplemental Figure 7. Subgroup meta-analysis: Differences by kinesiophobia measurement instrument.

