
Metric	Survey question
Overall satisfaction	1. I had an easy time signing up for the Genetic Causes of CP study (strongly agree, agree, uncertain, disagree, strongly disagree).
Comprehension (initial) retention (follow-up)	 2. Do I have to participate in the Genetic Causes of CP study? (Yes, No) 3. Are there any risks involved in participating in the study? (Yes,
(sum #2-6)	No) 4. Will the information learned in this study go on to help people in the CP community? (Yes, No)
	5. My samples and data may be shared with anyone who wants to use them, for any reason. (Yes, No)
	6. Other researchers who use my samples and data in the future will be provided with my personal information like my name, phone number, and address. (Yes, No)
Adequacy of information (#7)	7. The amount of information I was given about the study was (way too much, too much, just enough, not enough, not nearly enough)
Study clarity (#10)	8. How can we make the study better?
Privacy (#11)	 9. How can we make it easier to sign up for the study? 10. When I joined the study, I understood what I had to do to sign up (strongly agree, agree, uncertain, disagree, strongly disagree) 11. Are the privacy protections for the study strong enough? (Yes, No) 11-a. If not, what would help you feel more comfortable?
Study goals (#12)	12. When I signed up for the study, I understood that the researchers are trying to find changes in my family's DNA that cause cerebral palsy (strongly agree, agree, uncertain, disagree, strongly disagree).
Participant trust	13. Doctors who do medical research care only about what is best for each patient (agree, disagree).
(sum #13-16)	 14. Doctors tell their patients everything they need to know about being in a research study (agree, disagree) 15. Medical researchers treat people like "guinea pigs." 16. I completely trust doctors who do medical research (agree, disagree)

Supplemental Table 1. Follow-up Survey Questions

^aToo much indicates not concise; too little indicates not thorough



Supplemental Figure 2. Electronic iConsent App Representative Image

Consent Steps: 1 2	3 4 5 6 7 8 9 10		NEXT
	Image removed	Cerebral Palsy Research Network College of Medicin Neurological Institute College of CP iConsent Step 1	ne


Questions & Answers

Will you need access to the person with CP's medical records?

- "De-identified" (without any personal information) information about the person with CP's symptoms will be provided by CPRN doctors at the participating CPRN hospital.
- · Study personnel will not have direct access to medical records.
- We may contact you in the future about the chance to participate in other studies.

BACK $(\leftarrow) (\rightarrow)$ NEX

Supplemental Figure 3. Distribution of Initial (Comprehension) and Follow-up (Retention) Scores^a



^a Score represents the number of correct answers out of 5 questions

		Fathers			Mothers			Probands			Total		
Survey question	Correct response	Initial N=380	Follow-up N=66	pvalueª	Initial N=420	Follow-up N=113	pvalueª	Initial N=21	Follow-up N=6	pvalueª	Initial N=821	Follow-up N=185	pvalueª
Do I have to be part		N=164	N=29		N=183	N=53		N=10	N=4		N=357	N=86	
of the GCCP study? ^b	No	148 (90.2%)	23 (79.3%)	0.0877	104 (56.8%)	48 (90.6%)	<0.0001	10 (100.0%)	4 (100.0%)		262 (73.4%)	75 (87.2%)	0.0070
Is it my choice to be		N=216	N=37		N=237	N=60		N=11	N=2		N=464	N=99	
part of the GCCP study? ^b	Yes	212 (98.1%)	37 (100.0%)	0.4041	235 (99.2%)	57 (95.0%)	0.0254	11 (100.0%)	2 (100.0%)	-	458 (98.7%)	96 (97.0%)	0.2109
Are there risks involved in participating?	Yes	330 (86.8%)	14 (21.2%)	<0.0001	224 (53.3%)	39 (34.5%)	0.0004	12 (57.1%)	3 (50.0%)	0.7562	566 (68.9%)	56 (30.3%)	<0.0001
Will the information learned in this study be used to help people in the CP community?	Yes	367 (96.6%)	65 (98.5%)	0.4124	416 (99.0%)	112 (99.1%)	0.9474	19 (90.5%)	6 (100.0%)	0.4321	802 (97.7%)	183 (98.9%)	0.2892
My samples and data may be shared with anyone who wants to use them, and for any purpose.	No	350 (92.1%)	49 (74.2%)	<0.0001	303 (72.1%)	106 (93.8%)	<0.0001	12 (57.1%)	5 (83.3%)	0.2414	665 (81.0%)	160 (86.5%)	0.0792
Other researchers who use my samples and data in the future will be provided with my personal information like my name, phone number, and address.	No	339 (89.2%)	48 (72.7%)	0.0003	296 (70.5%)	95 (84.1%)	0.0037	13 (61.9%)	5 (83.3%)	0.3261	648 (78.9%)	148 (80.0%)	0.7459

Supplemental Table 2: Overall Initial and Follow-up Question Responses – Breakdown

^{*a*} Chi-square test ^{*b*} Question was changed during the study; participants only answered one version of the question

Supplementary Methods and Results. Sensitivity Analysis and Comparison of Initial and Follow-up Scores to Survey Question 1

Sensitivity analysis Given that many respondents were answering question 1 (Q1) incorrectly, the question was revised after qualitative analysis. This resulted in 34 respondents receiving one version for their initial survey and a different version for their follow-up survey. A sensitivity analysis was thus conducted to see whether the results were similar when these respondents were removed.

In the sensitivity analysis, 324 people were given the original version of the question on their initial surveys and 463 were given the updated version. On follow-up, 85 got the original and 66 got the updated version. All groups improved at follow-up except for fathers who had the original version of the question; 89.5% of them answered correctly on the initial survey and 79.3% answered correctly on follow-up (**Table S1**).

In the sensitivity analysis, the fathers' scores still dropped significantly (as seen with the primary analysis) at the time of follow-up (p<0.0001). The mother's scores increased as noted in the initial analysis, and the difference now exceeded the threshold for significance (p=0.0172; original p=0.0565). The proband's scores increased, but the difference was still not statistically significant (p=0.1175). Overall, the entire group of respondents had a small but significant decrease in scores (p=0.0027) (**Table S1**).

Looking at the actual change in scores for the N=151 patients who completed the follow-up survey (**Table S2b**) the father's scores still decreased significantly (p<0.0001), the mothers' scores increased significantly (p=0.0012) and the probands scores did not change significantly (p=0.500). Overall for the entire cohort, there was a small decrease in scores and this was not statistically significant (p=0.4196).

Table S1: Overall Comparison of Initial and Follow-up Scores^a, Excluding Patients Who Received Two Different Versions of Q1

	Fathers				Mothers Proba			Probands	bands		Total	
	Initial N=368	Follow-up N=54	pvalue ^b	Initial N=399	Follow-up N=92	pvalue⁵	Initial N=20	Follow-up N=5	pvalue ^b	Initial N=787	Follow-up N=151	pvalue ^b
Mean (SD)	4.6 (0.8)	3.6 (1.1)	<0.0001	3.7 (1.1)	4.1 (0.8)	0.0172	3.7 (1.2)	4.6 (0.5)	0.1175	4.1 (1.1)	4.0 (1.0)	0.0027
Median	5.0	4.0		4.0	4.0		3.5	5.0		5.0	4.0	
Q1, Q3	4.0, 5.0	3.0, 4.0		3.0, 5.0	4.0, 5.0		3.0, 5.0	4.0, 5.0		4.0, 5.0	3.0, 5.0	
Range	(1.0, 5.0)	(1.0, 5.0)		(1.0, 5.0)	(2.0, 5.0)		(2.0, 5.0)	(4.0, 5.0)		(1.0, 5.0)	(1.0, 5.0)	

^a Score is the total number of correct answers out of 5 questions

^bWilcoxon rank sum test

Table S2a: Overall Initial and Follow-up Question Responses, Excluding Patients Who Got Two Different Versions of Q1 2

Fathers				Mothers			Probands			Total			
Survey question	Correct Response	Initial N=152	Follow-up N=29	pvalue ^a	Initial N=163	Follow-up N=52	pvalueª	Initial N=9	Follow-up N=4	pvalue ^a	Initial N=324	Follow-up N=85	pvalueª
Do I have to be part of the GCCP study? ²	No	136 (89.5%)	23 (79.3%)	0.1248	91 (55.8%)	47 (90.4%)	<0.0001	9 (100.0%)	4 (100.0%)		236 (72.8%)	74 (87.1%)	0.0064

^aWilcoxon rank sum test

	Fathers			Mothers			Probands			Total			
Survey question	Correct Response	Initial N=216	Follow-up N=25	pvalueª	Initial N=236	Follow-up N=40	pvalueª	Initial N=11	Follow-up N=1	pvalueª	Initial N=463	Follow-up N=66	pvalue ^a
Is it my choice to be part of the GCCP study? ²	Yes	212 (98.1%)	25 (100.0%)	0.4926	234 (99.2%)	40 (100.0%)	0.5590	11 (100.0%)	1 (100.0%)		457 (98.7%)	66 (100.0%)	0.3523

^aWilcoxon rank sum test

Table S2b: Comparison of Initial and Follow-up Scores^a Among Respondents Who Completed Both Surveys, Excluding Patients Who Received Two Different Versions of Q1 (Score = number of correct answers out of 5 questions)

	FathersMothersN=54N=92								Probands N=5				Total N=151			
	Initial	Follow- up	Change (Initial – Follow-up)	pvalue ^b	Initial	Follow- up	Change (Initial – Follow-up)	pvalue⁵	Initial	Follow- up	Change (Initial – Follow-up)	pvalue ^b	Initial	Follow- up	Change (Initial – Follow-up)	pvalue ^b
Mean (SD)	4.7 (0.6)	3.6 (1.1)	1.0 (1.2)	<.0001	3.7 (1.1)	4.1 (0.8)	-0.4 (1.2)	0.0012	4.2 (1.1)	4.6 (0.5)	-0.4 (1.5)	0.5000	4.1 (1.1)	4.0 (1.0)	0.1 (1.4)	0.4196
Median	5.0	4.0	1.0		4.0	4.0	0.0		5.0	5.0	0.0		4.0	4.0	0.0	
Q1, Q3	4.0, 5.0	3.0, 4.0	0.0, 2.0		3.0, 5.0	4.0, 5.0	-1.0, 0.0		3.0, 5.0	4.0, 5.0	-2.0, 1.0		3.0, 5.0	3.0, 5.0	-1.0, 1.0	
Range	(3.0, 5.0)	(1.0, 5.0)	(-1.0, 4.0)		(1.0, 5.0)	(2.0, 5.0)	(-3.0, 2.0)		(3.0, 5.0)	(4.0, 5.0)	(-2.0, 1.0)		(1.0, 5.0)	(1.0, 5.0)	(-3.0, 4.0)	

^a Score is the total number of correct answers out of 5 questions ^b Wilcoxon rank sum test

			Overall satisfactio	n scores			
Factors	-2 Strongly dissatisfied	-1 Dissatisfied	0 Neutral	1 Satisfied	2 Strongly satisfied	Kruskal-Wallis P-value	
Comprehension ^b							
1	0 (0.0%)	0 (0.0%)	0 (0.0%)	1 (1.5%)	1 (0.9%)		
2	0 (0.0%)	0 (0.0%)	0 (0.0%)	4 (6.1%)	15 (13.4%)		
3	1 (100.0%)	0 (0.0%)	0 (0.0%)	5 (7.6%)	14 (12.5%)	0.0293	
4	0 (0.0%)	0 (0.0%)	2 (50.0%)	15 (22.7%)	40 (35.7%)		
5	0 (0.0%)	2 (100.0%)	2 (50.0%)	41 (62.1%)	42 (37.5%)		
Retention ^c		, , ,					
1	0 (0.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	2 (1.8%)		
2	0 (0.0%)	0 (0.0%)	0 (0.0%)	6 (9.1%)	11 (9.8%)		
3	0 (0.0%)	0 (0.0%)	0 (0.0%)	13 (19.7%)	20 (17.9%)	0.2600	
4	0 (0.0%)	0 (0.0%)	4 (100.0%)	24 (36.4%)	54 (48.2%)		
5	1 (100.0%)	2 (100.0%)	0 (0.0%)	23 (34.8%)	25 (22.3%)		
Clarity ^d	,	, ,					
-2	0 (0.0%)	0 (0.0%)	0 (0.0%)	1 (1.5%)	0 (0.0%)		
-1	0 (0.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	<.0001	
0	0 (0.0%)	1 (50.0%)	2 (50.0%)	5 (7.6%)	0 (0.0%)		
2	1 (100.0%) 0 (0.0%)	1 (50.0%) 0 (0.0%)	1 (25.0%) 1 (25.0%)	51 (77.3%) 9 (13.6%)	30 (26.8%) 82 (73.2%)		
Adequacy of information ^e							
-2	0 (0.0%)	0 (0.0%)	1 (25.0%)	1 (1.5%)	2 (1.8%)		
-1	0 (0.0%)	1 (50.0%)	1 (25.0%)	6 (9.1%)	4 (3.6%)	0.0552	
0	1 (100.0%)	1 (50.0%)	2 (50.0%)	59 (89.4%)	106 (94.6%)		
1	0 (0.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)		
2	0 (0.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)		
Privacy ^f	. ,			,		0.0364	
0	0 (0.0%)	1 (50.0%)	0 (0.0%)	3 (4.5%)	1 (0.9%)		
1	1 (100.0%)	1 (50.0%)	4 (100.0%)	63 (95.5%)	111 (99.1%)		

Supplemental Table 3. Association Between Study Elements and Overall Satisfaction Scores^a (Expanded Data)

	Overall satisfaction scores									
Factors	-2 Strongly dissatisfied	-1 Dissatisfied	0 Neutral	1 Satisfied	2 Strongly satisfied	 Kruskal-Wallis P-value 				
Study goals ^g										
-2	0 (0.0%)	0 (0.0%)	0 (0.0%)	1 (1.5%)	0 (0.0%)					
-1	0 (0.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)					
0	0 (0.0%)	0 (0.0%)	0 (0.0%)	1 (1.5%)	6 (5.4%)	<.0001				
1	0 (0.0%)	2 (100.0%)	4 (100.0%)	42 (63.6%)	34 (30.4%)					
2	1 (100.0%)	0 (0.0%)	0 (0.0%)	22 (33.3%)	72 (64.3%)					
Participant trust ^h										
0	0 (0.0%)	0 (0.0%)	0 (0.0%)	1 (1.5%)	1 (0.9%)					
1	0 (0.0%)	0 (0.0%)	0 (0.0%)	6 (9.1%)	10 (8.9%)	0.7400				
2	0 (0.0%)	1 (50.0%)	1 (25.0%)	7 (10.6%)	9 (8.0%)	0.7122				
3	0 (0.0%)	1 (50.0%)	1 (25.0%)	21 (31.8%)	31 (27.7%)					
4	1 (100.0%)	0 (0.0%)	2 (50.0%)	31 (47.0%)	61 (54.5%)					

^a Satisfaction score was scored from strongly disagree (-2) to strongly agree (2) on the survey question "I had an easy time signing up for the Genetic Causes of CP study."

^b Comprehension score was defined as the number of comprehension questions answered correctly on the initial survey out of 5 possible.

^c Retention score was defined as the number of comprehension questions answered correctly on the follow-up survey.

^d Adequacy of Information was scored from not nearly enough (-2) to way too much (2) based on the statement "The amount of information I was given about the study was ______."

e Clarity was scored from strongly disagree (-2) to strongly agree (2) on the survey question "When I joined the study, I understood what I had to do to sign up."

^f Privacy was scored as yes (1) or no (0) on the survey question "Are the privacy protections for the study strong enough?"

^g Study goal was scored from strongly disagree (-2) to strongly agree (2) on the survey question "When I signed up for the study, I understood that the researchers are trying to find changes in my family's DNA that cause cerebral palsy."

h Participant trust score was defined as the number of H-TBR scale questions answered as indicating trust. Answers indicating trust are as follows: "Doctors who do medical research care only about what is best for each patient." (Agree)

"Doctors tell their patients everything they need to know about being in a research study." (Agree)

"Medical researchers treat people like "guinea pigs." (Disagree)

"I completely trust doctors who do medical research." (Agree)