

Table S1: Global comparisons of tidal volumes (in cc) of MTC subgroup of EOS patients before and after surgery without distinguishing concave and convex sides of spinal curve. Each cell shows mean (upper number) and standard deviation (SD) (lower number) values. P values (pre-surgical vs. post-surgical) are shown using paired t-testing. Ratio r = ((post-pre)/pre) x 100%.

	Without age correction						With age correction					
EOS	LLtv	RLtv	LCWtv	RCWtv	LDtv	RDtv	LLtv	RLtv	LCWtv	RCWtv	LDtv	RDtv
Pre	31.63 19.05	41.94 22.87	20.64 13.15	24.41 13.25	12.28 7.90	19.45 12.01	46.91 20.24	44.58 23.39	25.56 12.57	23.29 8.96	25.01 9.86	25.28 13.37
Post	49.43 24.76	70.80 36.02	31.10 15.42	38.45 19.45	20.17 12.80	36.91 20.43	58.74 33.22	61.50 32.25	34.99 18.66	34.55 17.19	27.58 19.60	29.50 18.47
r (%)	56.28	68.81	50.68	57.52	64.25	89.77	25.22	37.95	36.89	48.35	10.28	16.69
p	0.004	<0.001	0.007	0.001	0.006	<0.001	0.071	0.005	0.034	0.001	0.427	0.224

MTC – main thoracic curve; EOS – early onset scoliosis; LLtv – left lung tidal volume; RLtv – right lung tidal volume; LCWtv – left chest wall tidal volume; RCWtv – right chest wall tidal volume; LDtv – left hemi-diaphragm tidal volume; RDtv – right hemi-diaphragm tidal volume.