


Supplemental Tables and Figures

**Supplemental Table 1. Genetic ancestry reporting from each genetic ancestry provider and Apex-reported race/ethnicity.**

Genetic Ancestry Groupings	Ancestry 1	Ancestry 2	Apex-Reported Race/Ethnicity
African (AFR)	African Ancestry (SW US) African Caribbean Luhya (Kenya) Mende (Sierra Leone) Gambian Mendinka Esan (S Nigeria) Yoruba (Nigeria & Benin)	Not specified	Black or African American
Admixed American (AMR)	Puerto Rican Peruvian Colombian Mexican	Not specified	Hispanic or Latino American Indian or Alaska Native
East Asian (EAS)	Northern Chinese Southern Chinese Vietnamese Dai (SW China) Japanese	Not specified	Asian Hawaii Native or Other Pacific Islander
European (EUR)	English & Scottish Italian Finnish N&W European Iberian (Spain & Portugal)	Not specified	White or Caucasian
South Asian (SAS)	Punjabi (Pakistan) Bengali Gujarati (W India) Tamil Nadu-India Telugu (SE India)	Not specified	Asian



**Supplemental Figure 1. Flowchart of the 3D Health Study participants active and recontacted to receive genetic ancestry reports and complete the additional ancestry perspectives survey.**

**Supplemental Table 2. 3D Health Study Participant Demographics.**

Variable	Completed Survey (N=166) n, %
<b>Gender</b>	
Man	64 (38.6)%
Woman	101 (60.8)%
Non-binary	0 (0.0)%
Prefer not to answer	1 (0.6)%
<b>Age</b>	
Below 30	7 (4.2)%
30-45	49 (29.5)%
45-60	27 (16.3)%
60 or above	82 (49.4)%
Prefer not to answer	1 (0.6)%
<b>Race/Ethnicity <sup>a</sup></b>	
American Indian/Alaska Native	1 (0.6)%
Asian	34 (20.5)%
Black/African American	4 (2.4)%
Hispanic/Latino	9 (5.4)%
Native Hawaiian/Pacific Islander	0 (0.0)%
White	124 (74.7)%
Other	10 (6.0)%
<b>Had Prior Genetic Ancestry testing</b>	
Yes	74 (44.6)%
No	89 (53.6)%
Missing	3 (1.8)%

<sup>a</sup> Participants were allowed to select more than one race/ethnicity.

**Supplemental Table 3. UCSF Provider demographics**

Variable	Completed Survey (N=31) n, %
<b>Gender</b>	
Man	5 (16.1)%
Woman	25 (80.6)%
Non-binary	0 (0.0)%
Prefer not to answer	1 (3.2)%
<b>Age</b>	
Below 30	0 (0.0)%
30-45	18 (58.1)%
45-60	11 (35.5)%
60 or above	2 (6.5)%
Prefer not to answer	0 (0.0)%
<b>Race/Ethnicity <sup>a</sup></b>	
American Indian/Alaska Native	0 (0.0)%
Asian	9 (29.0)%
Black/African American	2 (6.5)%
Hispanic/Latino	2 (6.5)%
Native Hawaiian/Pacific Islander	0 (0.0)%
White	21 (67.7)%
Other	3 (9.7)%
<b>Specialty</b>	
Internal Medicine	12 (38.7)%
Family Medicine	2 (6.5)%
Genetics	1 (3.2)%
Genetic Counseling	14 (45.2)%
Other	2 (6.5)%
<b>Had Prior Genetic Ancestry testing</b>	
Yes	5 (16.1)%
No	26 (83.9)%

<sup>a</sup> Participants were allowed to select more than one race/ethnicity.

**Supplemental Table 4. Concordance between genetic ancestry calculations.** Genetic ancestry results from two commercial genetic testing providers and two 3<sup>rd</sup>-party analyses were compared using genetic samples in the UCSF 3D Health Study. The agreement was determined by the % of overlapping ancestry from each ancestry location (European, African, Asian, and Admixed American). P values were determined using a Wilcoxon Rank sum test.

Ancestry 1 vs. Ancestry 3 (n=460)					P-values		
Agreement	Min	Median	Max	Mean			
Overall	0.00%	54.65%	94.91%	57.00%			
Genetic Ancestry					Afr	As	Adm
European (333)	26.00%	53.41%	78.03%	53.34%	6.0e-5**	6.6e-22**	0.63
African (9)	53.57%	79.99%	94.91%	76.18%		0.37	9.8e-4**
Asian (75)	36.45%	78.26%	89.94%	72.90%			5.7e-6**
Admixed American (18)	0.00%	53.34%	72.78%	52.52%			
Ancestry 1 vs. Ancestry 4 (n=460)							
Agreement	Min	Median	Max	Mean			
Overall	0.00%	54.79%	95.21%	57.26%			
Genetic Ancestry							
European (333)	32.27%	52.96%	79.67%	53.12%	2.5e-3**	1.6e-29**	1.3e-3**
African (9)	47.92%	79.99%	95.21%	73.69%		0.75	0.076
Asian (75)	38.66%	78.67%	89.94%	74.53%			9.4e-6**
Admixed American (18)	0.00%	62.70%	74.77%	58.21%			
Ancestry 2 vs. Ancestry 3 (n=465)							
Agreement	Min	Median	Max	Mean			
Overall	33.50%	99.87%	100.00%	95.80%			
Genetic Ancestry							
European (337)	33.50%	99.90%	100.00%	97.24%	7.5e-4**	4.0e-3**	1.3e-9**
African (9)	92.73%	94.53%	97.10%	94.97%		2.1e-4**	6.0e-4**
Asian (74)	78.60%	100.00%	100.00%	97.62%			3.7e-10**
Admixed American (18)	38.72%	65.15%	100.00%	66.83%			
Ancestry 2 vs. Ancestry 4 (n=465)							
Agreement	Min	Median	Max	Mean			
Overall	58.68%	100.00%	100.00%	95.70%			
Genetic Ancestry (n)							
European (337)	58.68%	100.00%	100.00%	97.33%	3.5e-5**	2.4e-15**	2.3e-12**
African (9)	83.13%	95.52%	98.62%	94.37%		0.040*	3.1e-3**
Asian (74)	71.67%	97.97%	100.00%	94.65%			6.3e-6**
Admixed American (18)	58.72%	75.65%	100.00%	78.32%			
Ancestry 3 vs. Ancestry 4 (n=474)							
Agreement	Min	Median	Max	Mean			
Overall	58.77%	99.41%	100.00%	97.05%			
Genetic Ancestry (n)							


European (344)	71.90%	99.72%	100.00%	98.76%	2.3e-4**	1.3e-13**	6.0e-9**
African (9)	81.30%	98.48%	99.01%	96.39%		0.90	6.2e-3*
Asian (76)	62.90%	97.91%	100.00%	93.54%			1.7e-3**
Admixed American (18)	58.77%	89.40%	100.00%	86.08%			

\*P < 0.05; \*\*P<0.005

**Supplemental Table 5. Pare-wise comparisons of the concordance of the four ancestries when compared to Apex-reported ancestry.** McNemar's test was used on each pairwise comparison.


Comparison	Threshold		
	50%	75%	90%
Ancestry 1/2	5.0e-24**	1.7e-70**	9.6e-74**
Ancestry 1/3	1.4e-21**	6.4e-71**	3.8e-81**
Ancestry 1/4	1.5e-22**	1.4e-71**	3.1e-82**
Ancestry 2/3	0.21	>0.99	4.0e-7**
Ancestry 2/4	0.23	0.45	1.1e-8**
Ancestry 3/4	>0.99	0.131	0.041*

\*P < 0.05; \*\*P<0.005



**Supplemental Figure 2. 3D Health Participant concerns about the possible unintended outcomes of genetic ancestry testing.**





**Figure 3. 3D Health Participant concerns on the use of genetic ancestry reports in the electronic health record.**

**Supplemental Table 6. Provider perspectives on the integration of genetic ancestry into the electronic health record.** Responses were totaled for the overall responses from all providers and then calculated based on the participant's self-identified race/ethnicity. Providers who selected more than one race/ethnicity that did not fit into one of the general ancestry categories were placed into other. A Fisher's Exact test was used to determine if there was a statistically significant difference in responses between ancestry groups.

Genetic ancestry reports should be in the electronic health record.	Strongly disagree	Somewhat disagree	Neutral	Somewhat agree	Strongly agree	P-value
	2 (6.5)%	6 (19.4)%	16 (51.6)%	7 (22.6)%	0 (0.0)%	
European (n=18)	1 (5.6)%	4 (22.2)%	8 (44.4)%	5 (27.8)%	0 (0.0)%	0.9913
African (n=1)	0 (0.0)%	0 (0.0)%	1 (100.0)%	0 (0.0)%	0 (0.0)%	
Asian (n=7)	1 (14.3)%	1 (14.3)%	4 (57.1)%	1 (14.3)%	0 (0.0)%	
Admixed American (n=1)	0 (0.0)%	0 (0.0)%	1 (100.0)%	0 (0.0)%	0 (0.0)%	
Other (n=4)	0 (0.0)%	1 (25.0)%	2 (50.0)%	1 (25.0)%	0 (0.0)%	