

**Supplemental Table S5: Data describing missense variant effects on NMDAR properties.**

<i>GRIN</i> -variant	Glutamate EC <sub>50</sub> (95% CI), μM	Glycine EC <sub>50</sub> (95% CI), μM	Mg <sup>2+</sup> IC <sub>50</sub> (95% CI), μM	P <sub>OPEN</sub>	τ <sub>WEIGHT</sub> (ms)	Surface Expression (% of total)
<b>WT 1/2A*</b>	<b>3.6</b>	<b>1.2</b>	<b>20</b>	<b>0.23</b>	<b>60</b>	<b>100</b>
<b>1-K531R/2A</b>	9.9 (8.9, 11.1)	2.7 (2.4, 2.9)	18 (14, 24)	0.098 ± 0.0052	26 ± 4.1	113 ± 12
<b>1-Q556R/2A</b>	1.1 (0.64, 1.9)	0.42 (0.34, 0.51)	29 (22, 39)	0.48 ± 0.018	200 ± 49	113 ± 3.1
<b>1-P557R/2A<sup>a</sup></b>	0.37 (0.34, 0.41)	0.088 (0.080, 0.10)	28 (20, 39)	0.026 ± 0.0016	395 ± 32	39 ± 0.099
<b>1-G638S/2A</b>	3.3 (2.6, 4.1)	1.1 (0.95, 1.3)	83 (58, 119)	0.24 ± 0.010	54 ± 5.6	60 ± 16
<b>1-C744G/2A</b>	0.71 (0.51, 1.0)	1.3 (1.1, 1.5)	26 (21, 30)	0.33 ± 0.011	163 ± 16	101 ± 6.7
<b>1-A806E/2A</b>	0.28 (0.23, 0.33)	0.073 (0.055, 0.10)	18 (16, 21)	0.30 ± 0.011	441 ± 64	116 ± 1.6
<b>1-R844L/2A</b>	2.4 (2.1, 2.9)	0.88 (0.76, 1.0)	54 (46, 63)	0.35 ± 0.022	98 ± 13	140 ± 10
<b>WT 1/2A*</b>	<b>3.2</b>	<b>1.2</b>	<b>26</b>	<b>0.17</b>	<b>54</b>	<b>100</b>
<b>2A-R518C</b>	3600 (3260, 4000)	1.5 (1.2, 1.9)	21 (17, 25)	tstm	tstm	32 ± 4.0
<b>2A-T690M</b>	5900 (4700, 7300)	1.4 (1.1, 1.7)	29 (24, 34)	tstm	tstm	30 ± 5.2
<b>2A-I694T<sup>b</sup></b>	9.7 (9.0, 11)	0.87 (0.63, 1.2)	28 (20, 40)	0.13 ± 0.0060	45 ± 3.7	72 ± 8.0
<b>2A-R695Q</b>	0.32 (0.28, 0.35)	1.2 (1.1, 1.4)	15 (11, 21)	0.20 ± 0.01	257 ± 34	77 ± 23
<b>2A-A716D</b>	tstm	tstm	tstm	tstm	tstm	n.a.
<b>2A-K772E<sup>b</sup></b>	4.8 (4.4, 5.2)	1.3 (1.2, 1.5)	13 (8.5, 21)	0.14 ± 0.005	47 ± 5.6	18 ± 2.0
<b>2A-L812M<sup>c</sup></b>	0.40 (0.34, 0.45)	0.14 (0.12, 0.15)	36 (32, 41)	0.40 ± 0.0098	443 ± 47	85 ± 13
<b>2A-M817V<sup>d</sup></b>	0.28 (0.20, 0.39)	0.14 (0.11, 0.17)	60 (41, 87)	0.93 ± 0.070	632 ± 103	100 ± 14
<b>2A-A818V</b>	1.6 (1.3, 2.0)	0.38 (0.32, 0.45)	40 (32, 50)	0.713 ± 0.022	86 ± 9.6	155 ± 55
<b>WT 1/2B*</b>	<b>1.3</b>	<b>0.3</b>	<b>23.0</b>	<b>0</b>	<b>716</b>	<b>100</b>
<b>2B-G543R</b>	0.22 (0.19, 0.26)	0.094 (0.08, 0.12)	26 (22, 31)	0.013 ± 0.0023	3530 ± 223	110 ± 11
<b>2B-V558I</b>	1.0 (0.84, 1.2)	0.39 (0.33, 0.47)	24 (20, 30)	0.0066 ± 0.001	376 ± 110	103 ± 7.5
<b>2B-W559C</b>	tstm	tstm	tstm	tstm	tstm	74 ± 12
<b>2B-W559R</b>	tstm	tstm	tstm	tstm	tstm	81 ± 1.9
<b>2B-N616I</b>	0.22 (0.14, 0.35)	0.039 (0.28, 0.55)	>5000	0.016 ± 0.0032	2240 ± 254	82 ± 25
<b>2B-N616S</b>	0.91 (0.81, 1.0)	0.23 (0.20, 0.28)	>5000	0.010 ± 0.00052	1140 ± 58	83 ± 8.1
<b>2B-M654I</b>	0.059 (0.045, 0.078)	0.034 (0.022, 0.055)	42 (30, 59)	0.032 ± 0.0060	5720 ± 647	67 ± 11
<b>2B-M654V</b>	0.057 (0.042, 0.077)	0.087 (0.062, 0.12)	38 (29, 50)	0.14 ± 0.020	1410 ± 97	25 ± 8.4
<b>2B-T691A</b>	1230 (1090, 1380)	0.46 (0.44, 0.48)	28 (21, 36)	tstm	tstm	96 ± 19
<b>2B-T691I</b>	2700 (2490, 2920)	0.39 (0.32, 0.46)	22 (16, 29)	tstm	tstm	119 ± 25
<b>2B-R696C</b>	0.22 (0.19, 0.25)	0.23 (0.20, 0.28)	28 (22, 34)	0.023 ± 0.0020	1880 ± 151	96 ± 25
<b>2B-A717E</b>	tstm	tstm	tstm	tstm	tstm	41 ± 17
<b>2B-C746W</b>	0.87 (0.72, 1.1)	2.7 (2.2, 3.3)	19 (16, 23)	0.013 ± 0.00089	597 ± 55	47 ± 8.2
<b>2B-L797F</b>	39 (26, 59)	0.67 (0.61, 0.74)	24 (20, 28)	0.051 ± 0.0031	38 ± 4.0	89 ± 5.6
<b>2B-L813V<sup>e</sup></b>	0.27 (0.23, 0.31)	0.021 (0.013, 0.034)	24 (21, 29)	0.021 ± 0.0023	1360 ± 101	83 ± 7.2
<b>2B-M818R</b>	tstm	tstm	tstm	tstm	tstm	39 ± 8.0
<b>2B-M818T<sup>e</sup></b>	0.42 (0.30, 0.58)	0.13 (0.10, 0.19)	15 (8.5, 26)	0.15 ± 0.015	1680 ± 440	63 ± 17
<b>2B-G820A</b>	1.3 (0.96, 1.9)	0.32 (0.26, 0.39)	22 (19, 24)	0.011 ± 0.0016	205 ± 10	84 ± 7.6

\* WT values were recorded on the same day for every variant, and the mean for each parameter is given

Mean values for variant receptors are given with 95% confidence intervals determined from the log EC<sub>50</sub> or IC<sub>50</sub> values, or with SEM. All data are from 8-23 oocytes, 4-15 HEK cells, or 2-10 independent surface expression

tstm indicates the response was too small to measure, n.a. indicates data is not available

a Part of the data set were reanalyzed from data in Ogden et al. 2017

- b Part of the data set were reanalyzed from data in Swanger et al. 2016
- c Part of the data set were reanalyzed from data in Yuan et al. 2014
- d Part of the data set were reanalyzed from data in Chen et al. 2017
- e Part of the data set were reanalyzed from data in Brock et al. 2023