

Supplemental Table S4: Classification of enw variants reported here according to the criteria of Myers et al. (2023)

GRIN -variant	Glutamate Potency	Glycine Potency	Mg ²⁺ IC ₅₀	P _{OPEN}	τ _{WEIGHT}	Surface Expression	Count (High, Mod)	Charge Transfer ^a	Classification
1-K531R/2A	H (0.46)	H (0.50)	1.0	H (0.38)	M (0.51)	1.1	1, 3	0.27, 0.19	Likely LoF
1-Q556R/2A	H (3.4)	H (3.1)	M (1.5)	H (2.3)	H (4.0)	1.1	4, 1	7.1, 13	Likely GoF
1-P557R/2A	H (10)	H (14)	1.3	H (0.12)	H (7.3)	M (0.38)	Conflict	0.93, 2.8	Possible GoF
1-G638S/2A	1.3	1.3	H (4.1)	1.2	0.77	M (0.60)	Conflict	2.9, 5.3	Possible GoF
1-C744G/2A	H (3.9)	0.80	1.2	M (1.6)	M (2.3)	1.0	2, 1	5.3, 13	Likely GoF
1-A806E/2A	H (10)	H (14)	0.89	1.2	H (8.7)	1.2	3, 0	9.1, 14	Likely GoF
1-R844L/2A	1.3	1.4	H (2.4)	1.4	1.4	1.4	subthreshold	3.7, 3.6	Possible GoF
2A-R518C	H (0.00053)	0.90	0.81	tstm	tstm	H (0.32)	tstm	tstm	Likely LoF*
2A-T690M	H (0.00050)	0.86	1.2	tstm	tstm	H (0.30)	tstm	tstm	Likely LoF*
2A-I694T	H (0.34)	1.3	1.1	0.72	0.80	0.72	1, 0	0.51, 0.16	Likely LoF
2A-R695Q	H (11)	1.0	1.0	1.2	H (3.6)	0.77	2, 0	2.9, 14	Likely GoF
2A-A716D	tstm	tstm	tstm	tstm	tstm	n.a.	tstm	tstm	Likely LoF*
2A-K772E	M (0.62)	0.9	M (0.63)	0.78	0.84	H (0.18)	1, 2	0.083, 0.054	Likely LoF
2A-L812M	H (8.7)	H (14)	M (2.0)	H (2.2)	H (9.4)	0.85	4, 1	32, 49	Likely GoF
2A-M817V	H (13)	H (6.7)	H (2.9)	H (5.81)	H (15)	1.0	5,	579, 860	Likely GoF
2A-A818V	M (2.4)	M (2.4)	M (2.4)	H (4.1)	M (1.7)	M (1.6)	1, 5	24, 43	Likely GoF
2B-G543R	H (5.4)	H (3.4)	1.2	M (0.65)	H (3.3)	1.1	Conflict	1.8, 3.7	Possible GoF
2B-V558I	1.2	0.92	0.86	0.06	0.53	1.0	1, 1	0.03, 0.07	Likely LoF
2B-W559C	tstm	tstm	tstm	tstm	tstm	0.74	tstm	tstm	Likely LoF*
2B-W559R	tstm	tstm	tstm	tstm	tstm	0.81	tstm	tstm	Likely LoF*
2B-N616I	H (5.1)	H (5.8)	H (1641)	0.75	H (3.2)	0.82	4, 0	95, 191	Likely GoF
2B-N616S	1.3	1.3	H (1778)	H (0.28)	M (1.6)	0.83	Conflict	11, 9.3	Possible GoF
2B-M654I	H (20)	H (9.9)	M (1.8)	0.86	H (9.8)	0.67	3, 1	6.1, 11	Likely GoF
2B-M654V	H (21)	H (3.9)	M (1.8)	H (3.8)	H (2.1)	H (0.25)	Conflict	2.1, 18	Possible GoF
2B-T691A	H (0.00073)	0.94	1.1	tstm	tstm	0.96	tstm	tstm	Likely LoF*
2B-T691I	H (0.0011)	0.81	1.1	tstm	tstm	1.2	tstm	tstm	Likely LoF*
2B-R696C	H (4.6)	0.96	0.99	0.82	H (2.7)	0.96	2, 0	2.8, 5.9	Likely GoF
2B-A717E	tstm	tstm	tstm	tstm	tstm	M (0.41)	tstm	tstm	Likely LoF*
2B-C746W	1.0	H (0.12)	0.71	0.68	0.86	M (0.47)	2, 0	0.078, 0.095	Likely LoF
2B-L797F	H (0.041)	M (0.65)	0.93	1.4	H (0.054)	0.89	2, 1	0.062, 0.019	Likely LoF
2B-L813V	H (5.4)	H (13)	0.89	1.1	M (1.9)	0.83	2, 1	1.2, 4.6	Likely GoF
2B-M818R	tstm	tstm	tstm	tstm	tstm	M (0.39)	tstm	tstm	Likely LoF*
2B-M818T	M (2.5)	M (2.2)	0.68	H (3.4)	H (2.4)	M (0.63)	Conflict	3.5, 4.3	Possible GoF
2B-G820A	1.0	1.0	0.91	H (0.49)	H (0.31)	0.84	2, 0	0.16, 0.53	Likely LoF

Fold change was calculated from ration of fitted parameters for mutant and WT controls recorded o the same day.

All calls were made by criteria of Myers et al., (2023). N.a.a indicates data not available.

tstm indicates that the current response was too small to measure despite measureable surface expression