

Can drinking water, sanitation, handwashing, and nutritional interventions reduce antibiotic use in young children?

Supplemental Information

Table of Contents

Item	Pages
Table S1. CONSORT Checklist	2-3
Table S2. Enrollment characteristics by intervention group within the EED substudy and the WASH Benefits parent trial in Bangladesh	4
Table S3. Enrollment characteristics by intervention group within the EED substudy and the WASH Benefits parent trial in Kenya	5
Table S4. Enrollment characteristics of children in the EED substudy included in follow-up vs. lost to follow-up in Bangladesh	6
Table S5. Enrollment characteristics of children in the EED substudy included in follow-up vs. lost to follow-up in Kenya	7
Table S6. Caregiver-reported antibiotic use in last 3 months by young children in subgroups of age and sex, Bangladesh	8
Table S7. Caregiver-reported antibiotic use in last 3 months by young children in subgroups of age and sex, Kenya	9
Table S8. WSH, nutrition and N+WSH interventions vs. controls on antibiotic use within last 3 months among children in Bangladesh	10
Table S9. Combined N+WSH intervention vs. WSH and nutrition interventions on antibiotic use within last 3 months among children in Bangladesh	11
Table S10. WSH, nutrition and N+WSH interventions vs. controls on antibiotic use within last 3 months among children in Kenya	12
Table S11. Combined N+WSH intervention vs. WSH and nutrition interventions on antibiotic use within last 3 months among children in Kenya	13
Table S12. Multiplicative interaction by age and sex for unadjusted intervention effects on antibiotic use within last 3 months, Bangladesh	14
Table S13. Additive interaction by age and sex for unadjusted intervention effects on antibiotic use within last 3 months, Bangladesh	15
Table S14. Multiplicative interaction by age and sex for unadjusted intervention effects on antibiotic use within last 3 months, Kenya	16
Table S15. Additive interaction by age and sex for unadjusted intervention effects on antibiotic use within last 3 months, Kenya	17
Table S16. Sensitivity analysis, unadjusted intervention effects on antibiotic use within last 2 weeks and last month, Bangladesh*	18
Figure S1. CONSORT Diagram for the WASH Benefits Bangladesh environmental enteric dysfunction substudy	19
Figure S2. CONSORT Diagram for the WASH Benefits Kenya environmental enteric dysfunction substudy	20

Table S1. CONSORT Checklist

Item	Description	Reported in Section
Title and Abstract		
1a	Identification as a randomized trial in the title; Identification as a cluster randomized trial in the title	Title
1b	Structured summary of trial design, methods, results, and conclusions	Abstract
Introduction		
Background and Objectives		
2a	Scientific background and explanation of rationale; Rationale for using a cluster design	Introduction (paragraph 1)
2b	Specific objectives or hypotheses; Whether objectives pertain to the cluster level, the individual participant level, or both	Introduction (paragraph 2)
Methods		
Trial Design		
3a	Description of trial design (such as parallel, factorial) including allocation ratio; Definition of cluster and description of how the design features apply to the clusters	Methods: Study design, Data collection (paragraph 1)
3b	Important changes to methods after trial commencement (such as eligibility criteria), with reasons	N/A
Participants		
4a	Eligibility criteria for participants; Eligibility criteria for clusters	Methods: Study design
4b	Settings and locations where the data were collected	Methods: Study design
Interventions		
5	The interventions for each group with sufficient details to allow replication, including how and when they were actually administered; Whether interventions pertain to the cluster level, the individual participant level, or both	Methods: Study design, Interventions (paragraph 1)
Outcomes		
6a	Completely defined pre-specified primary and secondary outcome measures, including how and when they were assessed; Whether outcome measures pertain to the cluster level, the individual participant level, or both	Methods: Data collection (paragraph 2), Statistical parameters and estimation strategy (paragraph 1)
6b	Any changes to trial outcomes after the trial commenced, with reasons	N/A
Sample Size		
7a	How sample size was determined; Method of calculation, number of cluster(s) (and whether equal or unequal cluster sizes are assumed), cluster size, a coefficient of intracluster correlation (ICC or k), and an indication of its uncertainty	Methods: Minimum detectable effect size
7b	When applicable, explanation of any interim analyses and stopping guidelines	N/A
Randomization		
Sequence Generation		
8a	Method used to generate the random allocation sequence	Methods: Study design
8b	Type of randomization; details of any restriction (such as blocking and block size); Details of stratification or matching if used	Methods: Study design
Allocation Concealment Mechanism		
9	Mechanism used to implement the random allocation sequence (such as sequentially numbered containers), describing any steps taken to conceal the sequence until interventions were assigned; Specification that allocation was based on clusters rather than individuals and whether allocation concealment (if any) was at the cluster level, the individual participant level, or both	Methods: Study design
Implementation		
10a	Who generated the random allocation sequence, who enrolled clusters, and who assigned clusters to interventions	Methods: Study design
10b	Mechanism by which individual participants were included in clusters for the purposes of the trial (such as complete enumeration, random sampling)	Methods: Study design
10c	From whom consent was sought (representatives of the cluster, or individual cluster members, or both) and whether consent was sought before or after randomization	Methods: Ethics
Blinding		
11a	If done, who was blinded after assignment to interventions (for example, participants, care providers, those assessing outcomes) ^[11] and how	Methods: Data collection (paragraph 2)
11b	If relevant, description of the similarity of interventions	N/A

Item	Description	Reported in Section
Statistical Methods		
12a	Statistical methods used to compare groups for primary and secondary outcomes; How clustering was taken into account	Methods: Statistical parameters and estimation strategy (paragraph 2)
12b	Methods for additional analyses, such as subgroup analyses and adjusted analyses	Methods: Statistical parameters and estimation strategy (paragraph 2), Subgroup analyses, Sensitivity analyses
Results		
Participant Flow		
13a	For each group, the numbers of participants/clusters who were randomly assigned, received intended treatment, and were analyzed for the primary outcome	Results: Enrollment, Fig S1, Fig S2
13b	For each group, losses and exclusions after randomization, together with reasons, for both clusters and individual cluster members	Results: Enrollment, Fig S1, Fig S2
Recruitment		
14a	Dates defining the periods of recruitment and follow-up	Results: Enrollment
14b	Why the trial ended or was stopped	N/A
Baseline Data		
15	A table showing baseline demographic and clinical characteristics for each group; Baseline characteristics for the individual and cluster levels as applicable for each group	Tables S2, S3
Numbers Analyzed		
16	For each group, number of participants/clusters (denominator) included in each analysis and whether the analysis was by the original assigned groups	Statistical parameters and estimation strategy (paragraph 2), Tables S6, S7
Outcomes and Estimation		
17a	For each primary and secondary outcome, results for each group, and the estimated effect size and its precision (such as 95% confidence interval); Results at the individual and cluster levels as applicable and a coefficient of intracluster correlation (ICC or k) for each primary outcome	Results: Intervention effects on antibiotic use, Fig 1, Fig2, Tables S8-S11
17b	For binary outcome, presentation of both absolute and relative effect sizes is recommended	Results: Intervention effects on antibiotic use, Tables S8-S11
Ancillary Analyses		
18	Results of any other analyses performed, including subgroup analyses and adjusted analyses, distinguishing pre-specified from exploratory	Results: Subgroup analyses, Secondary and sensitivity analyses, Fig 3, Fig 4, Tables S12-S16
Harms		
19	All important harms or unintended effects in each group	N/A
Discussion		
Limitations		
20	Trial limitations, addressing sources of potential bias, imprecision and, if relevant, multiplicity of analyses	Discussion
Generalizability		
21	Generalizability (external validity, applicability) of the trial findings; Generalizability to clusters and/or individual participants (as relevant)	Discussion
Interpretation		
22	Interpretation consistent with results, balancing benefits and harms, and considering other relevant evidence	Discussion
Other Information		
Registration		
23	Registration number and name of trial registry	Methods: Study design
Protocol		
24	Where the full trial protocol can be accessed, if available	N/A
Funding		
25	Sources of funding and other support (such as supply of drugs), role of funders	Funding

Table S2. Enrollment characteristics by intervention group within the EED substudy and the WASH Benefits parent trial in Bangladesh

	EED substudy				WASH Benefits parent trial			
	C	WSH	N	N+WSH	C	WSH	N	N+WSH
Number of compounds	454	429	450	450	1382	698	699	686
Socio-demographics								
Mother's age in years, mean (SD)	23 (5)	24 (5)	24 (5)	24 (5)	24 (5)	24 (5)	24 (5)	24 (6)
Mother's years of education, mean (SD)	7 (3)	6 (3)	6 (4)	6 (3)	6 (3)	6 (3)	6 (3)	6 (3)
Father's years of education, mean (SD)	5 (4)	5 (4)	5 (4)	5 (4)	5 (4)	5 (4)	5 (4)	5 (4)
Father works in agriculture, % (n)	23% (104)	29% (128)	34% (148)	28% (127)	30% (414)	31% (216)	33% (232)	30% (207)
Number of people in household, mean (SD)	5 (2)	5 (2)	5 (2)	5 (2)	5 (2)	5 (2)	5 (2)	5 (2)
Has electricity, % (n)	60% (269)	62% (278)	62% (269)	61% (272)	57% (784)	61% (426)	59% (409)	60% (412)
Has cement floor, % (n)	17% (75)	12% (55)	11% (50)	12% (53)	10% (145)	11% (77)	10% (67)	10% (72)
Acres of agricultural land owned, mean (SD)	0 (0)	0 (0)	0 (0)	0 (0)	0.15 (0.21)	0.15 (0.23)	0.16 (0.27)	0.14 (0.38)
Drinking water								
Primary water source is shallow tubewell, % (n)	73% (329)	76% (338)	71% (309)	71% (318)	75% (1038)	78% (546)	74% (519)	73% (504)
Stored water observed at home, % (n)	51% (230)	45% (200)	48% (209)	51% (229)	48% (666)	43% (304)	43% (301)	48% (331)
Reported treating water yesterday, % (n)	0% (1)	0% (0)	0% (0)	0% (1)	0% (4)	0% (0)	0% (0)	0% (2)
Distance to water source in minutes, mean (SD)	1 (2)	1 (6)	1 (2)	1 (2)	1 (3)	1 (5)	1 (3)	1 (2)
Sanitation								
Reported daily open defecation								
Adult men, % (n)	4% (19)	7% (30)	9% (38)	9% (38)	7% (97)	8% (54)	9% (59)	7% (50)
Adult women, % (n)	3% (12)	4% (16)	5% (23)	5% (21)	4% (62)	4% (29)	6% (39)	4% (24)
Children 8-15 yrs, % (n)	5% (9)	8% (17)	8% (13)	11% (22)	10% (53)	10% (30)	8% (23)	10% (28)
Children 3-8 yrs, % (n)	30% (64)	37% (90)	40% (90)	37% (92)	38% (267)	38% (137)	39% (129)	37% (134)
Children <3 yrs, % (n)	72% (71)	75% (73)	80% (68)	88% (79)	82% (245)	79% (123)	85% (128)	88% (123)
Latrine								
Owned by household, % (n)	60% (271)	55% (244)	54% (234)	51% (230)	54% (750)	53% (373)	54% (377)	53% (367)
Has concrete slab, % (n)	97% (426)	93% (401)	93% (382)	94% (399)	95% (1251)	93% (620)	94% (620)	94% (621)
Has functional water seal, % (n)	38% (157)	26% (95)	32% (114)	31% (111)	31% (358)	26% (152)	31% (183)	27% (155)
Owns child potty	8% (37)	4% (20)	6% (27)	5% (21)	4% (61)	4% (27)	5% (36)	4% (30)
Human feces observed								
In house, % (n)	6% (25)	8% (36)	9% (41)	8% (36)	8% (114)	7% (48)	8% (58)	7% (49)
In child's play area, % (n)	1% (5)	1% (4)	2% (7)	1% (6)	2% (21)	1% (7)	1% (8)	1% (7)
Handwashing								
Within 6 steps of latrine								
Has water, % (n)	21% (84)	13% (51)	10% (38)	13% (54)	14% (178)	10% (67)	10% (62)	11% (72)
Has soap, % (n)	7% (88)	7% (42)	5% (32)	6% (36)	11% (45)	8% (32)	6% (23)	7% (27)
Within 6 steps of kitchen								
Has water, % (n)	12% (48)	10% (4)	11% (43)	10% (42)	9% (118)	9% (61)	9% (61)	9% (60)
Has soap, % (n)	4% (18)	3% (11)	5% (19)	3% (14)	3% (33)	2% (15)	4% (23)	3% (18)
Nutrition								
Household is food secure, % (n)	71% (485)	74% (331)	67% (298)	71% (308)	67% (932)	69% (482)	69% (479)	71% (485)

Table S3. Enrollment characteristics by intervention group within the EED substudy and the WASH Benefits parent trial in Kenya

	EED substudy				WASH Benefits parent trial			
	C	WSH	N	N+WSH	C	WSH	N	N+WSH
Number of compounds	511	456	434	448	1919	912	843	921
Socio-demographics								
Mother's age in years, mean (SD)	26 (6)	26 (6)	26 (6)	26 (6)	26 (6)	26 (6)	26 (6)	26 (6)
Mother completed at least primary education, % (n)	45% (227)	44% (200)	50% (215)	49% (217)	48% (916)	47% (430)	49% (409)	48% (438)
Father completed at least primary education, % (n)	59% (279)	59% (252)	61% (248)	62% (258)	62% (1098)	61% (521)	64% (491)	62% (526)
Father works in agriculture, % (n)	47% (231)	45% (197)	49% (201)	44% (187)	41% (749)	43% (374)	43% (343)	43% (372)
Number of people in compound, mean (SD)	9 (6)	9 (5)	9 (6)	9 (6)	8 (5)	8 (5)	8 (7)	8 (5)
Has electricity, % (n)	4% (22)	8% (35)	7% (30)	6% (25)	6% (122)	7% (64)	7% (58)	7% (67)
Has cement floor, % (n)	5% (25)	5% (25)	5% (20)	4% (19)	6% (107)	5% (50)	6% (48)	6% (56)
Has iron roof, % (n)	60% (305)	59% (271)	66% (285)	62% (279)	68% (1302)	63% (574)	69% (581)	67% (615)
Drinking water								
Primary water source is improved, % (n)	79% (402)	71% (324)	71% (308)	80% (355)	76% (1446)	69% (624)	72% (603)	76% (697)
Reported treating currently stored water, % (n)	10% (43)	13% (46)	8% (27)	13% (46)	13% (196)	13% (97)	12% (79)	14% (106)
Distance to water source in minutes, mean (SD)	10 (10)	10 (11)	10 (11)	10 (12)	11 (12)	11 (13)	11 (12)	11 (12)
Sanitation								
Children <3 yrs daily defecate in the open, % (n)	77% (230)	76% (195)	78% (188)	79% (187)	78% (789)	77% (394)	79% (363)	78% (388)
Household owns latrine, % (n)	81% (415)	86% (390)	85% (370)	86% (384)	82% (1561)	83% (754)	83% (701)	83% (764)
Access to improved latrine, % (n)	17% (82)	17% (75)	16% (66)	15% (65)	17% (309)	18% (153)	15% (119)	16% (143)
Human feces observed in compound, % (n)	7% (36)	9% (39)	9% (40)	9% (42)	9% (163)	8% (73)	9% (73)	9% (87)
Handwashing location								
Has water within 2 m, % (n)	24% (121)	28% (127)	27% (117)	27% (121)	25% (487)	28% (251)	27% (228)	27% (249)
Has soap within 2 m, % (n)	8% (40)	13% (61)	11% (46)	10% (46)	9% (164)	13% (115)	11% (90)	9% (87)
Nutrition								
Moderate to severe household hunger, % (n)	11% (55)	9% (41)	12% (54)	11% (47)	11% (203)	11% (101)	12% (98)	11% (104)

Table S4. Enrollment characteristics of children in the EED substudy included in follow-up vs. lost to follow-up in Bangladesh

	Included	Missing from follow-up at age 14 months	Missing from follow-up at age 28 months
Number of compounds	1778	284	273
Socio-demographics			
Mother's age in years, mean (SD)	24 (5)	24 (5)	24 (5)
Mother's years of education, mean (SD)	6 (3)	6 (3)	6 (3)
Father's years of education, mean (SD)	5 (4)	5 (4)	5 (4)
Father works in agriculture, % (n)	29% (507)	24% (69)	26% (70)
Number of people in household, mean (SD)	5 (2)	5 (2)	5 (2)
Has electricity, % (n)	61% (1088)	60% (169)	56% (153)
Has cement floor, % (n)	13% (233)	14% (40)	15% (42)
Acres of agricultural land owned, mean (SD)	0 (0)	0 (0)	0 (0)
Drinking water			
Primary water source is shallow tubewell, % (n)	73% (1294)	70% (200)	70% (191)
Stored water observed at home, % (n)	49% (868)	51% (144)	52% (142)
Reported treating water yesterday, % (n)	0% (2)	0% (0)	0% (0)
Distance to water source in minutes, mean (SD)	1 (3)	1 (2)	1 (1)
Sanitation			
Reported daily open defecation			
Adult men, % (n)	7% (125)	8% (21)	7% (18)
Adult women, % (n)	4% (72)	5% (14)	3% (9)
Children 8-15 yrs, % (n)	8% (61)	11% (12)	14% (15)
Children 3-8 yrs, % (n)	36% (337)	39% (57)	37% (51)
Children <3 yrs, % (n)	79% (291)	83% (45)	81% (38)
Latrine			
Owned by household, % (n)	55% (979)	52% (148)	56% (154)
Has concrete slab, % (n)	94% (1608)	96% (258)	95% (249)
Has functional water seal, % (n)	32% (477)	31% (73)	36% (84)
Owns child potty	6% (105)	7% (20)	7% (20)
Human feces observed			
In house, % (n)	8% (138)	4% (12)	7% (18)
In child's play area, % (n)	1% (22)	1% (3)	2% (6)
Handwashing			
Within 6 steps of latrine			
Has water, % (n)	14% (227)	16% (42)	16% (40)
Has soap, % (n)	8% (127)	10% (27)	11% (27)
Within 6 steps of kitchen			
Has water, % (n)	11% (173)	12% (30)	9% (21)
Has soap, % (n)	4% (62)	5% (12)	2% (5)
Nutrition			
Household is food secure, % (n)	71% (1254)	68% (192)	68% (187)

Table S5. Enrollment characteristics of children in the EED substudy included in follow-up vs. lost to follow-up in Kenya

	Included	Missing from follow-up at age 17 months	Missing from follow-up at age 22 months
Number of compounds	1849	462	556
Socio-demographics			
Mother's age in years, mean (SD)	26 (6)	25 (6)	25 (6)
Mother completed at least primary education, % (n)	47% (859)	49% (225)	49% (271)
Father completed at least primary education, % (n)	60% (1037)	62% (267)	61% (310)
Father works in agriculture, % (n)	46% (816)	42% (181)	43% (223)
Number of people in compound, mean (SD)	9 (6)	8 (6)	8 (5)
Has electricity, % (n)	6% (112)	7% (31)	6% (33)
Has cement floor, % (n)	5% (89)	6% (27)	5% (29)
Has iron roof, % (n)	62% (1140)	60% (279)	61% (341)
Drinking water			
Primary water source is improved, % (n)	75% (1389)	74% (340)	76% (424)
Reported treating currently stored water, % (n)	11% (162)	11% (39)	12% (54)
Distance to water source in minutes, mean (SD)	10 (11)	10 (9)	10 (11)
Sanitation			
Children <3 yrs daily defecate in the open, % (n)	78% (800)	77% (178)	71% (205)
Household owns latrine, % (n)	84% (1559)	81% (375)	82% (454)
Access to improved latrine, % (n)	16% (288)	19% (83)	18% (93)
Human feces observed in compound, % (n)	9% (157)	9% (40)	10% (55)
Handwashing location			
Has water within 2 m, % (n)	26% (486)	26% (122)	27% (148)
Has soap within 2 m, % (n)	10% (193)	11% (52)	10% (58)
Nutrition			
Moderate to severe household hunger, % (n)	11% (197)	14% (65)	11% (59)

Table S6. Caregiver-reported antibiotic use in last 3 months by young children in subgroups of age and sex, Bangladesh

Arm	All observations		Mean child age						Sex			
	N	% (n)	3 months		14 months		28 months		Female		Male	
			N	% (n)	N	% (n)	N	% (n)	N	% (n)	N	% (n)
Used antibiotics ≥ 1 time												
C	1005	63.2 (635)	270	54.1 (146)	377	75.1 (283)	358	57.5 (206)	515	62.3 (321)	490	64.1 (314)
WSH	1077	57.0 (614)	285	45.6 (130)	395	70.6 (279)	397	51.6 (205)	510	54.7 (279)	567	59.1 (335)
N	1018	54.1 (551)	269	41.3 (111)	377	68.2 (257)	372	49.2 (183)	492	49.6 (244)	526	58.4 (307)
N+WSH	1058	54.2 (573)	278	36.0 (100)	379	71.5 (271)	401	50.4 (202)	565	49.7 (281)	493	59.2 (292)
Used antibiotics ≥ 2 times												
C	1005	24.7 (248)	270	24.8 (67)	377	34.8 (131)	358	14.0 (50)	515	22.1 (114)	490	27.4 (134)
WSH	1077	18.2 (196)	285	14.0 (40)	395	25.6 (101)	397	13.9 (55)	510	15.9 (81)	567	20.3 (115)
N	1018	16.4 (167)	269	12.6 (34)	377	24.4 (92)	372	11.0 (41)	492	15.7 (77)	526	17.1 (90)
N+WSH	1058	16.2 (171)	278	11.9 (33)	379	26.9 (102)	401	9.0 (36)	565	14.5 (82)	493	18.1 (89)
Arm	N	Mean (SD)	N	Mean (SD)	N	Mean (SD)	N	Mean (SD)	N	Mean (SD)	N	Mean (SD)
Episodes of antibiotic use												
C	1005	0.98 (0.98)	270	0.90 (1.06)	377	1.23 (0.99)	358	0.77 (0.84)	515	0.92 (0.94)	490	1.03 (1.02)
WSH	1077	0.81 (0.88)	285	0.65 (0.87)	395	1.05 (0.93)	397	0.68 (0.78)	510	0.74 (0.81)	567	0.87 (0.93)
N	1018	0.77 (0.90)	269	0.58 (0.82)	377	1.04 (0.99)	372	0.64 (0.79)	492	0.70 (0.86)	526	0.84 (0.93)
N+WSH	1058	0.77 (0.89)	278	0.56 (0.93)	379	1.08 (0.95)	401	0.61 (0.71)	565	0.70 (0.87)	493	0.84 (0.91)
Total days of antibiotic use												
C	995	5.06 (5.81)	265	4.77 (6.36)	373	6.27 (5.91)	357	4.01 (4.99)	511	4.77 (5.58)	484	5.37 (6.03)
WSH	1075	4.12 (5.05)	283	3.54 (5.18)	395	5.13 (5.20)	397	3.52 (4.64)	509	3.82 (4.83)	566	4.39 (5.22)
N	1016	4.02 (5.30)	268	3.00 (4.84)	376	5.34 (5.74)	372	3.41 (4.86)	492	3.67 (5.15)	524	4.34 (5.41)
N+WSH	1055	4.05 (5.39)	276	3.25 (6.01)	379	5.70 (5.88)	400	3.04 (3.86)	564	3.65 (5.14)	491	4.52 (5.63)

C: Control; WSH: Water, sanitation, hygiene; N: Nutrition, N+WSH: Nutrition+ water, sanitation, hygiene; SD: Standard deviation.

Table S7. Caregiver-reported antibiotic use in last 3 months by young children in subgroups of age and sex, Kenya

	All observations		Mean child age						Sex			
			6 months		17 months		22 months		Female		Male	
Arm	N	% (n)	N	% (n)	N	% (n)	N	% (n)	N	% (n)	N	% (n)
Used antibiotics ≥1 time												
C	1143	52.6 (601)	360	52.5 (189)	394	53.3 (210)	389	51.9 (202)	571	53.2 (304)	572	51.9 (297)
WSH	1043	50.3 (525)	366	49.7 (182)	345	53.9 (186)	332	47.3 (157)	568	49.5 (281)	475	51.4 (244)
N	1048	50.4 (528)	347	49.3 (171)	358	54.8 (196)	343	46.9 (161)	495	51.1 (253)	553	49.7 (275)
N+WSH	1046	53.1 (555)	365	51.8 (189)	352	55.7 (196)	329	51.7 (170)	574	52.8 (303)	472	53.4 (252)
Used antibiotics ≥2 times												
C	1143	13.3 (152)	360	17.2 (62)	394	11.7 (46)	389	11.3 (44)	571	12.8 (73)	572	13.8 (79)
WSH	1043	13.7 (143)	366	14.2 (52)	345	13.9 (48)	332	13.0 (43)	568	12.7 (72)	475	15.0 (71)
N	1048	11.6 (122)	347	14.7 (51)	358	12.0 (43)	343	8.2 (28)	495	12.7 (63)	553	10.7 (59)
N+WSH	1046	13.0 (136)	365	14.8 (54)	352	13.6 (48)	329	10.3 (34)	574	12.4 (71)	472	13.8 (65)
Arm	N	Mean (SD)	N	Mean (SD)	N	Mean (SD)	N	Mean (SD)	N	Mean (SD)	N	Mean (SD)
Episodes of antibiotic use												
C	1143	0.68 (0.75)	360	0.74 (0.85)	394	0.65 (0.69)	389	0.64 (0.69)	571	0.67 (0.72)	572	0.68 (0.77)
WSH	1043	0.66 (0.77)	366	0.68 (0.83)	345	0.69 (0.74)	332	0.61 (0.74)	568	0.64 (0.76)	475	0.69 (0.78)
N	1048	0.65 (0.78)	347	0.70 (0.93)	358	0.68 (0.72)	343	0.55 (0.65)	495	0.66 (0.77)	553	0.63 (0.79)
N+WSH	1046	0.69 (0.78)	365	0.71 (0.85)	352	0.72 (0.78)	329	0.63 (0.70)	574	0.68 (0.77)	472	0.71 (0.80)
Total days of antibiotic use												
C	1142	3.95 (12.1)	359	5.61 (18.8)	394	3.42 (9.11)	389	2.95 (3.90)	571	4.10 (12.7)	571	3.80 (11.5)
WSH	1043	4.14 (13.0)	366	4.06 (10.9)	345	5.37 (19.2)	332	2.95 (4.12)	568	4.53 (15.7)	475	3.68 (8.72)
N	1047	3.36 (8.45)	347	2.90 (4.69)	357	4.15 (12.0)	343	2.99 (6.67)	494	3.59 (9.60)	553	3.15 (7.27)
N+WSH	1046	3.71 (9.02)	365	3.93 (9.96)	352	3.56 (9.51)	329	3.61 (7.22)	574	3.60 (8.28)	472	3.84 (9.86)

C: Control; WSH: Water, sanitation, hygiene; N: Nutrition, N+WSH: Nutrition+ water, sanitation, hygiene; SD: Standard deviation.

Table S8. WSH, nutrition and N+WSH interventions vs. controls on antibiotic use within last 3 months among children in Bangladesh

			Relative effects				Absolute effects			
			Unadjusted		Adjusted		Unadjusted		Adjusted	
Arm	n	% (n)	PR (95% CI)	p-value	PR (95% CI)	p-value	PD (95% CI)	p-value	PD (95% CI)	p-value
Used antibiotics ≥1 time										
C	1005	63.2 (635)	ref	--	--	--	--	--	--	--
WSH	1077	57.0 (614)	0.90 (0.82, 0.99)	0.03	0.90 (0.82, 0.99)	0.03	-0.06 (-0.12, -0.01)	0.03	-0.07 (-0.12, 0.03)	0.03
N	1018	54.1 (551)	0.86 (0.78, 0.94)	<0.001	0.84 (0.76, 0.92)	<0.001	-0.09 (-0.14, -0.04)	<0.001	-0.10 (-0.16, -0.05)	<0.001
N+WSH	1058	54.2 (573)	0.86 (0.79, 0.93)	<0.001	0.86 (0.78, 0.93)	<0.001	-0.09 (-0.14, -0.04)	<0.001	-0.10 (-0.15, -0.04)	<0.001
Used antibiotics ≥2 times										
C	1005	24.7 (248)	ref	--	--	--	--	--	--	--
WSH	1077	18.2 (196)	0.74 (0.63, 0.87)	<0.001	0.73 (0.63, 0.85)	<0.001	-0.06 (-0.10, -0.03)	<0.001	-0.07 (-0.10, -0.04)	<0.001
N	1018	16.4 (167)	0.66 (0.56, 0.79)	<0.001	0.64 (0.53, 0.78)	<0.001	-0.08 (-0.12, -0.05)	<0.001	-0.09 (-0.13, -0.05)	<0.001
N+WSH	1058	16.2 (171)	0.65 (0.55, 0.78)	<0.001	0.66 (0.55, 0.79)	<0.001	-0.09 (-0.12, -0.05)	<0.001	-0.08 (-0.12, -0.05)	<0.001
Arm	N	Mean (SD)	CR (95% CI)	p-value	CR (95% CI)	p-value	CD (95% CI)	p-value	CD (95% CI)	p-value
Episodes of antibiotic use										
C	1005	0.98 (0.98)	ref	--	--	--	--	--	--	--
WSH	1077	0.81 (0.88)	0.83 (0.75, 0.92)	<0.001	0.82 (0.74, 0.90)	<0.001	-0.17 (-0.26, -0.07)	<0.001	-0.18 (-0.27, -0.09)	<0.001
N	1018	0.77 (0.90)	0.79 (0.72, 0.88)	<0.001	0.77 (0.69, 0.86)	<0.001	-0.20 (-0.29, -0.11)	<0.001	-0.22 (-0.32, -0.13)	<0.001
N+WSH	1058	0.77 (0.89)	0.79 (0.71, 0.87)	<0.001	0.78 (0.71, 0.87)	<0.001	-0.21 (-0.29, -0.12)	<0.001	-0.21 (-0.30, -0.12)	<0.001
Total days of antibiotic use										
C	995	5.06 (5.81)	ref	--	--	--	--	--	--	--
WSH	1075	4.11 (5.05)	0.81 (0.74, 0.90)	<0.001	0.81 (0.73, 0.89)	<0.001	-0.94 (-1.41, -0.48)	<0.001	-1.02 (-1.49, -0.54)	<0.001
N	1016	4.02 (5.30)	0.79 (0.71, 0.89)	<0.001	0.75 (0.67, 0.85)	<0.001	-1.05 (-1.54, -0.55)	<0.001	-1.19 (-1.74, -0.65)	<0.001
N+WSH	1055	4.05 (5.39)	0.80 (0.72, 0.89)	<0.001	0.80 (0.71, 0.89)	<0.001	-1.01 (-1.51, -0.51)	<0.001	-1.06 (-1.60, -0.53)	<0.001

PR: Prevalence ratio (for binary outcomes), CR: Count ratio (for count outcomes); PD: Prevalence difference (for binary outcomes); CD: Count difference (for count outcomes); CI: Confidence interval; C: Control; WSH: Water, sanitation, hygiene; N: Nutrition, N+WSH: Nutrition+ water, sanitation, hygiene.

Table S9. Combined N+WSH intervention vs. WSH and nutrition interventions on antibiotic use within last 3 months among children in Bangladesh

Arm	Relative effects				Absolute effects			
	Unadjusted		Adjusted		Unadjusted		Adjusted	
	PR (95% CI)	p-value	PR (95% CI)	p-value	PD (95% CI)	p-value	PD (95% CI)	p-value
Used antibiotics ≥1 time								
N+WSH vs. WSH	0.95 (0.87, 1.03)	0.22	0.95 (0.88, 1.04)	0.26	-0.03 (-0.07, 0.02)	0.22	-0.02 (-0.07, 0.02)	0.35
N+WSH vs. N	1.00 (0.91, 1.10)	0.99	1.01 (0.92, 1.12)	0.77	0.00 (-0.05, 0.05)	0.99	0.01 (-0.05, 0.06)	0.77
Used antibiotics ≥2 times								
N+WSH vs. WSH	0.89 (0.74, 1.06)	0.19	0.91 (0.76, 1.09)	0.32	-0.02 (-0.05, 0.01)	0.19	-0.01 (-0.04, 0.01)	0.40
N+WSH vs. N	0.99 (0.83, 1.17)	0.87	1.05 (0.88, 1.25)	0.62	-0.00 (-0.03, 0.03)	0.87	0.01 (-0.02, 0.04)	0.59
Episodes of antibiotic use								
N+WSH vs. WSH	0.95 (0.85, 1.06)	0.35	0.98 (0.88, 1.09)	0.68	-0.04 (-0.13, 0.04)	0.35	-0.02 (-0.10, 0.07)	0.67
N+WSH vs. N	0.99 (0.89, 1.10)	0.87	1.02 (0.91, 1.14)	0.76	-0.01 (-0.09, 0.08)	0.87	0.01 (-0.07, 0.10)	0.75
Total days of antibiotic use								
N+WSH vs. WSH	0.98 (0.87, 1.11)	0.79	0.99 (0.87, 1.11)	0.83	-0.07 (-0.55, 0.41)	0.79	0.04 (-0.45, 0.53)	0.87
N+WSH vs. N	1.01 (0.90, 1.13)	0.88	1.05 (0.93, 1.18)	0.41	0.03 (-0.42, 0.49)	0.88	0.17 (-0.30, 0.63)	0.48

PR: Prevalence ratio (for binary outcomes), CR: Count ratio (for count outcomes); PD: Prevalence difference (for binary outcomes); CD: Count difference (for count outcomes); CI: Confidence interval; WSH: Water, sanitation, hygiene; N: Nutrition, N+WSH: Nutrition+ water, sanitation, hygiene.

Table S10. WSH, nutrition and N+WSH interventions vs. controls on antibiotic use within last 3 months among children in Kenya

			Relative effects				Absolute effects			
			Unadjusted		Adjusted		Unadjusted		Adjusted	
Arm	N	% (n)	PR (95% CI)	p-value	PR (95% CI)	p-value	PD (95% CI)	p-value	PD (95% CI)	p-value
Used antibiotics ≥1 time										
C	1143	52.6 (601)	ref	--	--	--	--	--	--	--
WSH	1043	50.3 (525)	0.96 (0.86, 1.06)	0.41	0.97 (0.88, 1.07)	0.55	-0.02 (-0.08, 0.03)	0.41	-0.02 (-0.07, 0.04)	0.53
N	1048	50.4 (528)	0.96 (0.88, 1.04)	0.33	0.95 (0.87, 1.03)	0.19	-0.02 (-0.07, 0.02)	0.33	-0.03 (-0.07, 0.02)	0.22
N+WSH	1046	53.1 (555)	1.01 (0.92, 1.10)	0.84	1.00 (0.91, 1.09)	0.99	0.00 (-0.04, 0.05)	0.84	0.00 (-0.05, 0.05)	0.97
Used antibiotics ≥2 times										
C	1143	13.3 (152)	ref	--	--	--	--	--	--	--
WSH	1043	13.7 (143)	1.03 (0.80, 1.32)	0.81	1.03 (0.79, 1.35)	0.81	0.00 (-0.03, 0.04)	0.81	0.00 (-0.03, 0.04)	0.80
N	1048	11.6 (122)	0.88 (0.68, 1.12)	0.30	0.88 (0.68, 1.14)	0.33	-0.02 (-0.05, 0.01)	0.30	-0.02 (-0.05, 0.02)	0.33
N+WSH	1046	13.0 (136)	0.98 (0.78, 1.22)	0.84	0.94 (0.74, 1.19)	0.61	-0.00 (-0.03, 0.03)	0.84	-0.01 (-0.04, 0.02)	0.61
Arm	N	Mean (SD)	CR (95% CI)	p-value	CR (95% CI)	p-value	CD (95% CI)	p-value	CD (95% CI)	p-value
Episodes of antibiotic use										
C	1143	0.68 (0.75)	ref	--	--	--	--	--	--	--
WSH	1043	0.66 (0.77)	0.98 (0.87, 1.10)	0.77	0.99 (0.89, 1.12)	0.92	-0.01 (-0.09, 0.07)	0.77	-0.00 (-0.08, 0.07)	0.93
N	1048	0.65 (0.78)	0.96 (0.85, 1.07)	0.45	0.96 (0.86, 1.07)	0.41	-0.03 (-0.10, 0.05)	0.44	-0.03 (-0.10, 0.04)	0.40
N+WSH	1046	0.69 (0.78)	1.02 (0.92, 1.14)	0.66	1.00 (0.90, 1.12)	0.90	0.02 (-0.06, 0.09)	0.67	0.00 (-0.07, 0.08)	0.90
Total days of antibiotic use										
C	1142	3.95 (12.1)	ref	--	--	--	--	--	--	--
WSH	1043	4.14 (13.0)	1.05 (0.71, 1.54)	0.81	1.00 (0.73, 1.37)	0.99	0.19 (-1.37, 1.76)	0.81	0.25 (-1.37, 1.88)	0.76
N	1047	3.36 (8.45)	0.85 (0.65, 1.12)	0.24	0.83 (0.63, 1.09)	0.17	-0.59 (-1.61, 0.43)	0.25	-0.68 (-1.68, 0.32)	0.18
N+WSH	1046	3.71 (9.02)	0.94 (0.76, 1.17)	0.57	0.95 (0.77, 1.16)	0.60	-0.24 (-1.08, 0.60)	0.58	-0.29 (-1.16, 0.58)	0.51

PR: Prevalence ratio (for binary outcomes), CR: Count ratio (for count outcomes); PD: Prevalence difference (for binary outcomes); CD: Count difference (for count outcomes); CI: Confidence interval; C: Control; WSH: Water, sanitation, hygiene; N: Nutrition, N+WSH: Nutrition+ water, sanitation, hygiene.

Table S11. N+WSH intervention vs. WSH and nutrition interventions on antibiotic use within last 3 months among children in Kenya

Arm	Relative effects				Absolute effects			
	Unadjusted		Adjusted		Unadjusted		Adjusted	
	PR (95% CI)	p-value	PR (95% CI)	p-value	PD (95% CI)	p-value	PD (95% CI)	p-value
Used antibiotics ≥1 time								
N+WSH vs. WSH	1.05 (0.95, 1.17)	0.33	1.04 (0.93, 1.15)	0.50	0.03 (-0.03, 0.08)	0.33	0.02 (-0.04, 0.07)	0.54
N+WSH vs. N	1.05 (0.95, 1.17)	0.32	1.05 (0.95, 1.16)	0.36	0.03 (-0.03, 0.08)	0.32	0.03 (-0.03, 0.08)	0.35
Used antibiotics ≥2 times								
N+WSH vs. WSH	0.95 (0.71, 1.27)	0.72	0.95 (0.70, 1.28)	0.72	-0.01 (-0.05, 0.03)	0.73	-0.01 (-0.05, 0.03)	0.73
N+WSH vs. N	1.12 (0.82, 1.52)	0.48	1.12 (0.81, 1.54)	0.49	0.01 (-0.02, 0.05)	0.49	0.01 (-0.03, 0.05)	0.51
Episodes of antibiotic use								
N+WSH vs. WSH	1.04 (0.91, 1.19)	0.56	1.02 (0.89, 1.17)	0.72	0.03 (-0.06, 0.12)	0.56	0.02 (-0.08, 0.11)	0.73
N+WSH vs. N	1.07 (0.93, 1.23)	0.34	1.08 (0.93, 1.24)	0.31	0.05 (-0.05, 0.14)	0.34	0.05 (-0.05, 0.14)	0.33
Total days of antibiotic use								
N+WSH vs. WSH	0.90 (0.64, 1.26)	0.52	0.95 (0.72, 1.25)	0.70	-0.43 (-1.81, 0.94)	0.53	-0.48 (-1.86, 0.89)	0.49
N+WSH vs. N	1.10 (0.87, 1.40)	0.41	1.19 (0.95, 1.49)	0.13	0.35 (-0.48, 1.18)	0.41	0.50 (-0.33, 1.33)	0.24

PR: Prevalence ratio (for binary outcomes), CR: Count ratio (for count outcomes); PD: Prevalence difference (for binary outcomes); CD: Count difference (for count outcomes); CI: Confidence interval; WSH: Water, sanitation, hygiene; N: Nutrition, N+WSH: Nutrition+ water, sanitation, hygiene.

Table S12. Multiplicative interaction by age and sex for unadjusted intervention effects on antibiotic use within last 3 months, Bangladesh

	WSH vs. C		N vs. C		N+WSH vs. C		N+WSH vs. WSH		N+WSH vs. N	
	PR (95% CI)	p-value ^a	PR (95% CI)	p-value ^a	PR (95% CI)	p-value ^a	PR (95% CI)	p-value ^a	PR (95% CI)	p-value ^a
Used antibiotics ≥1 time										
All obs	0.90 (0.82, 0.99)		0.86 (0.78, 0.94)		0.86 (0.79, 0.93)		0.95 (0.87, 1.03)		1.00 (0.91, 1.10)	
Age 3 mo	0.84 (0.70, 1.02)	--	0.76 (0.63, 0.93)	--	0.67 (0.54, 0.82)	--	0.79 (0.62, 1.00)	--	0.87 (0.67, 1.13)	--
Age 14 mo	0.94 (0.85, 1.04)	0.26	0.91 (0.82, 1.00)	0.10 *	0.95 (0.86, 1.05)	<0.001 *	1.01 (0.92, 1.11)	0.06 *	1.05 (0.94, 1.17)	0.15 *
Age 28 mo	0.90 (0.78, 1.03)	0.57	0.85 (0.73, 1.00)	0.39	0.88 (0.76, 1.00)	0.04 *	0.98 (0.87, 1.10)	0.09 *	1.02 (0.89, 1.18)	0.30
Female	0.88 (0.77, 1.00)	--	0.80 (0.70, 0.90)	--	0.80 (0.71, 0.90)	--	0.91 (0.80, 1.03)	--	1.00 (0.87, 1.15)	--
Male	0.92 (0.83, 1.03)	0.49	0.91 (0.81, 1.02)	0.09 *	0.92 (0.83, 1.03)	0.05 *	1.00 (0.90, 1.11)	0.24	1.01 (0.91, 1.13)	0.88
Used antibiotics ≥2 times										
All obs	0.74 (0.63, 0.87)		0.66 (0.56, 0.79)		0.65 (0.55, 0.78)		0.89 (0.74, 1.06)		0.99 (0.83, 1.17)	
Age 3 mo	0.57 (0.37, 0.86)	--	0.51 (0.33, 0.78)	--	0.48 (0.33, 0.69)	--	0.85 (0.54, 1.32)	--	0.94 (0.61, 1.44)	--
Age 14 mo	0.74 (0.60, 0.90)	0.26	0.70 (0.57, 0.86)	0.18 *	0.77 (0.64, 0.94)	0.02 *	1.05 (0.84, 1.31)	0.40	1.10 (0.90, 1.36)	0.52
Age 28 mo	0.99 (0.75, 1.32)	0.04 *	0.79 (0.55, 1.13)	0.12 *	0.64 (0.42, 0.98)	0.29	0.65 (0.45, 0.93)	0.38	0.81 (0.53, 1.26)	0.64
Female	0.72 (0.55, 0.93)	--	0.71 (0.56, 0.90)	--	0.66 (0.52, 0.83)	--	0.91 (0.69, 1.20)	--	0.93 (0.70, 1.22)	--
Male	0.74 (0.60, 0.91)	0.85	0.63 (0.50, 0.79)	0.47	0.66 (0.51, 0.85)	0.97	0.89 (0.70, 1.13)	0.89	1.06 (0.81, 1.37)	0.54
	CR (95% CI)	p-value ^a	CR (95% CI)	p-value ^a	CR (95% CI)	p-value ^a	CR (95% CI)	p-value ^a	CR (95% CI)	p-value ^a
Episodes of antibiotic use										
All obs	0.83 (0.75, 0.92)		0.79 (0.72, 0.88)		0.79 (0.71, 0.87)		0.95 (0.85, 1.06)		0.99 (0.89, 1.10)	
Age 3 mo	0.72 (0.57, 0.92)	--	0.65 (0.50, 0.84)	--	0.62 (0.49, 0.80)	--	0.86 (0.67, 1.11)	--	0.96 (0.74, 1.25)	--
Age 14 mo	0.85 (0.75, 0.96)	0.19 *	0.85 (0.75, 0.96)	0.05 *	0.88 (0.79, 0.99)	0.01 *	1.03 (0.90, 1.18)	0.18 *	1.04 (0.91, 1.19)	0.59
Age 28 mo	0.89 (0.77, 1.03)	0.14 *	0.84 (0.71, 1.00)	0.11 *	0.80 (0.68, 0.94)	0.10 *	0.90 (0.78, 1.04)	0.75	0.95 (0.81, 1.13)	0.97
Female	0.80 (0.69, 0.93)	--	0.76 (0.66, 0.88)	--	0.76 (0.67, 0.86)	--	0.94 (0.81, 1.09)	--	1.00 (0.86, 1.16)	--
Male	0.84 (0.73, 0.97)	0.65	0.82 (0.71, 0.94)	0.48	0.82 (0.71, 0.95)	0.43	0.97 (0.84, 1.13)	0.75	1.00 (0.87, 1.16)	0.94
Total days of antibiotic use										
All obs	0.81 (0.74, 0.90)		0.79 (0.71, 0.89)		0.80 (0.72, 0.89)		0.98 (0.87, 1.11)		1.01 (0.90, 1.13)	
Age 3 mo	0.74 (0.58, 0.94)	--	0.63 (0.48, 0.83)	--	0.68 (0.52, 0.89)	--	0.92 (0.69, 1.22)	--	1.08 (0.81, 1.46)	--
Age 14 mo	0.82 (0.72, 0.93)	0.44	0.85 (0.75, 0.97)	0.05 *	0.91 (0.79, 1.05)	0.04 *	1.11 (0.96, 1.29)	0.19 *	1.07 (0.93, 1.23)	0.93
Age 28 mo	0.88 (0.74, 1.05)	0.30	0.85 (0.70, 1.03)	0.09 *	0.76 (0.64, 0.90)	0.54 *	0.86 (0.73, 1.01)	0.70	0.89 (0.74, 1.07)	0.29
Female	0.80 (0.67, 0.95)	--	0.77 (0.64, 0.92)	--	0.76 (0.66, 0.89)	--	0.95 (0.82, 1.12)	--	0.99 (0.84, 1.18)	--
Male	0.82 (0.72, 0.93)	0.87	0.81 (0.70, 0.94)	0.68	0.84 (0.72, 0.98)	0.36	1.03 (0.87, 1.22)	0.52	1.04 (0.88, 1.23)	0.72

PR: Prevalence ratio (for binary outcomes), CR: Count ratio (for count outcomes); CI: Confidence interval; C: Control; WSH: Water, sanitation, hygiene; N: Nutrition, N+WSH: Nutrition+ water, sanitation, hygiene.

Bolded values refer to estimates with a p-value ≤0.05.

^a p-value for interaction term between study arm and subgroup variable.

* Interaction p-values <0.20 indicate significant effect modification.

Table S13. Additive interaction by age and sex for unadjusted intervention effects on antibiotic use within last 3 months, Bangladesh

	WSH vs. C		N vs. C		N+WSH vs. C		N+WSH vs. WSH		N+WSH vs. N	
	PD (95% CI)	p-value ^a	PD (95% CI)	p-value ^a	PD (95% CI)	p-value ^a	PD (95% CI)	p-value ^a	PD (95% CI)	p-value ^a
Used antibiotics ≥1 time										
All obs	-0.06 (-0.12, -0.01)		-0.09 (-0.14, -0.04)		-0.09 (-0.14, -0.04)		-0.03 (-0.07, 0.02)		0.00 (-0.05, 0.05)	
Age 3 mo	-0.08 (-0.17, 0.01)	--	-0.13 (-0.22, -0.04)	--	-0.18 (-0.27, -0.09)	--	-0.10 (-0.19, 0.00)	--	-0.05 (-0.15, 0.05)	--
Age 14 mo	-0.04 (-0.12, 0.03)	0.43	-0.07 (-0.14, 0.00)	0.28	-0.04 (-0.11, 0.04)	0.01 *	0.01 (-0.06, 0.08)	0.09 *	0.03 (-0.04, 0.11)	0.12 *
Age 28 mo	-0.06 (-0.14, 0.02)	0.64	-0.08 (-0.16, -0.00)	0.49	-0.07 (-0.15, 0.00)	0.08 *	-0.01 (-0.07, 0.04)	0.12 *	0.01 (-0.06, 0.08)	0.32
Female	-0.08 (-0.15, -0.00)	--	-0.13 (-0.19, -0.07)	--	-0.13 (-0.19, -0.06)	--	-0.05 (-0.12, 0.02)	--	0.00 (-0.07, 0.07)	--
Male	-0.05 (-0.12, 0.02)	0.54	-0.06 (-0.13, 0.01)	0.12 *	-0.05 (-0.12, 0.02)	0.06 *	0.00 (-0.06, 0.06)	0.27	0.01 (-0.06, 0.07)	0.87
Used antibiotics ≥2 times										
All obs	-0.06 (-0.10, -0.03)		-0.08 (-0.12, -0.05)		-0.09 (-0.12, -0.05)		-0.02 (-0.05, 0.01)		0.00 (-0.03, 0.03)	
Age 3 mo	-0.11 (-0.18, -0.03)	--	-0.12 (-0.19, -0.05)	--	-0.13 (-0.19, -0.06)	--	-0.02 (-0.08, 0.04)	--	-0.01 (-0.06, 0.05)	--
Age 14 mo	-0.09 (-0.15, -0.03)	0.73	-0.10 (-0.16, -0.05)	0.68	-0.08 (-0.13, -0.02)	0.22	0.01 (-0.04, 0.07)	0.42	0.03 (-0.03, 0.08)	0.41
Age 28 mo	-0.00 (-0.04, 0.04)	0.02 *	-0.03 (-0.07, 0.01)	0.03 *	-0.05 (-0.10, -0.00)	0.04 *	-0.05 (-0.09, -0.01)	0.47	-0.02 (-0.06, 0.02)	0.71
Female	-0.06 (-0.11, -0.01)	--	-0.06 (-0.11, -0.02)	--	-0.08 (-0.12, -0.03)	--	-0.01 (-0.06, 0.03)	--	-0.01 (-0.05, 0.03)	--
Male	-0.07 (-0.12, -0.02)	0.82	-0.10 (-0.15, -0.05)	0.26	-0.09 (-0.15, -0.04)	0.64	-0.02 (-0.07, 0.02)	0.78	0.01 (-0.04, 0.06)	0.55
	CD (95% CI)	p-value ^a	CD (95% CI)	p-value ^a	CD (95% CI)	p-value ^a	CD (95% CI)	p-value ^a	CD (95% CI)	p-value ^a
Episodes of antibiotic use										
All obs	-0.17 (-0.26, -0.07)	<0.001	-0.20 (-0.29, -0.11)	<0.001	-0.21 (-0.29, -0.12)	<0.001	-0.04 (-0.13, 0.04)	0.35	-0.01 (-0.09, 0.08)	0.87
Age 3 mo	-0.25 (-0.43, -0.07)	--	-0.32 (-0.50, -0.14)	--	-0.34 (-0.51, -0.17)	--	-0.06 (-0.13, 0.02)	--	-0.02 (-0.10, 0.05)	--
Age 14 mo	-0.18 (-0.32, -0.04)	0.52	-0.19 (-0.32, -0.05)	0.23	-0.15 (-0.28, -0.02)	0.05 *	-0.03 (-0.10, 0.04)	0.64	0.02 (-0.05, 0.09)	0.37
Age 28 mo	-0.08 (-0.19, 0.03)	0.11 *	-0.12 (-0.24, -0.00)	0.09 *	-0.15 (-0.26, -0.04)	0.08 *	-0.00 (-0.05, 0.05)	0.23	-0.02 (-0.07, 0.03)	0.95
Female	-0.18 (-0.30, -0.06)	--	-0.22 (-0.33, -0.11)	--	-0.22 (-0.32, -0.12)	--	-0.04 (-0.15, 0.06)	--	-0.00 (-0.11, 0.10)	--
Male	-0.16 (-0.29, -0.03)	0.84	-0.19 (-0.32, -0.06)	0.70	-0.18 (-0.32, -0.05)	0.66	-0.02 (-0.15, 0.10)	0.81	0.00 (-0.12, 0.13)	0.94
Total days of antibiotic use										
All obs	-0.94 (-1.41, -0.48)	<0.001	-1.05 (-1.54, -0.55)	<0.001	-1.01 (-1.51, -0.51)	<0.001	-0.07 (-0.55, 0.41)	0.79	0.03 (-0.42, 0.49)	0.88
Age 3 mo	-1.23 (-2.20, -0.26)	--	-1.77 (-2.75, -0.79)	--	-1.52 (-2.55, -0.49)	--	-0.09 (-0.25, 0.06)	--	-0.02 (-0.17, 0.13)	--
Age 14 mo	-1.14 (-1.86, -0.43)	0.87	-0.93 (-1.69, -0.18)	0.17 *	-0.57 (-1.41, 0.27)	0.12 *	0.04 (-0.11, 0.18)	0.20	0.04 (-0.11, 0.19)	0.52
Age 28 mo	-0.49 (-1.16, 0.18)	0.25	-0.60 (-1.32, 0.12)	0.07 *	-0.98 (-1.60, -0.36)	0.41	-0.07 (-0.16, 0.02)	0.80	-0.03 (-0.13, 0.08)	0.95
Female	-0.95 (-1.72, -0.18)	--	-1.10 (-1.85, -0.35)	--	-1.12 (-1.75, -0.49)	--	-0.17 (-0.76, 0.42)	--	-0.02 (-0.65, 0.60)	--
Male	-0.98 (-1.63, -0.33)	0.96	-1.03 (-1.74, -0.32)	0.90	-0.85 (-1.60, -0.11)	0.58	0.13 (-0.62, 0.88)	0.54	0.17 (-0.55, 0.90)	0.70

PD: Prevalence difference (for binary outcomes); CD: Count difference (for count outcomes); CI: Confidence interval; C: Control; WSH: Water, sanitation, hygiene; N: Nutrition, N+WSH: Nutrition+ water, sanitation, hygiene.

Bolded values refer to estimates with a p-value ≤0.05.

^a p-value for interaction term between study arm and subgroup variable.

* Interaction p-values <0.20 indicate significant effect modification.

Table S14. Multiplicative interaction by age and sex for unadjusted intervention effects on antibiotic use within last 3 months, Kenya

	WSH vs. C		N vs. C		N+WSH vs. C		N+WSH vs. WSH		N+WSH vs. N	
	PR (95% CI)	p-value ^a	PR (95% CI)	p-value ^a	PR (95% CI)	p-value ^a	PR (95% CI)	p-value ^a	PR (95% CI)	p-value ^a
Used antibiotics ≥1 time										
All obs	0.96 (0.86, 1.06)	0.41	0.96 (0.88, 1.04)	0.33	1.01 (0.92, 1.10)	0.84	1.05 (0.95, 1.17)	0.33	1.05 (0.95, 1.17)	0.32
Age 6 mo	0.95 (0.83, 1.08)	--	0.94 (0.82, 1.08)	--	0.99 (0.84, 1.16)	--	1.04 (0.87, 1.25)	--	1.06 (0.89, 1.24)	--
Age 17 mo	1.01 (0.85, 1.20)	0.57	1.03 (0.90, 1.17)	0.34	1.04 (0.91, 1.20)	0.63	1.03 (0.91, 1.18)	0.93	1.02 (0.88, 1.17)	0.75
Age 22 mo	0.91 (0.76, 1.09)	0.74	0.90 (0.78, 1.04)	0.70	1.00 (0.85, 1.16)	0.94	1.09 (0.92, 1.30)	0.70	1.10 (0.94, 1.29)	0.67
Female	0.93 (0.81, 1.06)	--	0.96 (0.86, 1.07)	--	0.99 (0.85, 1.15)	--	1.07 (0.93, 1.22)	--	1.03 (0.89, 1.20)	--
Male	0.99 (0.86, 1.14)	0.50	0.96 (0.85, 1.08)	0.98	1.03 (0.92, 1.15)	0.72	1.04 (0.89, 1.21)	0.79	1.07 (0.96, 1.21)	0.67
Used antibiotics ≥2 times										
All obs	1.03 (0.80, 1.32)	0.81	0.88 (0.68, 1.12)	0.30	0.98 (0.78, 1.22)	0.84	0.95 (0.71, 1.27)	0.72	1.12 (0.82, 1.52)	0.48
Age 6 mo	0.83 (0.60, 1.14)	--	0.85 (0.59, 1.23)	--	0.86 (0.59, 1.26)	--	1.04 (0.67, 1.62)	--	1.01 (0.65, 1.56)	--
Age 17 mo	1.19 (0.77, 1.84)	0.20	1.03 (0.66, 1.60)	0.52	1.17 (0.79, 1.72)	0.31	0.98 (0.62, 1.55)	0.85	1.14 (0.76, 1.71)	0.65
Age 22 mo	1.15 (0.75, 1.75)	0.18 *	0.72 (0.48, 1.08)	0.54	0.91 (0.59, 1.41)	0.84	0.80 (0.48, 1.32)	0.39	1.27 (0.80, 1.99)	0.41
Female	0.99 (0.74, 1.33)	--	1.00 (0.72, 1.37)	--	0.97 (0.70, 1.34)	--	0.98 (0.71, 1.35)	--	0.97 (0.66, 1.43)	--
Male	1.08 (0.75, 1.56)	0.70	0.77 (0.53, 1.13)	0.31	1.00 (0.73, 1.37)	0.90	0.92 (0.59, 1.44)	0.83	1.29 (0.85, 1.96)	0.27
	CR (95% CI)	p-value ^a	CR (95% CI)	p-value ^a	CR (95% CI)	p-value ^a	CR (95% CI)	p-value ^a	CR (95% CI)	p-value ^a
Episodes of antibiotic use										
All obs	0.98 (0.87, 1.10)	0.77	0.95 (0.85, 1.07)	0.45	1.02 (0.92, 1.14)	0.66	1.04 (0.91, 1.19)	0.56	1.07 (0.93, 1.23)	0.34
Age 6 mo	0.92 (0.79, 1.08)	--	0.94 (0.78, 1.14)	--	0.96 (0.79, 1.17)	--	1.05 (0.83, 1.32)	--	1.02 (0.82, 1.28)	--
Age 17 mo	1.06 (0.87, 1.29)	0.31	1.05 (0.89, 1.22)	0.42	1.11 (0.95, 1.29)	0.31	1.05 (0.87, 1.26)	0.99	1.06 (0.89, 1.26)	0.77
Age 22 mo	0.96 (0.79, 1.18)	0.72	0.87 (0.75, 1.01)	0.45	1.00 (0.84, 1.18)	0.83	1.03 (0.83, 1.27)	0.91	1.14 (0.96, 1.36)	0.37
Female	0.96 (0.83, 1.11)	--	0.99 (0.86, 1.13)	--	1.01 (0.86, 1.19)	--	1.05 (0.90, 1.24)	--	1.03 (0.86, 1.23)	--
Male	1.01 (0.85, 1.19)	0.63	0.93 (0.79, 1.09)	0.53	1.04 (0.91, 1.19)	0.83	1.03 (0.84, 1.26)	0.84	1.12 (0.95, 1.32)	0.42
Total days of antibiotic use										
All obs	1.05 (0.71, 1.54)	0.81	0.85 (0.65, 1.12)	0.24	0.94 (0.76, 1.17)	0.57	0.90 (0.64, 1.27)	0.52	1.10 (0.87, 1.40)	0.41
Age 6 mo	0.72 (0.42, 1.25)	--	0.52 (0.35, 0.77)	--	0.70 (0.43, 1.14)	--	0.97 (0.62, 1.51)	--	1.36 (0.97, 1.90)	--
Age 17 mo	1.57 (0.79, 3.13)	0.08 *	1.21 (0.75, 1.95)	0.01 *	1.04 (0.74, 1.46)	0.26	0.66 (0.34, 1.30)	0.33	0.86 (0.56, 1.33)	0.12 *
Age 22 mo	1.00 (0.78, 1.28)	0.27	1.01 (0.75, 1.37)	0.01 *	1.22 (0.91, 1.65)	0.06 *	1.22 (0.92, 1.63)	0.42	1.21 (0.87, 1.67)	0.59
Female	1.10 (0.67, 1.83)	--	0.88 (0.62, 1.24)	--	0.88 (0.64, 1.20)	--	0.80 (0.50, 1.26)	--	1.00 (0.72, 1.39)	--
Male	0.97 (0.67, 1.40)	0.60	0.83 (0.57, 1.21)	0.83	1.01 (0.73, 1.39)	0.55	1.04 (0.74, 1.46)	0.27	1.22 (0.88, 1.70)	0.40

PR: Prevalence ratio (for binary outcomes), CR: Count ratio (for count outcomes); CI: Confidence interval; C: Control; WSH: Water, sanitation, hygiene; N: Nutrition, N+WSH: Nutrition+ water, sanitation, hygiene.

Bolded values refer to estimates with a p-value ≤0.05.

^a p-value for interaction term between study arm and subgroup variable.

* Interaction p-values <0.20 indicate significant effect modification.

Table S15. Additive interaction by age and sex for unadjusted intervention effects on antibiotic use within last 3 months, Kenya

	WSH vs. C		N vs. C		N+WSH vs. C		N+WSH vs. WSH		N+WSH vs. N	
	PD (95% CI)	p-value ^a	PD (95% CI)	p-value ^a	PD (95% CI)	p-value ^a	PD (95% CI)	p-value ^a	PD (95% CI)	p-value ^a
Used antibiotics ≥1 time										
All obs	-0.02 (-0.08, 0.03)	0.41	-0.02 (-0.07, 0.02)	0.33	0.00 (-0.05, 0.05)	0.84	0.03 (-0.03, 0.08)	0.33	0.03 (-0.03, 0.08)	0.32
Age 6 mo	-0.03 (-0.10, 0.04)	--	-0.03 (-0.10, 0.04)	--	-0.01 (-0.09, 0.08)	--	0.02 (-0.07, 0.11)	--	0.03 (-0.06, 0.11)	--
Age 17 mo	0.01 (-0.08, 0.10)	0.58	0.01 (-0.05, 0.08)	0.35	0.02 (-0.05, 0.10)	0.62	0.02 (-0.05, 0.09)	0.95	0.01 (-0.07, 0.09)	0.77
Age 22 mo	-0.05 (-0.13, 0.4)	0.75	-0.05 (-0.12, 0.02)	0.72	-0.00 (-0.08, 0.08)	0.94	0.04 (-0.04, 0.13)	0.71	0.05 (-0.03, 0.13)	0.68
Female	-0.04 (-0.11, 0.03)	--	-0.02 (-0.08, 0.04)	--	-0.00 (-0.08, 0.07)	--	0.03 (-0.04, 0.10)	--	0.02 (-0.06, 0.09)	--
Male	-0.01 (-0.08, 0.07)	0.50	-0.02 (-0.09, 0.04)	0.99	0.01 (-0.05, 0.07)	0.72	0.02 (-0.06, 0.10)	0.80	0.04 (-0.02, 0.10)	0.67
Used antibiotics ≥2 times										
All obs	0.00 (-0.03, 0.04)	0.81	-0.02 (-0.05, 0.01)	0.30	-0.00 (-0.03, 0.03)	0.84	-0.01 (-0.05, 0.03)	0.73	0.01 (-0.02, 0.05)	0.49
Age 6 mo	-0.03 (-0.08, 0.02)	--	-0.03 (-0.08, 0.03)	--	-0.02 (-0.08, 0.04)	--	0.01 (-0.06, 0.07)	--	0.00 (-0.06, 0.07)	--
Age 17 mo	0.02 (-0.04, 0.08)	0.20	0.00 (-0.05, 0.06)	0.48	0.02 (-0.03, 0.07)	0.31	-0.00 (-0.07, 0.06)	0.85	0.02 (-0.04, 0.07)	0.68
Age 22 mo	0.02 (-0.04, 0.07)	0.17 *	-0.03 (-0.07, 0.01)	0.86	-0.01 (-0.06, 0.04)	0.72	-0.03 (-0.08, 0.03)	0.42	0.02 (-0.02, 0.06)	0.55
Female	-0.00 (-0.04, 0.04)	--	-0.00 (-0.04, 0.04)	--	-0.00 (-0.05, 0.04)	--	-0.00 (-0.04, 0.04)	--	-0.00 (-0.05, 0.04)	--
Male	0.01 (-0.04, 0.07)	0.69	-0.03 (-0.08, 0.01)	0.32	-0.00 (-0.04, 0.04)	0.91	-0.01 (-0.08, 0.05)	0.81	0.03 (-0.02, 0.08)	0.28
	CD (95% CI)	p-value ^a	CD (95% CI)	p-value ^a	CD (95% CI)	p-value ^a	CD (95% CI)	p-value ^a	CD (95% CI)	p-value ^a
Episodes of antibiotic use										
All obs	-0.01 (-0.09, 0.07)	0.77	-0.03 (-0.10, 0.05)	0.44	0.01 (-0.06, 0.09)	0.67	0.03 (-0.06, 0.12)	0.56	0.05 (-0.05, 0.14)	0.34
Age 6 mo	-0.06 (-0.17, 0.05)	--	-0.04 (-0.18, 0.10)	--	-0.03 (-0.17, 0.12)	--	0.03 (-0.13, 0.19)	--	0.01 (-0.14, 0.17)	--
Age 17 mo	0.04 (-0.10, 0.17)	0.31	0.03 (-0.08, 0.13)	0.41	0.07 (-0.04, 0.18)	0.32	0.03 (-0.10, 0.16)	1.00	0.04 (-0.08, 0.16)	0.78
Age 22 mo	-0.02 (-0.15, 0.10)	0.67	-0.08 (-0.18, 0.01)	0.57	-0.01 (-0.11, 0.10)	0.82	0.02 (-0.11, 0.15)	0.89	0.08 (-0.03, 0.18)	0.45
Female	-0.02 (-0.12, 0.07)	--	-0.01 (-0.10, 0.08)	--	0.01 (-0.10, 0.12)	--	0.04 (-0.07, 0.14)	--	0.02 (-0.10, 0.14)	--
Male	0.01 (-0.11, 0.12)	0.64	-0.05 (-0.16, 0.06)	0.53	0.03 (-0.07, 0.12)	0.83	0.02 (-0.12, 0.16)	0.85	0.07 (-0.03, 0.18)	0.43
Total days of antibiotic use										
All obs	0.19 (-1.37, 1.76)	0.81	-0.59 (-1.61, 0.43)	0.25	-0.24 (-1.08, 0.60)	0.58	-0.43 (-1.81, 0.94)	0.53	0.34 (-0.48, 1.18)	0.41
Age 6 mo	-1.55 (-4.27, 1.17)	--	-2.71 (-4.88, -0.53)	--	-1.67 (-4.16, 0.81)	--	-0.13 (-1.91, 1.66)	--	1.03 (-0.20, 2.26)	--
Age 17 mo	1.95 (-1.47, 5.37)	0.11 *	0.73 (-1.08, 2.54)	0.02 *	0.14 (-1.03, 1.31)	0.26	-1.81 (-5.22, 1.60)	0.37	-0.59 (-2.30, 1.12)	0.15 *
Age 22 mo	0.00 (-0.72, 0.73)	0.27	0.04 (-0.84, 0.93)	0.03 *	0.66 (-0.35, 1.67)	0.09 *	0.66 (-0.32, 1.63)	0.47	0.62 (0.46, 1.70)	0.58
Female	0.43 (-1.79, 2.64)	--	-0.51 (-1.84, 0.82)	--	-0.50 (-1.72, 0.72)	--	-0.93 (-2.92, 1.06)	--	0.01 (-1.16, 1.18)	--
Male	-0.12 (-1.50, 1.27)	0.61	-0.65 (-1.99, 0.69)	0.88	0.04 (-1.18, 1.26)	0.54	0.16 (-1.11, 1.42)	0.29	0.69 (-0.47, 1.85)	0.41

PD: Prevalence difference (for binary outcomes); CD: Count difference (for count outcomes); CI: Confidence interval; C: Control; WSH: Water, sanitation, hygiene; N: Nutrition, N+WSH: Nutrition+ water, sanitation, hygiene.

Bolded values refer to estimates with a p-value ≤0.05.

^a Interaction p-value between study arm and subgroup variable.

* Interaction p-values <0.20 indicate significant effect modification.

Table S16. Sensitivity analysis, unadjusted intervention effects on antibiotic use within last 2 weeks and last month, Bangladesh*

Arm	N	% (n)	Compared to C		Compared to WSH		Compared to N	
			PR (95% CI)	p-value	PR (95% CI)	p-value	PR (95% CI)	p-value
Bangladesh								
Used antibiotics ≥ 1 time in last two weeks								
C	1000	21.5 (215)	--	--	--	--	--	--
WSH	1076	22.1 (238)	1.03 (0.84, 1.26)	0.79	--	--	--	--
N	1014	19.9 (202)	0.93 (0.77, 1.11)	0.42	--	--	--	--
N+WSH	1050	19.2 (202)	0.89 (0.73, 1.10)	0.29	0.87 (0.72, 1.06)	0.16	0.97 (0.78, 1.19)	0.74
Used antibiotics ≥ 1 time in last month								
C	1000	48.2 (482)	--	--	--	--	--	--
WSH	1076	43.2 (465)	0.90 (0.81, 0.99)	0.03	--	--	--	--
N	1014	40.4 (410)	0.84 (0.76, 0.93)	<0.001	--	--	--	--
N+WSH	1050	42.3 (444)	0.88 (0.79, 0.97)	0.01	0.98 (0.88, 1.08)	0.67	1.05 (0.93, 1.18)	0.45

PR: Prevalence ratio; CI: Confidence interval; C: Control; WSH: Water, sanitation, hygiene; N: Nutrition, N+WSH: Nutrition+ water, sanitation, hygiene.

* Not conducted for Kenya because the survey question for this variable in Kenya was designed to also capture non-antibiotic medications.

Figure S1. CONSORT Diagram for the WASH Benefits Bangladesh environmental enteric dysfunction substudy




Figure S2. CONSORT Diagram for the WASH Benefits Kenya environmental enteric dysfunction substudy

