

Supplement Table II. Univariate analysis of drivers of RVF clusters in East Africa, 2008 –2022

| Variable | Coefficient | IRR | 95% CI | S.E | p-value |
|-----------------------------------------------------------------------------|-------------|---------------|---------------|---------|---------|
| Country* | | | | | |
| <i>Kenya (Intercept)</i> | -3.86 | - | 0.01, 0.03 | 0.19 | <0.001 |
| <i>Uganda</i> | -1.55 | 0.21 | 0.12, 0.38 | 0.28 | <0.001 |
| <i>Tanzania</i> | 0.90 | 2.46 | 1.14, 5.66 | 0.39 | 0.02* |
| Livestock production system* | | | | | |
| <i>Extensive (Intercept)</i> | -3.41 | - | 0.02, 0.05 | 0.22 | <0.001 |
| <i>Intensive</i> | -0.63 | 0.531 | 1.13, 3.93 | 0.35 | 0.06 |
| <i>Semi-intensive</i> | -1.89 | 0.150 | 1.08, 1.32 | 0.31 | <0.001 |
| Elevation level* | | | | | |
| <i>High (> 1000m) (Intercept)</i> | -5.17 | - | 0.004, 0.008 | 0.15 | <0.001 |
| <i>Low (<500m)</i> | 2.29 | 9.96 | 5.81, 17.60 | 0.27 | <0.001 |
| <i>Medium (500 - 1000m)</i> | 0.47 | 1.61 | 0.68, 4.15 | 0.44 | 0.28 |
| Rainfall prior to outbreak (mm) | | | | | |
| 3 rd month | 0.22 | 1.25 | 1.16, 1.34 | 0.0385 | <0.001 |
| 2 nd month | 0.22 | 1.24 | 1.15, 1.34 | 0.0411 | <0.001 |
| 1 st month | 0.18 | 1.19 | 1.11, 1.30 | 0.0458 | <0.001 |
| Outbreak month | 0.21 | 1.2438 | 1.13, 1.36 | 0.0523 | <0.001 |
| Humidity outbreak month | 0.04 | 1.04 | 1.01, 1.08 | 0.01751 | 0.01* |
| Positive change in NDVI (1stmonth prior) | | | | | |
| <i>No (Intercept)</i> | - | - | | - | - |
| <i>Yes</i> | 0.31 | 1.36 | 0.71, 2.60 | 0.31 | 0.32 |
| NDVI change between 2nd month prior & outbreak month* | | | | | |
| <i>No (Intercept)</i> | - | - | | - | - |
| <i>Yes</i> | 4.60 | 99.74 | 13.01, 812.40 | 0.97 | <0.001 |

Note S.E= standard error, IRR = incident rate, significant variables are in bold