

Supplementary Table S4. List of variables considered and included in propensity-score model

Potential confounders	Selected into propensity-score model	Reason for omitting
Stroke/TIA	✓	
Peripheral artery disease	✓	
Coronary artery disease	✓	
Diabetes	✓	
High-risk diabetes	✓	
Age (years)	✓	
Sex	✓	
Ethnicity	✓	
BMI	✓	7.6% missing
SBP	✓	22.5% missing
DBP	✓	22.5% missing
Creatinine	✓	19.5% missing
Index of Multiple Deprivation (IMD)	✓	0.1% missing
Smoke status	✓	2.7% missing
Alcohol use		16.9% missing
Statin use	✓	
Nitrate use	✓	
Diabetic treatment use	✓	
Diuretic use	✓	
CCB use	✓	
Betablocker use	✓	
Aspirin use	✓	
Antiplatelet use	✓	
Digoxin use		Insufficient number of events
Anticoagulant use	✓	
Alpha-blocker use	✓	
No. of hospital admissions within 6 months prior	✓	
No. of GP appointments within 6 months prior	✓	
Year of start of eligible period	✓	
Time since first eligible period (days)	✓	

Notes: TIA: transient ischaemic attack; BMI: body-mass index; SBP: systolic blood pressure; DBP: diastolic blood pressure.

Variables are measured at start of trial-eligible period or before.

Peripheral artery disease includes limb bypass surgery or angioplasty, limb/foot amputation, or intermittent claudication.

Coronary artery disease includes previous MI, angina, coronary angioplasty, or CABG.

SBP and DBP are measured within 6 months prior to start of trial-eligible period.

Medication use is within 3 months prior to start of trial-eligible period.

SBP and DBP had 22.5% missing but this variable was included as believed to be an important confounder.

Similarly, baseline creatinine had 19.5% missing but due to the known differences in prescribing based on baseline creatinine this variable was also included in the model.

Balance after weighting was assessed for all variables listed including those not included in the propensity-score model but could be assumed to be missing at random. If imbalance remained analysis was adjusted for these variables.