

Supplementary Table 1. OPHION Opioid Use Sub-Group Characteristics

Characteristic	Active injection	Methadone	Suboxone	Chronic Pain	No opioid	Total*
Participant, N	4	4	12	4	12	36
Age (years) Median (IQR) [†]	56 (46-61)	59 (55-65)	61 (57-67)	61 (57-67)	64 (59-65)	59 (54-65)
Sex Male N (%)	4 (100%)	2 (50%)	9 (75%)	2 (50%)	7 (58%)	24 (67%)
Race Black N (%)	1 (25%)	1 (25%)	3 (25%)	1 (25%)	9 (75%)	15 (42%)
White N (%)	1 (25%)	2 (50%)	6 (50%)	2 (50%)	2 (17%)	11 (31%)
Hispanic/Latino N (%)	0	1 (25%)	1 (8.3%)	0	1 (8%)	3 (8%)
American Indian N (%)	0	0	0	1 (25%)	0	1 (3%)
Declined/NA	2 (50%)	2 (50%)	2 (17%)	0	0	6 (17%)
Ethnicity Non-Hispanic N (%)	2 (50%)	1 (25%)	7 (58%)	3 (75%)	11 (92%)	24 (67%)
Hispanic N (%)	2 (50%)	3 (75%)	5 (42%)	1 (25%)	1 (8%)	12 (33%)
ART Duration (months) Median (IQR)	146 (96-209)	94 (79-111)	132 (81-167)	173 (161-188)	162 (106-204)	144 (97-179)
Duration of virus suppression (months) Median (IQR)	76 (51-96)	62 (35-95)	79 (64-89)	77 (41-101)	70 (38-91)	75 (44-93)

*Percentage totals may not add up to 100 due to rounding. [†]IQR, interquartile range