Supplementary materials

Supplementary tables

Set	Number	Age (Mean	Gender	Dataset	Country	Education years
		± std)				(Mean \pm std)
Training	91	60 ± 14	F=47;	PRISM=16;	NL=83;	7 ± 5
			M=44	HO=75	ES=8	
Validation	156	58 ± 12	F=93;	PRISM=12;	NL=151;	6 ± 3
			M=63	HO=144	ES=5	

Table S1: Demographics of healthy controls with training/validation split.

*std: standard deviation; F: female, M: male; NL: the Netherlands, ES: Spain.

Table S2: Results from linear regression models predicting Loneliness score from dwell time.

Group	Predictor	Estimate	Standard error	t value	<i>p</i> -value
All validation	Dwell time	-0.0006	0.0023	-0.2782	0.7821
(n=49)	Age	-0.0047	0.0029	-1.6400	0.1078
Healthy control	Dwell time	-0.0013	0.0015	-0.8399	0.4227
(validation)	Age	0.0063	0.0021	3.0350	0.0141
(n=12)					
Schizophrenia	Dwell time	-0.0023	0.0050	-0.4706	0.6447
(n=18)	Age	-0.0051	0.0152	-0.3366	0.7411
Alzheimer's	Dwell time	0.0033	0.0029	1.1651	0.2611
disease (n=19)	Age	0.0076	0.0048	1.5824	0.1331

Table S3: Binomial logistic regression age sensitivity analysis results; age cut-offs for healthy controls (HC) matched to each diagnostic group: HC vs SZ, AGE < 42; HC vs AD, AGE > 50; HC vs SCC, AGE > 43.

Group	Coefficient	Standard	Odds	z value	<i>p</i> -value	FDR
		error				corrected
						<i>p</i> -value
Schizophrenia	-0.0034	0.0184	0.9966	-0.1858	0.8526	1.0000
Alzheimer's disease	-0.0576	0.0148	0.9440	-3.8989	0.0001	0.0003*
Healthy/subjective	-0.0165	0.0082	0.9837	-2.0152	0.0439	0.1316
cognitive						
complaints						

Group	Estimate	Standard error	t value	<i>p</i> -value	FDR corrected
					<i>p</i> -value
All validation	-0.0416	0.0296	-1.4059	0.1708	0.5123
Healthy control	-0.0098	0.0144	-0.6826	0.5120	1.0000
(validation)					
Alzheimer's	0.0439	0.0569	0.7708	0.4521	1.0000
disease					

Table S4: Age-related results from linear regression models predicting MMSE score.

Table S5:1 Results from linear regression models predicting PANSS (sub)scores in participants with schizophrenia (n=18) (PANSS_PT: Positive Scale, PANSS_NT: Negative Scale, PANSS_GT: General Psychopathology Scale, PANSS_CS: Composite Scale).

Response	Predictor	Estimate	Standard error	t value	<i>p</i> -value
PANSS_PT	Dwell time	-0.0264	0.0491	-0.5368	0.5993
	Age	-0.0099	0.1499	-0.0658	0.9484
PANSS_NT	Dwell time	0.0043	0.0907	0.0472	0.9630
	Age	-0.0852	0.2770	-0.3077	0.7625
PANSS_GT	Dwell time	0.0653	0.0866	0.7542	0.4624
	Age	0.2041	0.2643	0.7722	0.4520
PANSS_CS	Dwell time	-0.0306	0.0736	-0.4166	0.6829
	Age	0.0754	0.2246	0.3357	0.7417
PANSS_TOTAL	Dwell time	0.0432	0.1995	0.2166	0.8315
	Age	0.1090	0.6090	0.1789	0.8604

Supplementary figures




Figure S1: Age distributions of training participants and all validation participants.




Figure S2: SFS scores of training participants and all validation participants (possible range: 55-135).




Figure S3: SFS scores of validation participants, split by diagnostic group.




Figure S4: Loneliness scores of training participants and all validation participants (possible range: 0-11).




Figure S5: Loneliness scores of validation participants, split by diagnostic group.




Figure S6: MMSE scores of training participants and all validation participants (maximum score = 30).




Figure S7: MMSE scores of validation participants, split by diagnostic group.




Figure S8: PANSS scores, split by (sub)scale. Positive (PT) and Negative (NT) Scales range: 7-49; General Psychopathology (GT) Scale range: 16-112; Total PANSS score: sum of the sub scales; Composite Scale (CS) range: -42 to +42 (subtract negative score from positive score).




Figure S9: Distributions of hourly percentage values per channel for all time series; these distributions are very zero-inflated.




Figure S10: Percentage of overall data availability (indicating missingness) for training participants and all validation participants.




Figure S11: Percentage of overall data availability (indicating missingness) for validation participants, split by diagnostic group.




Figure S12: Percentage of GPS data availability (indicating missing GPS data) for training participants and all validation participants.




Figure S13: Percentage of GPS data availability (indicating missing GPS data) for validation participants, split by diagnostic group.




Figure S14: Days of data for training and validation participants (i.e. time series length).




Figure S15: Days of data for validation participants (i.e. time series length).