

**Supplementary Table 1. Subject characteristics at baseline and changes observed after 12 weeks of exercise.**

	Baseline		%-change	
	NW (n = 13)	OW (n = 13)	NW (n = 13)	OW (n = 13)
Age (years)	50 (7)	53 (6)		
<b>Body composition</b>				
Weight (kg)	78.5 (8.2)	95.4 (10.2) *	-0.3 (2.1)	-1.7 (2.4) †
BMI (kg/m <sup>2</sup> )	23.5 (2.0)	29.0 (2.4) *	0.0 (2.0)	-1.2 (4.5)
FFM (kg) <sup>a</sup>	34.9 (3.5)	37.7 (5.0)	6.4 (3.8) †	5.3 (2.7) †
ScWAT (kg) <sup>a</sup>	10.3 (2.7)	18.0 (4.2) *	-6.6 (9.2) †	-7.3 (6.0) †
IAAT (kg) <sup>a</sup>	4.0 (2.0)	8.8 (2.6) *	-16.9 (15.1) †	-19.4 (10.8) †
Hepatic fat (AU) <sup>b</sup>	2.8 (2.2)	9.1 (5.9) *	-23.3 (50.7) †	-27.4 (15.8) †
Thigh muscle area (AU) <sup>a</sup>	20.3 (2.9)	24.4 (3.1) *	9.7 (4.7) †	7.1 (6.7) †
<b>Physical fitness</b>				
VO <sub>2</sub> max (ml/kg/min)	44.1 (4.4)	37.1 (4.9) †	13.2 (9.7) †	13.3 (7.7) †
Chest press (kg)	65.6 (16.8)	68.7 (13.7)	18.4 (8.7) †	13.6 (8.4) †
Pull down (kg)	68.8 (9.3)	75.6 (15.1)	18.3 (10.1) †	13.7 (7.3) †
Leg press (kg)	199.6 (36.9)	248.7 (30.3) *	9.8 (7.6) †	12.5 (8.4) †
<b>Glucose metabolism</b>				
HbA1c (mmol/mol)	33 (4)	37 (4) *	N.A.	N.A.
HbA1c (%)	5.2 (0.2)	5.5 (0.4) *	N.A.	N.A.
Glucose (mmol/l)	5.4 (0.5)	5.9 (0.3) *	3.1 (4.5) †	1.8 (6.8)
C-Peptide (pmol/l)	588.0 (117.8)	932.8 (248.9) *	7.3 (23.8)	12.3 (45.3)
Insulin (pmol/l)	38.5 (18.6)	65.3 (27.1) *	15.1 (49.2)	27.6 (66.2)
GIR (mg/kg/min)	7.6 (1.6)	4.2 (1.8) *	37.8 (30.1) †	44.4 (58.8) †

**Notes:** <sup>a</sup> n = 12 control, <sup>b</sup> n = 10 control, and n = 9 OW. \* p < 0.05 between groups (OW vs. NW), and † p < 0.05 baseline vs. 12 w within group. Between groups comparison were performed using unpaired t-tests, and within group comparisons were performed using paired t-tests. Logarithmic transformation was performed to approximate normal distribution, if necessary, and back transformed for presentation. Data represent means (SD).

**Abbreviations:** NW = normal weight. OW = overweight. N.A. = not available, AU = arbitrary units, BMI = body mass index, FFM = fat free mass, AT = adipose tissue, Sc = subcutaneous, IA = intra-abdominal, GIR = glucose infusion rate.

*Supplementary Table 5: Descriptive statistics UK biobank*

	All individuals (n=52,981)		Women (n=28,579)		Men (n=24,402)		Units
	Mean	SD	Mean	SD	Mean	SD	
Age	56.19	8.34	55.83	8.26	56.61	8.41	Years
Body mass index	27.27	4.69	26.85	5.09	27.76	4.11	kg/m <sup>2</sup>
Glucose	5.10	1.99	5.04	1.05	5.17	1.34	mmol/L
HbA1c	35.81	6.44	35.48	5.80	36.20	7.09	mmol/mol
Triglycerides	1.01	1.73	1.53	0.86	1.96	1.13	mmol/L
Type 2 diabetes	2,618 cases	-	865 cases	-	1753 cases	-	Yes/No.
Sample storage time	665.40	43.69	665.60	43.47	665.30	43.95	Weeks

*Supplementary Table 6. Serum CD300LG cis-pQTLs*

SNP	Chromosome	Base position	Effect allele	Other allele	Effect allele frequency	INFO score	Effect	SE	<i>p</i> -value	F-statistic
rs72831096	17	41469246	G	A	0.7937	0.9978	0.0312	0.0033	$6.3 \times 10^{-21}$	88.0903
rs851057	17	41837264	G	C	0.1283	0.9681	-0.0981	0.0041	$3.8 \times 10^{-129}$	584.5963
rs231474	17	42016477	T	G	0.4528	0.9942	0.0327	0.0027	$1.1 \times 10^{-33}$	146.4119

SNP = single nucleotide polymorphism. SE = standard error.

Supplementary Table 7. Serum CD300LG cis-pQTLs

SNP	CHR	Base position	Effect allele	Other allele	Effect allele frequency	INFO score	Effect	SE	p-value
rs76230830	1	23784293	A	G	0.918135	0.986397	0.027182	0.004972	4.60E-08
rs4651034	1	1.79E+08	A	C	0.663704	0.996121	-0.01747	0.002868	1.10E-09
rs3753847	1	2.05E+08	T	C	0.151267	0.973821	0.021328	0.003809	2.20E-08
rs12040264	1	2.2E+08	G	T	0.397163	0.99713	0.016136	0.002766	5.40E-09
rs1260326	2	27730940	T	C	0.391047	1	-0.02078	0.002772	6.70E-14
rs75166367	2	1.63E+08	G	A	0.93965	1	0.03759	0.005696	4.10E-11
rs12328675	2	1.66E+08	T	C	0.881422	0.99872	-0.02457	0.004186	4.30E-09
rs56256300	2	2.27E+08	A	AT	0.35506	0.999036	0.017272	0.002841	1.20E-09
rs1899951	3	12394840	C	T	0.877417	0.997266	-0.0388	0.004135	6.30E-21
rs77572134	3	52895934	T	C	0.858413	0.997026	-0.0235	0.003886	1.50E-09
rs11943206	4	56437010	T	C	0.700938	0.99321	-0.01638	0.002954	2.90E-08
rs13127398	4	1.03E+08	T	A	0.92883	0.985517	-0.03515	0.005331	4.30E-11
rs574745983	5	53302258	A	AT	0.752631	0.981441	0.017945	0.003171	1.50E-08
rs164514	5	1.41E+08	A	G	0.658169	0.996885	0.016234	0.002874	1.60E-08
rs1064627	6	30698541	A	G	0.8	0.999598	-0.01976	0.003383	5.20E-09
rs614008	6	31840794	C	T	0.421165	0.996933	0.017261	0.002743	3.10E-10
rs75104038	6	34190104	G	A	0.938486	0.992843	0.031943	0.005663	1.70E-08
rs998584	6	43757896	C	A	0.520758	0.994408	0.03954	0.002717	5.40E-48
rs9402846	6	1.37E+08	G	C	0.410445	0.990684	-0.01628	0.002767	4.00E-09
rs28588745	8	10647044	A	T	0.792571	0.995694	0.047092	0.003354	8.60E-45
rs13267474	8	23784587	C	T	0.590946	0.971999	-0.01559	0.002791	2.30E-08
rs2980888	8	1.27E+08	T	C	0.300429	0.999267	-0.02082	0.002966	2.20E-12
rs8176747	9	1.36E+08	C	G	0.937707	0.999944	-0.10105	0.005604	1.10E-72
rs4418728	10	94839724	G	T	0.549296	0.998989	-0.02406	0.002728	1.10E-18
rs11231698	11	63877163	C	T	0.945696	0.986634	0.046951	0.006031	7.00E-15
rs79719909	12	20474706	A	G	0.794547	0.977208	-0.01872	0.003388	3.30E-08
rs6489191	12	1.23E+08	A	G	0.415161	1	-0.01658	0.002765	2.00E-09
rs7956959	12	1.25E+08	C	T	0.880147	0.99506	-0.02507	0.004168	1.80E-09
rs9513116	13	29016715	G	A	0.679419	0.98658	0.017647	0.002923	1.60E-09
rs261290	15	58678720	T	C	0.346128	0.996264	0.025009	0.00287	3.00E-18
rs1077834	15	58723479	T	C	0.779062	0.995244	-0.02278	0.003279	3.80E-12
rs36060036	16	20361950	C	T	0.834888	0.994847	0.027024	0.003673	1.90E-13
rs56228609	16	56987765	C	T	0.68325	0.995814	-0.02783	0.002932	2.30E-21
rs2925979	16	81534790	T	C	0.301454	1	-0.02147	0.002973	5.20E-13
rs2229611	17	41063466	T	C	0.214629	1	-0.02415	0.003276	1.70E-13
17:42300634_CAAAAA_C	17	42300634	CAAAAA	C	0.616177	0.982143	-0.02467	0.002791	9.70E-19
rs2306828	17	43213772	A	C	0.164834	0.988568	-0.02401	0.003651	4.90E-11
rs4969142	17	76398304	G	A	0.510431	0.99707	-0.01551	0.002703	9.50E-09
rs7412	19	45412079	C	T	0.917544	1	-0.03758	0.004889	1.50E-14
rs6088669	20	30177582	C	A	0.843559	0.997585	-0.02721	0.003746	3.70E-13

SNP = single nucleotide polymorphism. CHR = chromosome. SE = standard error.

*Supplementary Table 8. Results of cis-only pQTLs MR analysis for serum protein level (log2) on outcomes of interest (2-hour post OGTT glucose (mmol/L), fasting glucose(mmol/L), fasting insulin(mmol/L), and HbA1c (%)).*

Outcome	N SNPs	Effect estimate	SE	p-value	Q heterogeneity	Q p-value
2-hour post OGTT glucose	3	-0.1285	0.1150	0.2640	0.9401	0.6250
Fasting glucose	3	-0.0203	0.0346	0.5580	3.8401	0.1466
Fasting insulin	3	-0.0681	0.0277	0.0139	0.9232	0.6303
HbA1c	3	-0.0357	0.0319	0.2626	6.1030	0.0473

Supplementary Table 9. Results of cis- and trans-combined pQTLs MR analysis for outcomes of interest (2-hour post OGTT glucose, fasting glucose, fasting insulin, and HbA1c).

Outcome	Method	N SNPs	Effect estimate	SE	p-value	Q heterogeneity	Q p-value	Intercept	Intercept p-value
2-hour post OGTT glucose	IVW	39	-0.3722	0.0998	$1.92 \times 10^{-4}$	128.8568	$8.22 \times 10^{-12}$		
2-hour post OGTT glucose	MR Egger	39	-0.0455	0.1957	0.8174			-0.0115	0.0628
2-hour post OGTT glucose	Weighted median	39	-0.1163	0.0916	0.2039				
2-hour post OGTT glucose	Simple mode	39	-0.2194	0.1702	0.2051				
2-hour post OGTT glucose	Weighted mode	39	-0.1241	0.0831	0.1436				
Fasting glucose	IVW	39	-0.0307	0.0358	0.3909	349.5152	$5.59 \times 10^{-52}$		
Fasting glucose	MR Egger	39	-0.0327	0.0749	0.6645			$6.89 \times 10^{-5}$	0.9758
Fasting glucose	Weighted median	39	-0.0292	0.0202	0.1488				
Fasting glucose	Simple mode	39	-0.1040	0.0385	0.0102				
Fasting glucose	Weighted mode	39	-0.0175	0.0221	0.4342				
Fasting insulin	IVW	39	-0.0870	0.0559	0.1188	666.0874	$9.63 \times 10^{-116}$		
Fasting insulin	MR Egger	39	0.0354	0.1126	0.7547			-0.0043	0.2195
Fasting insulin	Weighted median	39	-0.0657	0.0255	0.0100				
Fasting insulin	Simple mode	39	-0.0280	0.0428	0.5164				
Fasting insulin	Weighted mode	39	-0.0347	0.0328	0.2972				
HbA1c	IVW	39	-0.0485	0.0155	0.0017	115.4975	$9.53 \times 10^{-10}$		
HbA1c	MR Egger	39	-0.0610	0.0319	0.0638			0.0004	0.6565
HbA1c	Weighted median	39	-0.0513	0.0159	0.0013				
HbA1c	Simple mode	39	-0.0522	0.0344	0.1380				
HbA1c	Weighted mode	39	-0.0465	0.0213	0.0351				

*Supplementary Table 10: SNPs and their prior associations*

SNP	CHR	BP	ALLELE1	ALLELE0	A1FREQ	BETA	SE	P	Phenoscaner
rs36060036	16	20361950	C	T	0.835	0.027	0.004	1.9E-13	Blood pressure,
rs6088669	20	30177582	C	A	0.844	-0.027	0.004	3.7E-13	Blood pressure,
rs28588745	8	10647044	A	T	0.793	0.047	0.003	8.2E-45	Body fat percentage
rs2306828	17	43213772	A	C	0.165	-0.024	0.004	4.9E-11	Body mass,
rs614008	6	31840794	C	T	0.421	0.017	0.003	3.0E-10	Body mass, Hip circumference, blood pressure
rs75104038	6	34190104	G	A	0.938	0.032	0.006	1.7E-08	Body mass, Waist circumference
rs9513116	13	29016715	G	A	0.679	0.018	0.003	1.5E-09	CAD
rs56228609	16	56987765	C	T	0.683	-0.028	0.003	2.3E-21	CAD, HDL,
rs261290	15	58678720	T	C	0.346	0.025	0.003	3.0E-18	Cholesterol
rs7412	19	45412079	C	T	0.918	-0.038	0.005	1.5E-14	Cholesterol, APOB, CAD, Weight, MI, Waist circumference, BMI
rs1077834	15	58723479	T	C	0.779	-0.023	0.003	3.7E-12	Cholesterol, Triglycerides,
rs8176747	9	136131315	C	G	0.938	-0.101	0.006	1.1E-72	Chronic kidney disease
rs1899951	3	12394840	C	T	0.877	-0.039	0.004	6.3E-21	Fat mass, diabetes type 2, BMI, WHR, insulin sensitivity
rs1064627	6	30698541	A	G	0.800	-0.020	0.003	5.1E-09	Fat mass/fat-free mass, blood pressure, hip circumference, cholesterol, diabetes
rs231492	17	41978756	G	T	0.916	0.155	0.005	4.1E-218	HDL
rs2925979	16	81534790	T	C	0.301	-0.021	0.003	5.2E-13	HDL, WHR, Diabetes type 2, triglycerides, BMI
rs11231698	11	63877163	C	T	0.946	0.047	0.006	7.0E-15	Hypertention
rs4418728	10	94839724	G	T	0.549	-0.024	0.003	1.1E-18	Triglycerides, body fat percentage
rs1260326	2	27730940	T	C	0.391	-0.021	0.003	6.5E-14	Triglycerides, cholesterol, CRP, glucose, fat mass, alcohol intake, weight and more
rs2980888	8	126507308	T	C	0.300	-0.021	0.003	2.3E-12	Triglycerides, cholesterol, fat percentage, hip circumference, CAD
rs998584	6	43757896	C	A	0.521	0.040	0.003	5.3E-48	WHR, Hip circumference, Triglycerides, Cholesterol, BMI