

1 **SYNCHRONOUS GROUP-BASED TELE-EXERCISE VERSUS**
2 **COMMUNITY-BASED INTERVENTIONS: EFFECTS ON**
3 **PHYSICAL FUNCTIONING AND ADHERENCE IN OLDER**
4 **ADULTS AT RISK OF FALLS - A RANDOMIZED CONTROLLED**
5 **TRIAL**

6 **ABSTRACT**

7 **Objective** This study aimed to assess the effectiveness of synchronous group-based
8 tele-exercise (TE) in improving physical functioning and exercise adherence among
9 older adults at risk of falls, comparing it to a community-based group (CB).

10 **Methods**

11 91 community-dwelling older adults at risk of falls were recruited from 10 community
12 centres in Hong Kong. Participants were randomly assigned to either the TE or CB
13 group based on their community centre. Both groups received modified Otago exercise
14 training for three months, with the difference in the mode of delivery (tele-exercise at
15 home or face-to-face at a community centre). The primary outcomes assessed Fall
16 Efficacy Scale (FES-I), 6-meter walk test, Timed Up and Go test (TUG), Berg Balance
17 Scale (BBS), Functional Reach Test (FRT), Appendicular Skeletal Muscle Mass Index
18 (ASMI) and percentage of body fat. Secondary outcomes included exercise adherence,
19 dropout rate, and Physical Activity Enjoyment Scale (PACES). The primary analysis
20 was conducted using a modified intention-to-treat approach, utilizing all available data.

21 **Results**

22 Both groups showed significant improvements in FES-I, 6-meter walk test, TUG, and
23 BBS over time ($p < 0.05$). High adherence rates (TE: 85%; CB: 90%), low dropout rates
24 (TE: 4.4%; CB: 0%), and high PACES scores (TE: 108.6 ± 9.3 ; CB: 108.2 ± 7.9) were
25 observed in both groups. TE and CB were comparable in most outcomes at baseline
26 and at three months ($p > 0.05$). No serious adverse events were reported.

27

28 **Conclusions**

29 The findings revealed synchronous group-based tele-exercise training demonstrated
30 comparable effectiveness to face-to-face community training in terms of physical
31 functioning, adherence, and enjoyment.

32 **Trial registration number** ChiCTR2200063370

33 **Keywords:** synchronous tele-exercise, falls, older adults, exercise adherence,

34

1 **SYNCHRONOUS GROUP-BASED TELE-EXERCISE VERSUS**
2 **COMMUNITY-BASED INTERVENTIONS: EFFECTS ON**
3 **PHYSICAL FUNCTIONING AND ADHERENCE IN OLDER**
4 **ADULTS AT RISK OF FALLS - A RANDOMIZED CONTROLLED**
5 **TRIAL**

6

7 **SUMMARY BOX**

8 What is already known on this topic - summarise the state of scientific knowledge on
9 this subject before you did your study and why this study needed to be done

- 10 • Community-based exercise programme has been proven effective in reducing
11 the risk of falls and preventing fall-related injuries.
12 • Tele-exercise presents a viable solution to address the challenges faced by
13 community- dwelling older adults at risk of falls such as limited exercise space
14 in the community and travel constraints.

15 Currently, there is a plethora of published studies on the effectiveness of tele-exercise.
16 However, these studies are primarily small-scale, non-randomized, pilot feasibility
17 studies, resulting in a generally low-to-moderate methodological quality of evidence.

18 What this study adds - summarise what we now know as a result of this study that we
19 did not know before

- 20 • This is the first rigorously designed RCT with sufficient statistical power to
21 investigate the effectiveness of the tele-exercise group compared with the
22 community-based group.
23 • Synchronous group-based tele-exercise training demonstrated comparable
24 effectiveness to face-to-face community training in terms of physical
25 functioning, adherence, and enjoyment.
26 • Tele-exercise training proved to be safe, feasible and enjoyable for older adults
27 at risk of falls, including those with diverse age ranges (65-91), educational
28 backgrounds, and technological proficiency.

29 How this study might affect research, practice or policy - summarise the implications
30 of this study

- 31 • Incorporating well-designed tele-exercise programmes with professional
32 support and technical assistance is crucial to effectively engage older adults,
33 promote their physical functions, improve exercise adherence and reduce the
34 risk of falls.
35 • These findings will inform future research on tele-exercise interventions, guide
36 the implementation of tele-exercise programmes in practice, and influence
37 policy decisions regarding the application of tele-exercise for older adults at risk
38 of falls.

39

1 INTRODUCTION

2 Falls among older people are a significant concern, with a global prevalence of 26.5%.¹
3 Falls can lead to substantial costs, ranging from 2,044 to 25,955 USD per fall victim,
4 1,059 to 10,913 USD per fall event, and 5,654 to 42,840 USD per fall-related
5 hospitalization.² Hospitalizations due to falls are five times more common than those
6 resulting from other injuries.³ Falls also have adverse consequences such as hip
7 fractures, rehospitalization, extended hospital stays, and increased mortality rates.^{4 5}
8 Therefore, implementing preventive strategies is vital in mitigating age-related
9 impairments and promoting independence in older adults.

10 Traditional exercise interventions have shown promise in reducing fall risk and
11 preventing fall-related injuries.⁶ However, older adults face significant challenges due
12 to limited exercise space in the community and travel constraints,⁷ especially for those
13 who are susceptible to falls. To address these challenges, tele-exercise has emerged as
14 a viable alternative. Tele-exercise encompasses various forms, including video, audio,
15 or text communication technologies, delivered synchronously or asynchronously.^{8 9}
16 Chan, *et al*¹⁰ reported promising results in reducing falls among community-dwelling
17 older adults through telehealth combined with exercise interventions and highlighted
18 the safety, feasibility, and acceptability of telehealth interventions. The presence of real-
19 time guidance and supervision of video conferencing-based online exercise
20 programmes was found to be highly beneficial for older adults.⁷ Granet, *et al*¹¹
21 demonstrated that a higher ratio of live training resulted in greater improvements in
22 muscle function and a lower drop-out rate compared to a higher ratio of recorded
23 sessions, emphasizing the relevance of synchronous delivery of tele-exercise.

24 The existing studies on synchronous tele-exercise for community dwelling older adults
25 and older adults at risk of falls have limitations. They predominantly focused on older
26 adults who already had an internet connection and the necessary devices for online
27 exercise programmes as inclusion criteria.¹¹⁻¹³ Granet, *et al*¹¹ noted that the study
28 samples were highly educated, with over 67% of participants having attained a
29 university level of education, which may limit the generalizability of the results to the
30 broader population. In addition, the lack of proper sample size estimation in some
31 studies also highlights the need for improved study quality.^{11 13}

32 In a recent systematic review conducted by Gamble, *et al*⁸, the methodological quality
33 of the evidence from 26 randomized controlled trials about the effectiveness of
34 telerehabilitation such as tele-exercise for community dwelling older adults was
35 critiqued. Among the included trials, five¹⁴ specifically targeted older adults with an
36 increased risk of falls.¹³⁻¹⁷ It is noteworthy that four of these studies^{13 14 16 17} exhibited
37 a high risk of bias based on the revised Cochrane Risk-of-Bias 2 tool,¹⁸ emphasizing
38 the necessity for more rigorous and controlled investigations in this particular field.
39 Interestingly, only two of these studies incorporated a videoconferencing setup^{13 14},
40 which is an essential component of synchronous tele-exercise.

41 Therefore, the aim of the current study was to address these methodological limitations
42 by investigating the effectiveness of synchronous tele-exercise (TE) on physical
43 functioning and exercise adherence among community-dwelling older adults at risk of
44 falls in comparison with a community-based group (CB).

1 **METHODS**

2 **Study design**

3 This was a 3-month single-blind, randomized controlled trial (RCT) comparing
4 synchronous group-based tele-exercise training (TE) with face-to-face community-
5 based exercise training (CB) for community-dwelling older people at risk of falls. More
6 detailed information, including details of the outcome measures and intrinsic
7 differences between the two groups, can be obtained from our study protocol.¹⁹
8 Reporting follows the CONSORT 2010 statement.²⁰

9

10 **Research ethics**

11 Approval for the study was granted by the Ethics Committee of the College of
12 Professional and Continuing Education (RC/ETH/H/0029). The study was registered in
13 the Chinese Clinical Trial Registry (ChiCTR2200063370) on 5 September 2022. In
14 accordance with the WMA declaration of Helsinki, informed consent was obtained from
15 all participants.

16

17 **Sample size calculation**

18 Sample size calculations were conducted using G*Power software, assuming an effect
19 size of 0.367 based on a previous study by Benavent-Caballer, *et al*²¹ that focused on
20 the Time Up and Go Test. With a confidence level of 95%, a power of 80%, and an
21 estimated attrition rate of 20% (including mortality)²², a minimum sample size of 92
22 participants was determined to be necessary.

23

24 **Participants and eligibility criteria**

25 The participants were recruited from 10 elderly community centres located in various
26 locations across Hong Kong. The inclusion and exclusion criteria are listed in table 1.

27

28 **Randomization and blinding**

29 Ten community centres will be coded and then randomly assigned to either the tele-
30 exercise group (TE) or community-based group (CB) with a 1:1 allocation in a
31 concealed, blinded fashion to minimize cross-over contamination between the groups.
32 Randomization procedure and allocation sequence was supervised by an independent
33 assistant. The project assistant enrolled participants in the community centres. Trained
34 outcome assessors, who were blinded to group assignments, performed the assessments.
35 Participants were aware of their unique training modes but had no contact with
36 participants from other groups due to different locations. Instructors and researchers
37 were not blinded and were instructed not to discuss group assignments with the
38 participants and outcome assessors.

39

1 **Intervention programmes**

2 Both interventions have been described in detail in the published protocol.¹⁹ All
3 participants engaged in exercise training sessions three times a week for a duration of
4 three months, either through tele-exercise (TE) or face-to-face at the community centres
5 (CB). The participants in both groups were divided into classes consisting of 8 to 11
6 individuals. The exercise programme followed a structured format, which included a
7 warm-up, Otago exercise programme (OEP),²³ "workouts on-demand," and a cool-
8 down.

9 *Intervention- tele-exercise group*

10 For the TE group, the exercise programme was delivered via Zoom (Zoom Video
11 Communications Inc.) in the participants' homes, allowing for real-time online
12 communication with the participants. Participants without suitable devices were
13 provided with all the necessary devices.

14

15 *Safety measures and technological support for the tele-exercise group*

16 Prior to the intervention, home visits were conducted to ensure a safe home
17 environment, assist with setting up online communication applications, and optimize
18 the positioning of the tablet, chair, and participant for effective visual interaction. The
19 first lesson took place at the community centre to familiarize participants with exercise
20 safety rules and the OEP, followed by online training sessions. Participants were
21 reminded to turn on their tablets and Wi-Fi before each class so that we could control
22 their tablet remotely to promptly address technical issues. Participants performed all
23 OEP exercises, starting at the beginner level with support for standing and walking.
24 Camera and microphone activation were required to enhance interaction, with regular
25 progress monitoring by the research team. Participants were reminded to wear
26 appropriate exercise gear and maintain hydration.

27

28 *Intervention- community-based group*

29 For the CB group, the exercise classes were arranged at the community centres based
30 on the residence of the recruited participants. The training programme was the same as
31 that of TE group, except that it was conducted face-to-face at the community centre.

32

33 **Outcome measures**

34 Baseline assessments and post-intervention assessments at the three-month were
35 conducted by trained and blinded assessors at the community centres. Participants
36 received HK\$100 gift vouchers at 3 months for completing the training and assessments.
37 In addition to the assessments, demographic information and background variables such
38 as sex, age, and history of falls were collected. Cognitive function at baseline was
39 evaluated using the Mini-Mental State Examination (MMSE).²⁴ The Chinese version of
40 the Physical Activity Scale for the Elderly (PASE-C) was utilized to assess the physical
41 activity level of older adults.²⁵

1 *Primary outcomes*

2 Fall efficacy was evaluated using the Chinese version of the Fall Efficacy Scale-
3 International (FES-I).²⁶ Participants were asked to proactively record any instances of
4 falling and any adverse events in their diaries. The primary outcome measures also
5 included SARC-CalF, grip strength assessments for both the right and left hands, a 6-
6 meter walk test, a timed up and go test (TUG), the Berg Balance Scale score (BBS), the
7 functional reach test (FRT), appendicular skeletal muscle mass index (ASMI)
8 percentage of body fat and Physical Activity Scale for the Elderly-Chinese (PASE-C).
9 The definition of possible sarcopenia, sarcopenia and severe sarcopenia were based on
10 Chen, *et al*²⁷

11

12 *Secondary outcomes*

13 The secondary outcomes evaluate older people's involvement and experience of the
14 exercise programmes including exercise adherence, dropout rate and Physical Activity
15 Enjoyment Scale (PACES).²⁸ The exercise adherence rate was defined as the percentage
16 of attended classes out of the total 36 classes.

17

18 **Data analysis**

19 Baseline characteristics including age, adherence rate, and dropout rate, were analyzed
20 using an independent t-test. An intention-to-treat analysis, following the guidelines for
21 randomized controlled trials,²⁹ was employed, including all originally randomized
22 participants in the study. The normal distribution of all measurement variables was
23 confirmed using the Kolmogorov-Smirnov one-sample test. For the outcome measures,
24 a two-way repeated measures ANOVA was conducted, taking into account the group
25 (TE and CB) and the interaction between group and time. The differences between
26 categorical variables such as education level and % participants with possible
27 sarcopenia were analyzed by Chi-square test. A significance level of 0.05 was set for
28 all tests. The statistical analysis was carried out using SPSS, version 26.0 (IBM Co.,
29 Armonk, NY).

30 Sensitivity analyses were conducted to validate the statistical findings. These included
31 utilizing multiple imputation to address missing data, which confirmed consistent
32 results with the original analysis.

33 **RESULTS**

34 **Flow of participants**

35 During the screening period from October 2021 to June 2022, a total of 212 older adults
36 were identified for preliminary screening. Out of these, 120 individuals were excluded
37 based on the inclusion and exclusion criteria of the study. Eventually, 91 participants
38 successfully met the inclusion criteria. The TE group comprised 45 participants, while
39 the CB group consisted of 46 participants (Figure 1).

40

1 **Participant characteristics**

2 The baseline characteristics of the participants did not show significant differences
3 between the CB and TE groups, as summarized in Table 2.

4

5 **Technological accessibility and technological support for the TE participants**

6 In terms of smartphone ownership, the TE group exhibited a higher percentage at 84.4%,
7 while the CB group reported 71.7% ($p < 0.05$). Similarly, in terms of home Wi-Fi
8 ownership, the TE group also had a higher percentage at 71.1%, compared to the CB
9 group at 58.7%, but this difference did not reach a significant level. Additionally, only
10 17.8% of TE participants had previous experience with tele-exercise (Table 2).

11 Baseline home visits were conducted for all participants before the intervention, with
12 an average of 3.7 visits per participant. Additional visits addressed technological issues
13 and provided support, totaling 260 visits (including baseline visits) for the 45
14 participants in the TE group, averaging 5.8 visits per participant (range: 0-12 visits).

15

16 **Primary outcomes- falls incidence, fall risks and sarcopenia status**

17 There was no significant difference between the groups in the falls incidence. Prior to
18 the intervention, the average number of falls per participant was 1.57 ± 0.83 over the
19 past 12 months in the CB and 1.78 ± 0.77 in the TE. The majority of falls were classified
20 as minor incidents (CB: 97.2% vs TE: 97.8%).³⁰ During the intervention, the CB group
21 reported no falls, whereas one participant in the TE group reported experiencing a fall
22 outside of the lessons, which resulted in a minor injury (Table 3). The FES-I scores
23 improved significantly in both groups, indicating a reduction in fear of falling. ($p < 0.05$)
24 (Table 4).

25 There were no significant differences between the two groups or between the two time
26 points in the proportion of participants with possible sarcopenia, sarcopenia, and severe
27 sarcopenia. However, there was a trend towards a decrease in participants with possible
28 sarcopenia and severe sarcopenia in the CB group over time (Table 3).

29

30 **Primary outcomes- physical function and activity level**

31 Although no significant difference was observed between the CB and TE interventions,
32 most outcomes showed improvement after the 3-month intervention (Table 4). The 6-
33 meter walk test, the TUG scores and the BBS scores improved significantly in both
34 groups over the 3-month period ($p < 0.05$).

35 SARC-CalF scores and the FRT scores demonstrated significant improvement in the
36 CB group compared with its baseline ($p < 0.05$). Handgrip strength, measured in all
37 participants, did not show significant changes between or within groups. However,
38 within the female subgroup, CB participants showed a significant increase in handgrip
39 strength compared with its baseline ($p < 0.001$). Activity levels, assessed by the Physical

1 Activity Scale for the Elderly (PASE-C), significantly improved in both groups
2 ($p < 0.001$).

3

4 **Primary outcomes- body composition changes among female participants**

5 Due to the different standards for body composition between genders and the significant
6 proportion of female participants in both groups, the body composition changes among
7 female participants (CB $n=41$; TE $n=39$) were analyzed and presented. The results
8 showed minimal differences between the CB and the TE (Table 5). Calf circumference,
9 BMI and appendicular skeletal muscle mass index (ASMI) remained relatively stable
10 in both groups over the 3-month period. However, weight and % body fat exhibited a
11 significant decrease in the TE group ($p < 0.05$).

12

13 **Secondary outcomes- adherence rate, dropout rate and PACES**

14 Adherence rates were high in both groups, with the CB group averaging 90.0%
15 adherence and the TE group averaging 85.0%. The dropout rate was 0% in the CB group,
16 while the TE group had a dropout rate of 4.4%. The scores on the PACES were 108.6
17 in the CB group and 108.2 in the TE group, out of a total of 126 (Table 2).

18

19 **Adverse events reporting**

20 One participant in the TE group reported general tiredness and pain during screening
21 phase, but her doctor found no issues after thorough examinations and tests. Living
22 alone and rarely engaging in social activities, she insisted on participating in our
23 programme to maintain her physical functioning.

24 During the tele-exercise class, she reported tiredness and knee and lower back pain at
25 the beginning. The instructor advised her to rest when needed and practice at an easier
26 level. With the frequent reminders to lower the intensity, her problems subsided. She
27 completed the programme with an adherence rate of 72.2%.

28 **DISCUSSION**

29 **Key findings**

30 To our knowledge, this study is the first rigorously designed RCT with sufficient
31 statistical power to investigate the effectiveness of the synchronous group-based tele-
32 exercise compared with the community-based group in older adults at risk of falls. In
33 contrast to previous trials^{14 15}, our study includes a comprehensive range of outcomes
34 such as fall efficacy, fall incidence, sarcopenia status, physical function, activity level,
35 body composition, exercise adherence, dropout, enjoyment, adverse events,
36 technological accessibility and their education backgrounds. The study findings
37 indicate that both the CB and TE interventions were effective in decreasing fear of
38 falling, and enhancing physical function including mobility, balance and gait speed,
39 among the participants. These results regarding the effectiveness of the modified OEP

1 are consistent with the findings of systematic review conducted by Martins, *et al*³¹ and
2 Wu, *et al*³² The positive outcomes can be attributed to the fact that our intervention
3 adhered to the recommended effective programme requirements identified by Wu, *et*
4 *al*³², with an average duration of 45 minutes, three times per week for 12 weeks.

5 The study employed synchronous tele-exercise, utilizing video-linked appointments
6 supervised by healthcare professionals, and emphasized three main strategies: self-
7 efficacy, self-confidence, and empowerment, to enhance the exercise experience of the
8 participants.^{19 33 34} Immediate feedback after each class facilitated timely personalized
9 guidance, addressing participants' concerns. All classes were led by the same
10 knowledgeable instructor who had over 10 years of experience in teaching older adults
11 and people with chronic diseases. Our training programme effectively mitigated the
12 challenges associated with remote exercise sessions, as evidenced by the lack of
13 significant differences between the two groups in physical function.

14

15 The study demonstrated high adherence and enjoyment among participants in both the
16 CB and TE interventions, with low dropout rates and high PACES scores. CB achieved
17 90% adherence and TE reached 85%, surpassing the approximately 70% reported in a
18 systematic review of community-based exercise programmes.³⁵ As recommended, an
19 80% adherence rate is considered feasible.³⁶ Consistent with previous studies,
20 community-dwelling older adults expressed high satisfaction with online exercise
21 training.³⁷⁻³⁹ The success of the tele-exercise programme can be attributed to factors
22 such as professional support, individualized technology orientation sessions, and
23 standby technical assistance.^{39 40} Our programme also provided prompt technical
24 assistance, averaging 5.8 home visits per participant throughout the 36 lessons,
25 accounting for approximately 16.1% of the sessions. These measures ensured prompt
26 problem-solving, exercise safety, and the absence of serious adverse event, indicating a
27 high margin of exercise safety with the synchronous online delivery of modified OEP.

28

29 **Upper body strength improvements**

30 While the OEP primarily focuses on strength and balance exercises for the lower
31 extremities, it has been shown to prevent a decline in handgrip strength when compared
32 to no intervention control⁴¹ and conventional care.⁴² However, recent meta-analysis
33 reviews did not find clear benefits of the OEP on upper body strength in older adults.³²
34 These analyses did not analyze the changes based on gender, as there are gender
35 differences in the norm of handgrip strength. Nevertheless, our study observed
36 significant improvements in upper body strength exclusively among female participants
37 within the CB group. The primary purpose of engaging the upper limb muscles was to
38 provide support and maintain balance during exercises. While this aspect may appear
39 minor, it holds significant importance for older adults who are at risk of falls. Previous
40 research has demonstrated that even small volumes of resistance exercise, performed
41 briefly and infrequently, can lead to substantial increases in muscular strength.⁴³ The
42 improvement in handgrip strength in the CB may explain the decreasing trend of
43 possible sarcopenia and severe sarcopenia among the CB participants over time.

1

2 **Impact on body weight and fat**

3 Overall, the impact of the interventions on body composition parameters was modest,
4 with slight variations between CB and TE groups. However, subgroup analysis revealed
5 that female participants in the TE group experienced a significant decrease in weight
6 and body fat percentage after three months, compared to baseline. This aligns with
7 Duckham, *et al*⁴⁴, who demonstrated fat mass reduction with an Otago-based exercise
8 programme. The significant decrease in body fat percentage in our study may be due to
9 increased physical activity levels, as indicated by higher PASE-C scores. Regular
10 participation in Otago-based exercise can serve as an effective energy expenditure
11 method for older adults at risk of falls, particularly for those who do not have
12 established exercise habits and maintain very low activity levels. Another study found
13 improvements in lower limb muscle mass with resistance exercise delivered via video
14 conferencing.⁴⁵ Unlike our programme, Hong, *et al*⁴⁵ also included nutrition education,
15 which is important for preventing and managing sarcopenia.⁴⁶ This difference in
16 approach may explain the significant increase in muscle mass observed in their study
17 but not in ours.

18 **Bridging technological and education gaps in tele-exercise**

19 Most tele-exercise studies primarily recruited individuals with existing internet access
20 at home, favoring those who were technologically competent.^{11-13 47} A strength of our
21 study was that technological competence and device ownership were not inclusion
22 criteria. We provided participants with the necessary devices, welcoming individuals
23 with various technological backgrounds. Our sample consisted of less educated
24 individuals, with over 50% completed primary education or below, and only one TE
25 participant completing university. Our findings indicated that older adults, regardless
26 of education and technological proficiency, could enjoy tele-exercise when
27 accompanied by well-designed programmes, technical support, and timely feedback.
28 Furthermore, the utilization of commonly available telecommunications software in the
29 trial, instead of custom-built systems¹³, enhanced the generalizability of the findings.⁴⁸

30

31 **Technological exposure and potential selection bias**

32 Our findings revealed a significantly higher rate of smartphone ownership in the TE
33 group (84.4%) compared to the CB (71.7%) ($p < 0.05$), indicating greater technological
34 exposure among TE participants. Additionally, the TE group had a higher proportion of
35 participants with Wi-Fi at home (72.1% vs. CB: 58.7%, N.S.). The TE group also
36 exhibited slightly higher levels of education, suggesting potential selection bias towards
37 technologically experienced and higher-educated individuals. It is important to note that
38 three participants declined participation before the intervention phase due to family
39 objections and concerns about unsafe home environments. This highlights the presence
40 of skepticism towards the use of tele-exercise among older adults, indicating that some
41 individuals and their families may have reservations about engaging in exercise
42 programmes remotely.

1 **Limitations**

2 One noteworthy concern with tele-exercise is the potential impact on the proper
3 execution of exercises, especially for older adults with specific health conditions or
4 mobility issues. Due to intrinsic differences between the delivery modes of CB and TE,
5 the instructor could only provide verbal corrections for participants' movements,
6 without the ability to physically assist them.¹⁹ It is worth noting that conducting tele-
7 exercise sessions in the limited space of most Hong Kong residences presents
8 challenges. Census and Statistics Department⁴⁹ indicate that the median floor areas for
9 domestic households and individuals in Hong Kong are 40 and 16 m², respectively. This
10 spatial constraint could impact the ability to capture a full view of participants while
11 they were standing during tele-exercise sessions. To overcome this issue, we made
12 efforts during home visits to find the optimal placement and angle for tablets with built-
13 in cameras to maximize visibility during the exercise sessions. To ensure understanding
14 of the exercise programme and promote exercise safety, all participants were required
15 to attend the first class at the community centre. This allowed for in-person instructions
16 and reminders before transitioning to tele-exercise sessions.¹⁹ It is important to note
17 that the success of this program may not be transferable to other exercises that involve
18 more dynamic movements such as standing, sitting, and lying down, within a single
19 lesson. From a technological standpoint, it would be beneficial for the technology
20 company to explore the possibility of incorporating tablets with wide-angle front
21 cameras or establishing a seamless connection between a tablet and an external camera
22 without any delays. These enhancements would help improve visibility during tele-
23 exercise sessions.


24

25 **Conclusion**

26 To our knowledge, this is the first rigorously designed RCT with sufficient statistical
27 power and a comprehensive set of outcomes to explore the effectiveness of the tele-
28 exercise group compared with the community-based group. Both CB and TE
29 interventions yielded positive outcomes, high participant satisfaction, and no serious
30 adverse events. These results highlighted the effectiveness of these approaches for falls
31 prevention and physical improvement in older adults at risk of falls, including those
32 with varying education levels and technological competence, even among those who
33 were new to tele-exercise. The study provides valuable insights and potential strategies
34 for safely and effectively delivering synchronous tele-exercise training among
35 community-dwelling older adults at risk of falls. These findings have significant
36 implications for enhancing older adults' health and well-being, especially when in-
37 person interventions are challenging or inaccessible.

38

1 TABLES AND FIGURE



24 Figure 1 Flow of Participants.

25

1

2

Inclusion criteria	Exclusion criteria
<ul style="list-style-type: none">• permanent residents of Hong Kong• aged 65 years or above living independently at home• diagnosed with possible sarcopenia and/or having had either a fall in the past year or fear of falling with a score of ≥ 20 points on FES-I• able to speak and listen to Cantonese• able to walk without mobility aids in their home	<ul style="list-style-type: none">• diagnosed with dementia or cognitive impairment, Mini-Mental State (MMSE) < 25• severe vision impairments that do not permit the use of a tablet, reading of the exercise booklet, and completing the activity diaries• unable to pass the preparticipation Health Screening of the ACSM and the subsequent medical clearance• regular physical activities in the past 6 months based on the definition of WHO⁵⁰

3 Table 1 Inclusion and exclusion criteria (adopted from Chan, *et al*¹⁹)

4

5

	Community-based (CB)	Tele-exercise (TE)
Baseline Characteristics		
n (excluding dropout)	46	45
Age (Range)	77.3 ± 6.7 (65-93)	74.8 ± 6.2 (65-91)
Proportion of female participants	89.1% (41)	86.6% (39)
Weight (kg)	56.6 ± 12.0	58.8 ± 11.6
Height (cm)	150.3 ± 6.3	153.6 ± 8.2
BMI (kg/m ²)	25.5 ± 0.7	24.2 ± 0.7
Mini Nutritional Assessment (MNA)	24.9 ± 0.3	25.1 ± 0.3
Mini Mental State Examination (MMSE)	27.59 ± 1.89	27.84 ± 1.78
Education Level		
Junior primary or below	47.8%	22.2%
Senior primary	21.7%	35.6%
Junior Secondary	21.7%	17.8%
Senior Secondary	8.6%	22.2%
University or above	0%	2.2%
Technological accessibility		
Smartphone ownership	71.7%	84.4%*
Home Wi-Fi ownership	58.7%	71.1%
Previous experiences with tele-exercise	N/A	17.8%
Participant involvement and experience		
Dropout during intervention	0%	4.4% (2)
Adherence rate	90.0 ± 7.1 %	85.0 ± 11.8%
Physical Activity Enjoyment Scale (PACES)	108.6 ± 9.3%	108.2 ± 7.9%

1 **p*<0.05 compared with CB

2 Table 2. Demographic characteristics and satisfaction of the 3-month training
3 programme

1
2
3
4
5
6

	Baseline (last 12 months before intervention))		3 month	
Fall incidence				
	CB	TE	CB	TE
Average no. of falls per participant	1.57 ± 0.83	1.78 ± 0.77	0	0.02 ± 0.15
Minor injuries, %	97.2%	97.8%	0%	100%
Serious injuries, hospitalisation %	0.03%	2.5%	0%	0
Sarcopenia analysis				
	CB	TE	CB	TE
% of participants with Possible Sarcopenia	41.3%	33.3%	26.1%	35.6%
% of participants with Sarcopenia	39.1%	28.9%	39.1%	28.9%
% of participants with Severe Sarcopenia	28.3%	15.6%	13.0%	17.8%

7

8 *Possible sarcopenia: SARC-CalF \geq 11, handgrip strength M:<28 kg, F<18 kg

9 Sarcopenia: SARC-CalF \geq 11, ASMI M<7.0 kg/m², F<5.7 kg/m², handgrip strength
10 M:<28 kg; F<18 kg or 6-metre walk<1.0 m/s

11 Severe Sarcopenia: SARC-CalF \geq 11, ASMI M<7.0 kg/m², F<5.7 kg/m², handgrip
12 strength M:<28 kg; F<18 kg and 6-metre walk<1.0 m/s

13

14

15 Table 3 Falls incidence and sarcopenia analysis

Outcomes	Baseline ^a (mean ± SE)	3- month ^b (mean ± SE)	Mean difference (95% CI)	p-value Within Group
FES-I				
CB	36.41 ± 1.06	33.37 ± 1.16	-3.04 (-4.36 to -1.73)	<0.001
TE	34.98 ± 1.08	32.49 ± 1.17	-2.49 (-3.81 to -1.16)	<0.001
Mean difference (95% CI)	-1.44 (-4.44 to 1.57)	-0.88 (-4.14 to 2.38)		
<i>p-value Between Groups</i>	0.345	0.594		
SARC-CalF				
CB	8.04 ± 0.70	7.04 ± 0.64	-1 (-1.79 to -0.21)	<0.05
TE	8.56 ± 0.70	8.04 ± 0.65	-0.52 (-1.31 to 0.28)	0.2
Mean difference (95% CI)	0.51 (-1.45 to 2.48)	0.10 (-0.82 to 2.81)		
<i>p-value Between Groups</i>	0.606	0.278		
6-meter Walk (m/s)				
CB	0.69 ± 0.03	0.75 ± 0.23	0.06 (0.04 to 0.10)	<0.05
TE	0.68 ± 0.23	0.76 ± 0.04	0.08 (0.04 to 0.11)	<0.05
Mean difference (95% CI)	-0.01 (-0.08 to 0.07)	0.00 (-0.07 to 0.07)		
<i>p-value Between Groups</i>	0.871	0.955		
TUG				
CB	15.59 ± 1.16	13.88 ± 1.30	-1.71 (-3.05 to -0.38)	<0.05
TE	12.89 ± 1.17	11.15 ± 1.31	-1.74 (-3.10 to -0.39)	<0.05
Mean difference (95% CI)	-2.70 (-5.97 to 0.58)	-2.73 (-6.38 to 0.93)		
<i>p-value Between Groups</i>	0.105	0.142		
BBS				
CB	46.20 ± 1.09	49.74 ± 1.11	3.54 (1.40 to 5.68)	<0.01
TE	47.22 ± 1.10	50.67 ± 1.13	3.45 (1.28 to 5.61)	<0.01
Mean difference (95% CI)	1.03 (-2.06 to 4.11)	0.93 (-2.21 to 4.07)		
<i>p-value Between Groups</i>	0.510	0.557		
FRT (cm)				
CB	21.67 ± 1.04	24.41 ± 1.04	2.74 (0.66 to 4.81)	<0.05
TE	26.59 ± 1.05	26.75 ± 1.05	0.16 (-1.94 to 2.25)	0.881
Mean difference (95% CI)	4.92 (1.98 to 7.87)	2.34 (-0.58 to 5.26)		
<i>p-value Between Groups</i>	0.001	0.115		
Handgrip (kg) (all participants)				
CB	16.98 ± 4.51	18.18 ± 4.73	1.2 (-0.20 to 2.60)	0.092
TE	19.25 ± 6.28	19.81 ± 6.05	0.56 (-0.86 to 1.98)	0.434
Mean difference (95% CI)	2.27 (-0.00 to 4.55)	1.63 (-0.63 to 3.89)		
<i>p-value Between Groups</i>	0.050	0.155		
Handgrip (kg) (Female only)				
CB	16.14 ± 0.72	18.46 ± 0.79	2.32 (1.04 to 3.60)	<0.001
TE	17.95 ± 0.75	18.73 ± 0.82	0.78 (-0.53 to 2.10)	0.235
Mean difference (95% CI)	1.80 (-0.68 to 3.86)	0.27 (-1.98 to 2.52)		
<i>p-value Between Groups</i>	0.087	0.809		

PASE-C				
CB	122.02 ± 9.39	157.63 ± 9.12	35.61 (31.48 to 39.74)	<0.001
TE	119.51 ± 9.50	152.65 ± 9.23	33.14 (28.97 to 37.32)	<0.001
Mean difference (95% CI)	-2.51 (-29.05 to 24.03)	-4.98 (-30.73 to 20.78)		
<i>p-value Between Groups</i>	0.851	0.702		

1 FES-I: Fall Efficacy Scale; TUG: Timed Up and Go Test; BBS: Berg Balance Scale score; FRT: Functional Reach Test; PASE-C:
 2 Chinese version of the Physical Activity Scale for the Elderly;

3 Table 4. Comparison of fall risk, physical function and activity level at baseline and
 4 after the intervention

5

1

Outcomes	Baseline ^a (mean ± SE)	3- month ^b (mean ± SE)	Mean difference (95% CI)	p-value Within Group
Calf Circumference				
CB	33.98 ± 0.54	33.96 ± 0.54	-0.01 (-0.33 to 0.31)	0.940
TE	33.86 ± 0.56	33.92 ± 0.56	0.06 (-0.26 to 0.39)	0.698
Mean difference (95% CI)	-0.12 (-1.66 to 1.43)	-0.04 (-1.59 to 1.51)		
<i>p-value Between Groups</i>	0.881	0.959		
Weight				
CB	55.85 ± 1.8	55.48 ± 1.98	-0.37 (-2.52 to 1.77)	0.730
TE	57.53 ± 1.85	54.94 ± 2.03	-2.59 (-4.79 to -0.40)	<0.05
Mean difference (95% CI)	1.68 (-3.46 to 6.81)	-0.55 (-6.19 to 5.10)		
<i>p-value Between Groups</i>	0.518	0.848		
% Body fat				
CB	36.23 ± 1.04	36.69 ± 1.14	0.46 (-0.99 to 1.90)	0.532
TE	35.81 ± 1.07	33.78 ± 1.78	-2.03 (-3.51 to -0.55)	<0.05
Mean difference (95% CI)	-0.42 (-3.37 to 2.53)	-2.91 (-6.16 to 0.35)		
<i>p-value Between Groups</i>	0.778	0.079		
BMI				
CB	25.43 ± 0.75	24.94 ± 0.80	-0.49 (-1.59 to 0.61)	0.381
TE	24.99 ± 0.77	24.10 ± 0.82	-0.89 (-2.02 to 0.24)	0.122
Mean difference (95% CI)	-0.44 (-2.59 to 1.70)	-0.84 (-3.11 to 1.43)		
<i>p-value Between Groups</i>	0.683	0.463		
ASMI				
CB	5.74 ± 0.13	5.70 ± 0.21	-0.04 (-0.34 to 0.27)	0.815
TE	6.06 ± 0.14	6.25 ± 0.22	0.19 (-0.13 to 0.50)	0.237
Mean difference (95% CI)	0.32 (-0.05 to 0.70)	0.55 (-0.05 to 1.14)		
<i>p-value Between Groups</i>	0.092	0.070		

2 BMI: body mass index; ASMI: Appendicular skeletal muscle mass index ASMI

3 Table 5 Body composition changes among female participants (CB n=41; TE n=39)

4

1

2 **Author affiliations**

3 ¹College of Professional and Continuing Education, The Hong Kong Polytechnic
4 University, Hong Kong SAR, China

5 ²School of Management, Tung Wah College, Hong Kong SAR, China

6 ³Department of Rehabilitation Sciences, The Hong Kong Polytechnic University,
7 Hong Kong SAR, China

8 ⁴Hong Kong Telemedicine Association, Hong Kong SAR, China

9

10 **Acknowledgement**

11 The authors are grateful to the Centre for Ageing and Healthcare Management Research
12 (CAHMR), College of Professional and Continuing Education, The Hong Kong
13 Polytechnic University for coordination with the elderly services providers i.e. Sik Sik
14 Yuen. Thank Sik Sik Yuen for recruiting older people and all the participants for joining
15 this study. The authors would like to express their gratitude to Mr. Au Wing Ho for his
16 exceptional exercise teaching provided to all the participants. Additionally, the authors
17 extend their heartfelt thanks to the student helpers who offered technical support to the
18 participants.

19

20 **Author's contribution**

21 KOWC, PPY, BYFF, VTSL, FSFN, WCPF and TKCN conceived the study. KOWC
22 and JLCC wrote the main manuscript text. ISC collected the data. KOWC and ISC
23 performed statistical analysis. All authors reviewed the manuscript and approved the
24 final version of the manuscript.

25

26 **Funding**

27 The study was funded by the Health and Medical Research Fund (HMRF) under the
28 Health Bureau of the Government of the Hong Kong Special Administrative Region of
29 the People's Republic of China (05200028). The funder has no role on the study design,
30 protocol preparation, data collection and management, or future data analysis and
31 interpretation, writing of the report, and the decision to submit the report for publication.
32 They have no authority over any of these activities.

33

34 **Competing interests**

35 The authors declare that they have no competing interests.

36 **Patient and public involvement statement:**

1 Patients and/or the public were not involved in the design, or conduct, or reporting, or
2 dissemination plans of this research.

3 **Consent for publication** Obtained.

4 **Ethics approval and consent to participate**

5 The study conforms to the Declaration of Helsinki. The study involves human
6 participants and was approved by the Ethics Committee of the College of Professional
7 and Continuing Education, The Hong Kong Polytechnic University (RC/ETH/H/0029).
8 All participants provided informed consent to participate in the study prior to their
9 involvement.

10 **Availability of data and materials**

11 Data available upon reasonable request if permitted by the respective ethics vote.

12 **ORCID iDs**

13 Karly O.W. CHAN: 0000-0001-7098-519X

14 Peter P.M. YUEN: 0000-0002-0134-0141

15 Ben Y.F. FONG: 0000-0003-0909-0674

16 Vincent T.S. LAW: 0000-0001-8204-3925

17 Fowie S.F. NG: 0000-0002-6634-0345

18 Janet L.C. Lee: 0000-0001-8447-6855

19 I.S. CHEUNG: N/A

20 Tommy K.C. Ng: 0000-0002-7715-9533

21 Wilson C.P. FUNG: N/A

22 **References**

- 23 1. Salari N, Darvishi N, Ahmadipanah M, et al. Global prevalence of falls in the older adults:
24 a comprehensive systematic review and meta-analysis. *J Orthop Surg Res* 2022;17(1)
25 doi: 10.1186/s13018-022-03222-1
- 26 2. Heinrich S, Rapp K, Rissmann U, et al. Cost of falls in old age: a systematic review.
27 *Osteoporos Int* 2010;21:891-902.
- 28 3. Alexander BH, Rivara FP, Wolf ME. The cost and frequency of hospitalization for fall-
29 related injuries in older adults. *Am J Public Health* 1992;82(7):1020-23. doi:
30 10.2105/ajph.82.7.1020
- 31 4. Wang AQ-L, Ng BH-P, Cheung LP-C, Chin RP-H. Factors Affecting Mortality and
32 Hospital Admissions after Hip Surgery among Elderly Patients with Hip Fracture in
33 Hong Kong – Review of a Three-Year Follow-Up. *Hong Kong J Occup Ther*
34 2017;30(1):6-13. doi: 10.1016/j.hkjot.2017.10.004
- 35 5. Wong TM, Leung FKL, Lau TW, et al. Effectiveness of a Day Rehabilitation Program in
36 Improving Functional Outcome and Reducing Mortality and Readmission of Elderly
37 Patients With Fragility Hip Fractures. *Geriatr Orthop Surg Rehabil* 2018;9:1-8. doi:
38 10.1177/2151459318759355
- 39 6. El-Khoury F, Cassou B, Charles M-A, Dargent-Molina P. The effect of fall prevention
40 exercise programmes on fall induced injuries in community dwelling older adults:
41 systematic review and meta-analysis of randomised controlled trials. *BMJ* 2013;347

- 1 7. Lee JLC, Lou VWQ, Kwan RYC. The Experience of Participating in Remotely Delivered
2 Online Exercise Classes During the COVID-19 Pandemic Among Community-
3 Dwelling Older Adults and Its Postpandemic Implications. *J Aging Phys Act*
4 2023;1(aop):1-9.
- 5 8. Gamble C, Van Haastregt J, Van Dam Van Isselt E, et al. Effectiveness of guided
6 telerehabilitation on functional performance in community-dwelling older adults: A
7 systematic review. *Clin Rehabil* 2024;38(4):457-77. doi:
8 10.1177/02692155231217411
- 9 9. Costa RRG, Dorneles JR, Veloso JH, et al. Synchronous and asynchronous tele-exercise
10 during the coronavirus disease 2019 pandemic: Comparisons of implementation and
11 training load in individuals with spinal cord injury. *J Telemed Telecare*
12 2023;29(4):308-17. doi: 10.1177/1357633x20982732
- 13 10. Chan JKY, Klainin-Yobas P, Chi Y, et al. The effectiveness of e-interventions on fall,
14 neuromuscular functions and quality of life in community-dwelling older adults: A
15 systematic review and meta-analysis. *Int J Nurs Stud* 2021;113:103784. doi:
16 <https://doi.org/10.1016/j.ijnurstu.2020.103784>
- 17 11. Granet J, Peyrusqué E, Ruiz F, et al. Online physical exercise intervention in older adults
18 during lockdown: Can we improve the recipe? *Aging Clin Exp Res* 2023;35(3):551-
19 60. doi: 10.1007/s40520-022-02329-z
- 20 12. Zengin Alpozgen A, Kardes K, Acikbas E, et al. The effectiveness of synchronous tele-
21 exercise to maintain the physical fitness, quality of life, and mood of older people-a
22 randomized and controlled study. *Eur Geriatr Med* 2022;13(5):1177-85.
- 23 13. Wu G, Keyes L, Callas P, et al. Comparison of telecommunication, community, and home-
24 based Tai Chi exercise programs on compliance and effectiveness in elders at risk for
25 falls. *Arch Phys Med Rehabil* 2010;91(6):849-56.
- 26 14. Hong J, Kong H-J, Yoon H-J. Web-Based Telepresence Exercise Program for Community-
27 Dwelling Elderly Women With a High Risk of Falling: Randomized Controlled Trial.
28 *JMIR mHealth and uHealth* 2018;6(5):e132. doi: 10.2196/mhealth.9563
- 29 15. Bernocchi P, Giordano A, Pintavalle G, et al. Feasibility and clinical efficacy of a
30 multidisciplinary home-telehealth program to prevent falls in older adults: a
31 randomized controlled trial. *J Am Med Dir Assoc* 2019;20(3):340-46.
- 32 16. Yi D, Yim J. Remote home-based exercise program to improve the mental state, balance,
33 and physical function and prevent falls in adults aged 65 years and older during the
34 COVID-19 pandemic in Seoul, Korea. *Med Sci Monit* 2021;27:e935496-1.
- 35 17. Light K, Bishop M, Wright T. Telephone calls make a difference in home balance training
36 outcomes: a randomized trial. *J Geriatr Phys Ther* 2016;39(3):97-101.
- 37 18. Sterne JA, Savović J, Page MJ, et al. RoB 2: a revised tool for assessing risk of bias in
38 randomised trials. *BMJ* 2019;366
- 39 19. Chan KOW, Yuen PP, Fong BYF, et al. Effectiveness of telehealth in preventive care: a
40 study protocol for a randomised controlled trial of tele-exercise programme involving
41 older people with possible sarcopenia or at risk of fall. *BMC Geriatr* 2023;23(1):845.
42 doi: 10.1186/s12877-023-04535-4
- 43 20. Schulz KF, Altman DG, Moher D. CONSORT 2010 statement: updated guidelines for
44 reporting parallel group randomised trials. *J Pharmacol Pharmacother*
45 2010;1(2):100-07.
- 46 21. Benavent-Caballer V, Rosado-Calatayud P, Segura-Ortí E, et al. The effectiveness of a
47 video-supported group-based Otago exercise programme on physical performance in
48 community-dwelling older adults: a preliminary study. *Physiotherapy*
49 2016;102(3):280-86. doi: <https://doi.org/10.1016/j.physio.2015.08.002>
- 50 22. Nyman SR, Victor CR. Older people's participation in and engagement with falls
51 prevention interventions in community settings: an augment to the cochrane
52 systematic review. *Age Ageing* 2011;41(1):16-23. doi: 10.1093/ageing/afr103
- 53 23. Campbell A, Robertson MC. Otago exercise programme to prevent falls in older adults.
54 *Wellington: ACC Thinksafe* 2003

- 1 24. Tombaugh TN, McIntyre NJ. The mini-mental state examination: a comprehensive
2 review. *J Am Geriatr Soc* 1992;40(9):922-35.
- 3 25. Ngai S, Cheung R, Lam P, et al. Validation and reliability of the Physical Activity Scale
4 for the Elderly in Chinese population. *J Rehabil Med* 2012;44(5):462-65. doi:
5 10.2340/16501977-0953
- 6 26. Kwan MMS, Tsang WWN, Close JCT, Lord SR. Development and validation of a
7 Chinese version of the Falls Efficacy Scale International. *Arch Gerontol Geriatr*
8 2013;56(1):169-74. doi: <https://doi.org/10.1016/j.archger.2012.10.007>
- 9 27. Chen L-K, Woo J, Assantachai P, et al. Asian Working Group for Sarcopenia: 2019
10 consensus update on sarcopenia diagnosis and treatment. *J Am Med Dir Assoc*
11 2020;21(3):300-07. e2.
- 12 28. Chung P-K, Leung K-M. Psychometric properties of eight-item physical activity
13 enjoyment scale in a Chinese population. *J Aging Phys Act* 2019;27(1):61-66.
- 14 29. White IR, Horton NJ, Carpenter J, Pocock SJ. Strategy for intention to treat analysis in
15 randomised trials with missing outcome data. *BMJ* 2011;342
- 16 30. Garrard L, Boyle DK, Simon M, et al. Reliability and validity of the NDNQI® injury falls
17 measure. *West J Nurs Res* 2016;38(1):111-28.
- 18 31. Martins AC, Santos C, Silva C, et al. Does modified Otago Exercise Program improves
19 balance in older people? A systematic review. *Prev Med Rep* 2018;11:231-39. doi:
20 <https://doi.org/10.1016/j.pmedr.2018.06.015>
- 21 32. Wu S, Guo Y, Cao Z, et al. Effects of Otago exercise program on physical function in
22 older adults: A systematic review and meta-analysis of randomized controlled trials.
23 *Arch Gerontol Geriatr* 2024;124:105470. doi:
24 <https://doi.org/10.1016/j.archger.2024.105470>
- 25 33. Hager A-GM, Mathieu N, Lenoble-Hoskovec C, et al. Effects of three home-based
26 exercise programmes regarding falls, quality of life and exercise-adherence in older
27 adults at risk of falling: protocol for a randomized controlled trial. *BMC Geriatr*
28 2019;19(1):1-11.
- 29 34. Picorelli AMA, Pereira LSM, Pereira DS, et al. Adherence to exercise programs for older
30 people is influenced by program characteristics and personal factors: a systematic
31 review. *J Physiother* 2014;60(3):151-56. doi:
32 <https://doi.org/10.1016/j.jphys.2014.06.012>
- 33 35. Farrance C, Tsofliou F, Clark C. Adherence to community based group exercise
34 interventions for older people: A mixed-methods systematic review. *Prev Med*
35 2016;87:155-66.
- 36 36. Martel D, Lauzé M, Agnoux A, et al. Comparing the effects of a home-based exercise
37 program using a gerontechnology to a community-based group exercise program on
38 functional capacities in older adults after a minor injury. *Exp Gerontol* 2018;108:41-
39 47. doi: <https://doi.org/10.1016/j.exger.2018.03.016>
- 40 37. Brustio PR, Klonova A, Rudi D, et al. Effects of online and face-to-face exercise training
41 compared in healthy older adults: a feasibility study. *Sport Sci Health* 2024 doi:
42 10.1007/s11332-024-01166-z
- 43 38. Schwartz H, Har-Nir I, Wenhoda T, Halperin I. Staying physically active during the
44 COVID-19 quarantine: exploring the feasibility of live, online, group training
45 sessions among older adults. *Transl Behav Med* 2021;11(2):314-22. doi:
46 10.1093/tbm/ibaa141
- 47 39. Dawson R, Oliveira JS, Kwok WS, et al. Exercise Interventions Delivered Through
48 Telehealth to Improve Physical Functioning for Older Adults with Frailty, Cognitive,
49 or Mobility Disability: A Systematic Review and Meta-Analysis. *Telemedicine and e-
50 Health* 2023
- 51 40. Gell N, Hoffman E, Patel K. Technology Support Challenges and Recommendations for
52 Adapting an Evidence-Based Exercise Program for Remote Delivery to Older Adults:
53 Exploratory Mixed Methods Study. *JMIR Aging* 2021;4(4):e27645. doi:
54 10.2196/27645

- 1 41. García-Gollarte F, Mora-Concepción A, Pinazo-Hernandis S, et al. Effectiveness of a
2 Supervised Group-Based Otago Exercise Program on Functional Performance in Frail
3 Institutionalized Older Adults: A Multicenter Randomized Controlled Trial. *J Geriatr*
4 *Phys Ther* 2023;46(1)
- 5 42. Liew LK, Tan MP, Tan PJ, et al. The Modified Otago exercises prevent grip strength
6 deterioration among older fallers in the Malaysian falls assessment and intervention
7 trial (MyFAIT). *J Geriatr Phys Ther* 2019;42(3):123-29.
- 8 43. Fisher J, Steele J, McKinnon P, McKinnon S. Strength Gains as a Result of Brief,
9 Infrequent Resistance Exercise in Older Adults. *J Sports Med* 2014;2014:1-7. doi:
10 10.1155/2014/731890
- 11 44. Duckham RL, Masud T, Taylor R, et al. Randomised controlled trial of the effectiveness
12 of community group and home-based falls prevention exercise programmes on bone
13 health in older people: the ProAct65+ bone study. *Age Ageing* 2015;44(4):573-79.
14 doi: 10.1093/ageing/afv055
- 15 45. Hong J, Kim J, Kim SW, Kong H-J. Effects of home-based tele-exercise on sarcopenia
16 among community-dwelling elderly adults: Body composition and functional fitness.
17 *Exp Gerontol* 2017;87:33-39. doi: <https://doi.org/10.1016/j.exger.2016.11.002>
- 18 46. Robinson SM, Reginster JY, Rizzoli R, et al. Does nutrition play a role in the prevention
19 and management of sarcopenia? *Clin Nutr* 2018;37(4):1121-32. doi:
20 <https://doi.org/10.1016/j.clnu.2017.08.016>
- 21 47. Beauchamp MR, Hulteen RM, Ruissen GR, et al. Online-Delivered Group and Personal
22 Exercise Programs to Support Low Active Older Adults' Mental Health During the
23 COVID-19 Pandemic: Randomized Controlled Trial. *J Med Internet Res*
24 2021;23(7):e30709. doi: 10.2196/30709
- 25 48. Hinman RS, Campbell PK, Kimp AJ, et al. Telerehabilitation consultations with a
26 physiotherapist for chronic knee pain versus in-person consultations in Australia: the
27 PEAK non-inferiority randomised controlled trial. *The Lancet*
28 2024;403(10433):1267-78.
- 29 49. Census and Statistics Department HKSAR. 2021 Population Census - Main Results. In:
30 Census and Statistics Department, ed. Hong Kong, 2022.
- 31 50. World Health Organization. Physical activity 2020 [Available from:
32 <https://www.who.int/news-room/fact-sheets/detail/physical-activity> accessed 23 Nov
33 2022.

34