

## Successful strategies that address gender-related barriers and promote bodily autonomy within efforts to scale up and sustain post-pregnancy contraception: A scoping review

Arachu Castro<sup>1\*</sup>, Rita Kabra<sup>2</sup>, Anna Coates<sup>2</sup>, James Kiarie<sup>2</sup>

<sup>1</sup> Tulane University School of Public Health and Tropical Medicine

<sup>2</sup> World Health Organization

\* Corresponding author: Department of International Health and Sustainable Development, 1440 Canal Street, New Orleans, LA 70112, USA, [acastrol@tulane.edu](mailto:acastrol@tulane.edu).

### ABSTRACT

**Introduction:** Acknowledging the integral role of bodily autonomy in advancing gender equality, our study aimed to assess the extent to which strategies used in postpartum and post-abortion contraception have effectively equipped women, girls, and gender-diverse individuals with the tools, knowledge, and resources required to make autonomous decisions that align with their diverse life experiences.

**Methods:** We conducted a scoping review using the databases PubMed, EBSCOhost, EMBASE, and SciSpace. We included implementation, evaluation, and experimental studies published in any language between 2013 and 2023 and excluded studies not meeting these criteria. We used a WHO scale to determine the level of gender responsiveness.

**Results:** We found 30 implementation, evaluation, and experimental studies published in any language between 2013 and 2023. We categorized the strategies as gender-transformative (4 studies), gender-specific (24 studies), and gender-sensitive (2 studies). None of the studies reported strategies hindering reproductive health and rights. All strategies involved women and girls, and none explicitly targeted gender-diverse people capable of childbearing.

**Conclusion:** Gender-transformative strategies shared a commitment to empowering women with the autonomy to make informed decisions about post-pregnancy contraception through a) delivering personalized counseling that respected each woman's reproductive goals and ensured privacy during these discussions; b) integrating the cultural and familial context of women's health decisions; and c) promoting a rights-based approach that prioritized informed consent and defended women's reproductive rights. These interventions were delivered in contexts of quality improvement strategies integrating contraception services into the continuum of post-pregnancy care, offering women continuous access to information. Gender-specific strategies focused on informed contraceptive decision-making while recognizing the potential impact of gender dynamics on contraceptive use without actively challenging the underlying gender norms or power relations. Gender-sensitive strategies, although indicated gender awareness, did not address the process of informed contraceptive decision-making nor emphasize the provision of supportive environments that respect and enhance bodily autonomy.

### KEYWORDS

Pregnancy, Reproductive Health, Gender Equity, Reproductive Rights, Contraception.

## KEY MESSAGES

**What is already known on this topic:** Previous scoping and systematic reviews have explored strategies to increase post-pregnancy contraception uptake globally. However, none have specifically focused on strategies that promote bodily autonomy while addressing gender-related barriers. Our study addresses this gap by providing a comprehensive understanding of such strategies and their impact on scaling up and sustaining post-pregnancy contraception.

**What this study adds:** This study provides new insights by being the first scoping review to focus on strategies promoting bodily autonomy in addressing gender-related barriers to scaling up and sustaining post-pregnancy contraception. The gender-transformative strategies reported in the studies shared a commitment to empowering women with the autonomy to make informed decisions about post-pregnancy contraception through a) delivering personalized counseling that respected each woman's reproductive goals and ensured privacy during these discussions; b) integrating the cultural and familial context of women's health decisions; and c) promoting a rights-based approach that prioritized informed consent and defended women's reproductive rights.

**How this study might affect research, practice, or policy:** This study highlights the importance of integrating gender-transformative activities into post-pregnancy contraceptive strategies. It underscores the necessity of understanding and addressing local gender norms and the broader health system context to effectively promote bodily autonomy. The findings suggest that success should not be solely measured by contraceptive uptake but also by how well interventions address gender-related barriers. Future research should focus on developing and validating indicators that evaluate these barriers and promote bodily autonomy, ensuring comprehensive strategies that truly empower women, girls, and gender-diverse individuals with the means, abilities, and assets to make informed choices that resonate with the broader spectrum of their lives.

## INTRODUCTION

Within a framework of sexual and reproductive health (SRH) and rights, the World Health Organization (WHO) aims to fortify the gender-responsive approaches in the escalation and preservation of post-pregnancy contraceptive services that contribute to a fulfilling fertility desire (1). This endeavor stems from recognizing that reproductive and sexual freedom is foundational to gender equality and that “any individual has the necessary agency to determine their reproductive decisions and sexuality, regardless of gendered norms and gender-based discrimination” (2). Such freedom paves the way for women, girls, and those of diverse gender identities to gain comprehensive control over their bodies and life trajectories, irrespective of prevailing restrictive gender perceptions related to sexuality, reproduction, or any gender-related biases. Consequently, the availability of gender-responsive contraception, which goes beyond merely aiming to improve uptake or access to contraceptives but more ambitiously aims to promote personal bodily sovereignty, becomes indispensable in our quest for gender equality.

Accordingly, our study aims consist of evaluating if and how the structure and provision of postpartum and post-abortion contraceptive services empower women, girls, and gender-diverse individuals with the means, abilities, and assets to make informed choices that resonate with the broader spectrum of their lives and challenges harmful gender norms, which may restrict reproductive and life choices. We focus on discerning how such interventions can effectively and intentionally foster bodily autonomy as a primary goal. The evidence generated through this review will advance the knowledge of strategies that successfully address gender-related barriers to scaling up and sustaining postpartum and post-abortion contraception in ways that promote the exercise of bodily autonomy.

## METHODOLOGY

This scoping review follows the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) Statement (3), the PRISMA Extension for Scoping Reviews guidelines (4), and an adaptation of the Arksey and O'Malley framework for conducting scoping reviews (5). The scoping review and its protocol is registered in the OSF repository (6). The guidelines include several stages:

### 1. Identifying the research question:

We searched for evidence of addressing gender-related barriers and promoting bodily autonomy in successful strategies to scale up and sustain postpartum and post-abortion contraception. In the absence of other measures in most impact evaluations, we defined success according to the metrics defined for most interventions, that is, increased contraceptive uptake post-pregnancy, therefore, increasing access to sexual and reproductive health services.

### 2. Identifying relevant studies:

We searched PubMed, EBSCOhost (Academic Search Complete), and EMBASE online databases. Search terms “in any field” included: ((postpartum contraception) OR (postpartum family planning) OR (post abortion contraception) OR (post abortion family planning) OR (prenatal contraception) OR (prenatal family planning) OR (antenatal contraception) OR (antenatal family planning)) AND ((gender barrier) OR (gender issues) OR (cultural issues)). We also searched the database SciSpace, asking: “What are successful strategies for expanding postpartum or post-abortion contraception?” We included implementation, evaluation, and experimental studies published in any language between 2013 and 2023. We excluded papers that did not meet these criteria. One researcher (AC) conducted all the searches and the following steps in October 2023.

### 3. Study selection:

We refined the research strategy in a set of steps. First, we imported all the references to Endnote 20 and deleted duplicates. Second, we imported the remaining references to Covidence and conducted title and abstract screening. We removed publications that were either irrelevant, not found, abstract only, editorials, or letters. Third, we imported the full version of the remaining references plus additional references found through citation search to SciSpace and asked: “Does the paper identify successes to postpartum or post-abortion contraception?” If positive, we refined the search by asking two follow-up questions tested multiple times to capture as many eligible papers as possible: “Were any successes identified?” and “Was the intervention successful?” We excluded the papers that yielded a negative response. We selected the remaining papers for full review and classified them for full review according to the type of studies. We retained implementation, evaluation, and experimental studies and discarded reviews and observational studies. We assessed as eligible the publications that have successfully addressed barriers to the broad adoption and ongoing use of post-pregnancy contraceptive methods, emphasizing the reinforcement of bodily autonomy. One researcher (AC) collected and reviewed the data.

### 4. Charting the data:

For each publication, we extracted information on the country where the research was conducted, the population studied, and the methodological design. We charted the data according to four themes:

1. The level of gender responsiveness. To determine the level of each study, we used WHO's Gender Assessment Tool and Gender Responsive Assessment Scale (7) and additional considerations specific to sexual and reproductive health (2), shown in **Box 1**.

2. Characteristics of gender-responsive strategies that reported success in expanding post-pregnancy contraception, especially those for which impact evaluations have been conducted.
3. Key components of gender-transformative strategies that can be adapted to different contexts and/or scaled up.
4. Factors associated with using digital tools to promote post-pregnancy contraceptive use.

*Box 1: Levels of gender responsiveness according to the WHO’s Gender Responsive Assessment Scale and gender responsiveness of post-pregnancy contraception services*

Gender Responsive Assessment Scale				Gender responsiveness of post-pregnancy contraception programs
Level 1	Gender-unequal	Perpetuates gender inequality by reinforcing unbalanced norms, roles and relations and often leads to one sex enjoying more rights or opportunities than the other.	Hinder the achievement of gender equality and health equity.	Hinder the achievement of sexual and reproductive health and rights.
Level 2	Gender-blind	Ignores gender norms, roles and relations and very often reinforces gender-based discrimination. By ignoring differences in opportunities and resource allocation for women and men, such policies are often assumed to be “fair” as they claim to treat everyone the same.		
Level 3	Gender-sensitive	Indicates gender awareness, although no remedial action is developed.	Represents a starting point for gender programming.	Designed to increase access to adequate post-pregnancy contraception without addressing gender norms and relations. Results are only based on service delivery and specific health outcomes without considering autonomy or informed decision-making.
Level 4	Gender-specific	Considers women’s and men’s specific needs and intentionally targets and benefits a specific group of women or men to achieve certain policy or program goals or meet certain needs. Such policies often make it easier for women and men to fulfill duties that are ascribed to them based on their gender roles but do not address underlying causes of gender differences.	Represent characteristics of gender-responsive health policies and programs.	Designed to increase access to appropriate sexual and reproductive health resources through the provision of post-pregnancy contraception. They address gender norms and relations and/or promote informed decision-making.
Level 5	Gender-transformative	Addresses the causes of gender-based health inequities by including ways to transform harmful gender norms, roles, and relations. The objective of such programs is often to promote gender equality and foster progressive changes in power relationships between women and men.		Designed to promote empowering reproductive and sexual agency and bodily autonomy through post-pregnancy contraception by transforming gender norms and relations, including those related to the relationship between sexuality and reproduction, addressing gender-based discrimination, and promoting informed decision-making.

*Source:* Prepared by the authors based on WHO’s Gender Responsive Assessment Scale (7) and Coates A, Allotey P. Global health, sexual and reproductive health and rights, and gender: square pegs, round holes. *BMJ Glob Health.* 2023;8(1) (2).

We used the acronyms shown in **Box 2**.

*Box 2: List of acronyms*

ANC	Antenatal care
BCS	Balanced Counseling Strategy
CHW	Community health worker
FIGO	International Federation of Gynecology and Obstetrics
FP	Family planning
IPP	Immediate postpartum
IMCH	Integrated Maternal and Child Health
IUD	Intrauterine device
L&D	Labor and delivery
LAM	Lactation amenorrhea method
LARC	Long-acting reversible contraceptive
MCH	Maternal and child health
MOH	Ministry of Health
NGO	Nongovernmental organization
PAC	Postabortion care
PAFP	Postabortion family planning
PNC	Postnatal care
PPFP	Postpartum family planning
PPIUD	Postpartum intrauterine device
RMNCH	Reproductive, maternal, newborn, and child health
SRH	Sexual and reproductive health
WHO	World Health Organization


## 5. Patient and Public Involvement:

There was no patient or public involvement in this study.

## RESULTS

The PRISMA flow chart (**Figure 1**) shows the identification, screening, and inclusion process of the 30 studies retained. Twenty-seven studies were conducted in a single country and 3 in multiple countries. Combined, the studies were conducted in: Afghanistan (1), Bangladesh (1), Benin (1), Chad (2), China (1), Côte d'Ivoire (1), the Democratic Republic of the Congo (1), Djibouti (1), Ethiopia (3), Ghana (2), Kenya (2), Malawi (1), Mali (1), Nepal (3), Niger (1), Nigeria (2), Pakistan (1), Rwanda (2), Senegal (1), Somalia (1), South Africa (1), Spain (1), Sri Lanka (1), Tanzania (3), Togo (1), and United States (4). All the studies were published in English.

Figure 1: PRISMA flow diagram



Source: Prepared by the authors following the PRISMA 2020 statement (3).

### 1. Level of gender responsiveness

In **Table 1**, we present the 30 studies according to the level of gender responsiveness reported, starting with level 5, in chronological order of publication. Four studies reported gender-transformative strategies (level 5), 24 were gender-specific (level 4), and two were gender-sensitive (level 3). Therefore, none of the strategies hindered the achievement of reproductive health and rights. All the strategies included women and girls, and none were explicitly targeted at gender-diverse people capable of childbearing. Additional information for each study is available in **Annex 1**.

Table 1: Level of gender-responsiveness of studies that report successful strategies in expanding post-pregnancy contraception

Reference and year	Country	Target population	Description of Intervention or strategy
<b>Level 5 – Gender-transformative</b>			
Tawfik, Y., et al. (2014) (8)	Afghanistan	Postpartum women	Integrates FP into postpartum care by providing counseling and addressing each woman's reproductive intentions in private spaces. Involved husbands and mothers-in-law with the aim of actively challenge power dynamics and promoting women's reproductive autonomy.
Curry, D. W., et al. (2015) (9)	Chad, the Democratic Republic of the Congo, Djibouti, Mali, and Pakistan	Women of reproductive age and women receiving post-abortion care	Focuses on improving FP in crisis-affected regions. Engages men and women in discussions, promotes informed contraceptive decision-making by women, and addresses gender relations.
Samuel, M., et al. (2016) (10)	Ethiopia	Women receiving abortion care	Employs a rights-based approach to postabortion FP, emphasizing women's reproductive autonomy and informed contraceptive decision-making.
Mbehero, F., et al. (2021) (11)	Kenya	Women receiving abortion care	Addresses the provision of safe abortion, post-abortion care, and contraception, focusing on women's health services and youth-friendly services, promoting sexual and reproductive rights, and challenging gender norms.
<b>Level 4 – Gender-specific</b>			
Adanikin, A. I., et al. (2013) (12)	Nigeria	Pregnant women in the 3 <sup>rd</sup> trimester	Focuses on the specific needs of pregnant women, offering tailored contraceptive counseling and promoting informed contraceptive decision-making. It acknowledges gender-specific reproductive roles but does not provide information on how they addressed underlying gender norms or power relations
Cooper, C. M., et al. (2014) (13)	Bangladesh	Postpartum women	Focuses on the specific needs of postpartum women to promote contraceptive uptake and informed contraceptive decision-making. It acknowledges gender-specific reproductive roles and involves husbands and mothers-in-law but does not challenge underlying gender norms or power relations.
Huang, Y., R., et al. (2014) (14)	China	Women receiving childbirth care	Focuses on the specific needs of rural-to-urban migrant women regarding postpartum contraception. The intervention provides free contraceptive counseling and a choice of methods, which helps to overcome financial barriers and empowers single and married women to make informed decisions about their reproductive health but does not actively seek to transform gender relations or norms.
Pleah, T., et al. (2016) (15)	Benin, Chad, Côte d'Ivoire, Niger, Senegal, and Togo	Women during pregnancy, labor, and immediate postpartum	Targets specific needs of postpartum women for IUD services, addresses cultural barriers, and promotes informed contraceptive decision-making, but does not substantially challenge underlying gender norms or roles.
Chukwumalu, K., et al. (2017) (16)	Somalia	Women receiving post-abortion care	Focuses on service provision and informed contraceptive decision-making without actively challenging prevailing gender norms.
Gbagbo, F. Y., et al. (2018) (17)	Ghana	Women receiving post-abortion care	Focuses on increasing post-abortion contraception through a comprehensive model that relies on informed contraceptive decision-making but does not explicitly address gender relations or aim to transform them. While the study acknowledges the presence of partner pressure, it does not explicitly analyze how these factors might be impacting the program's effectiveness or propose specific strategies to address them.
Ingabire, R., et al. (2018) (18)	Rwanda	Women during pregnancy, labor, immediate postpartum, infant vaccination visits, or in the community	Implements a multi-level intervention for PPIUD services involving male partners, but primarily focuses on increasing women's contraceptive uptake without fundamentally changing gender dynamics.
Lori, J. R., et al. (2018) (19)	Ghana	Pregnant women	Increases FP use through group ANC, promoting women's informed contraceptive decision-making, without directly challenging gender norms.
Torres, L. N., et al. (2018) (20)	United States	Postpartum women immediately after giving birth to a premature infant	Enhances knowledge and use of postpartum LARC methods, focusing on women's informed contraceptive decision-making, but without addressing underlying causes of gender roles or relations.
Wendot, S., et al. (2018) (21)	Kenya	Women receiving safe abortion and post-abortion care	Improves quality of post-abortion FP counseling and method uptake by using BCS (which promotes informed decision-making), targeting women's informed contraceptive decision-making without actively addressing gender power dynamics. While the study acknowledges the potential impact of gender dynamics (by asking for relationship status and partner support for contraceptive use), it does not explicitly analyze how these factors might be impacting women's decisions regarding PAFP.
Karra, M., et al. (2019) (22)	Sri Lanka	Women receiving childbirth care	Increases counseling rates and informed contraceptive decision-making but focuses on women's health services without challenging underlying gender norms or how these norms might be impacting women's experiences with the program or their choices regarding PPIUD.
Mossie, M. Y., et al. (2019) (23)	Ethiopia	Postpartum women	Focuses on tracking PFPF counseling and informed contraceptive decision-making through the continuum of care. The study acknowledges couple or family discussion of PFPF as an input to uptake but does not analyze how gender dynamics or social norms might be impacting women's experiences with PFPF or their decision-making processes.

Pradhan, E., et al. (2019) (24)	Nepal	Women in the immediate postpartum	Increases counseling and uptake of PPIUD, increasing women's contraceptive knowledge without explicitly analyzing how gender dynamics or social norms might be impacting women's experiences with the program or their choices regarding PPIUD.
Stephens, B., et al. (2019) (25)	Tanzania	Women receiving post-abortion care	Focuses on expanding post-abortion contraceptive coverage with BCS (which promotes informed contraceptive decision-making) and respectful care but does not actively seek to transform gender relations.
Haider, S., et al. (2020) (26)	United States	Postpartum women within 4.5 months of giving birth	Offers co-located contraceptive services with infant care to enhance informed contraceptive decision-making. Considers marital status or living arrangements, acknowledging the potential impact of gender dynamics, but does not develop actions to change gender norms or relations.
Huber-Krum, S., et al. (2020) (27)	Nepal	Women during pregnancy, early labor, and immediate postpartum	Focuses on training providers in counseling and PPIUD insertion, promoting informed contraceptive decision-making. Acknowledges gender norms and social pressure to have children but does not analyze how these might be impacting women's experiences with the program or their decision-making processes.
Lacy, M. M., et al. (2020) (28)	United States	Women seeking immediate postpartum or post-abortion LARC	Increases access to IPP LARC, focusing on women's health needs and informed contraceptive decision-making but does not analyze how gender dynamics or social norms might be impacting women's experiences with IPP LARC or their decision-making processes.
Pearson, E., et al. (2020) (29)	Tanzania	Women during pregnancy and after admission for delivery	Increases counseling and choice of PPIUD, targeting women's informed contraceptive decision-making. Considers marital status or living arrangements, acknowledging the potential impact of gender dynamics, but does not develop actions to change gender norms or relations.
Reyes-Lacalle, A., L., et al. (2020) (30)	Spain	Pregnant women	Enhances contraceptive use through additional counseling, focusing on women's informed contraceptive decision-making. Considers the partner's involvement and the marital or civil union status and acknowledges the study limitation that intimate partner violence was not analyzed. However, it does not analyze how gender dynamics or social norms might be impacting women's experiences with contraception or their decision-making processes.
Sitrin, D., et al. (2020) (31)	Ethiopia	Pregnant women	Integrates PPF into community-level services, focusing on women's contraceptive informed contraceptive decision-making and use. Considers marital status but does not analyze how gender dynamics or social norms might be impacting women's experiences with the program or their decision-making processes.
Wu, W. J., et al. (2020) (32)	Nepal	Women in the 8 <sup>th</sup> month of pregnancy and at 1, 5, and 10 months postpartum	Increases contraceptive use through BCS (which promotes informed contraceptive decision-making), community-based counseling, targeting women's needs without explicitly addressing gender dynamics.
Espey, J., et al. (2021) (33)	Rwanda	Women during pregnancy, early labor & delivery, and infant vaccination visits	Implements a multilevel intervention to increase uptake of postpartum LARC methods, promoting informed contraceptive decision-making and male engagement but without seeking to transform gender norms and relations.
Karra, M., et al. (2022) (34)	Malawi	Women during pregnancy or within 6 months postpartum	Increases contraceptive use and birth spacing, targeting women informed contraceptive decision-making. As acknowledged in the study limitations, while the study provided guidance to women on ways to involve their husband or male partner via counseling and dialogue, leaving up to the women how much she wanted to involve him in any part of the counseling or intervention, it did not specifically address the social norms concerning community or partner perspectives on FP, alleging that it would have required a larger, cluster randomized, multiarmed trial.
Ijarotimi, O., et al. (2023) (35)	Nigeria	Postpartum women attending infant vaccination clinics.	Integrates FP education with infant vaccination, targeting women's informed contraceptive decision-making. Considers the marital status without addressing the underlying gender dynamics that might play a role in contraceptive choices.
<b>Level 3 – Gender-sensitive</b>			
Melnick, A. L., et al. (2016) (36)	United States	Vulnerable, low-income, first-time mothers at risk for unintended and short-interval pregnancies	Focuses on decreasing women's gaps in use of effective methods after childbirth to reduce the incidence of unintended and short-interval pregnancies but does not address informed contraceptive decision-making nor how gender dynamics might be impacting women's experiences with the program.
Furey, A., et al. (2019) (37)	Tanzania and South Africa	Women admitted for childbirth	Focuses on improving the uptake and choice of postpartum FP but does not address informed contraceptive decision-making nor how gender dynamics might be impacting women's experiences with the program.

## 2. Characteristics of gender-responsive strategies that reported success in expanding post-pregnancy contraception

The studies that reported gender-transformative strategies successful in expanding post-pregnancy contraception (8-11) shared the goal of improving the quality of contraception counseling and service provision. These interventions were delivered in contexts of quality improvement strategies that integrated contraception services into the continuum of post-pregnancy care and that offered women

continuous access to information. In Afghanistan (8), the intervention involved integrating FP into postpartum care using quality improvement approaches focused on improving the quality of counseling and turning the process user-centered. This included tailoring the counseling process to address each woman's specific reproductive intentions and contraceptive needs, finding private spaces for PFP counseling, and involving husbands and mothers-in-law in counseling. Follow-up calls from FP counselors clarified contraceptive methods and underscored essential FP messages. These enhanced approaches successfully integrated FP into postpartum care, resulting in more women receiving counseling on postpartum FP and acquiring their chosen method before leaving the hospital. The strategy addressed gender-based factors by involving husbands and mothers-in-law in FP counseling. Before the strategy, the implementation team had identified that husbands were not allowed in the maternity ward and that some women wanted to consult with their husbands or mothers-in-law before making their decision on contraception use. Creating private spaces for postpartum FP counseling where the husband or the mother-in-law could participate, in person or via mobile phone, was a gender-transformative approach that contributed to meeting women's needs and a rise in joint decision-making on SRH needs.

In Chad, the Democratic Republic of the Congo, Djibouti, Mali, and Pakistan (9), the choice of contraceptives among women of reproductive age and those receiving post-abortion care varied by country, with LARCs being more prevalent in specific regions and other methods like oral contraceptives and injectables being dominant in others. The study discussed the role of community norms related to gender and fertility in implementing FP services. It emphasized the importance of engaging men and women in discussions about rights, fertility, and contraception at the local level. The strategy also highlighted the role of religious leaders in promoting modern contraception, indicating an understanding of the gender dynamics within religious contexts. Improving FP accessibility and education in crisis-affected regions involved several components: a) facility assessments and stocking: conducting health facility assessments to identify infrastructure gaps, reviewing the presence and condition of basic needs, and reequipping facilities with essential supplies and equipment; b) competency-based training: providing comprehensive training to health care providers, using materials from Jhpiego and the Population Council; this includes both theoretical sessions and practical training; c) supervision and quality improvement: engaging with facility teams and community leaders to ensure a clean, efficient, and respectful environment and regular supervisory visits to assess clinic conditions, stock levels, and data consistency; d) supply chain management: ensuring a steady supply of contraceptive methods and related items by both procuring necessary items and working to strengthen supply chain management in collaboration with government systems; and e) community mobilization: raising awareness at the community level through various means such as radio, participatory theater, and group dialogues, and engaging religious leaders across faiths to raise awareness and ensure women's rights to access health services. The key to community mobilization's success was motivating and involving the entire community to recognize the advantages and possibilities in planning and spacing childbirth and facilitating access for women to utilize services.

In Ethiopia (10), the strategy consisted of a package of interventions for women seeking abortion care that promoted a rights-based approach to informed consent and decision-making. The strategy improved the quality of PAFP counseling, increased the contraceptive method mix, emphasized and prioritized respect for women's reproductive autonomy in the provision of PAFP services, promoted youth-friendly services, ensured providers' commitment to the sexual and reproductive rights of young women, promoted privacy and confidentiality during service delivery, and strengthened community outreach through health extension workers. These strategies resulted in an increase in postabortion choice of a contraceptive method and a greater proportion choosing the more effective, long-acting methods.

In Kenya (11), health providers were trained on abortion values clarification, attitude transformation, methods of uterine evacuation, management of abortion complications, and post-abortion contraception. Training and mentorship of providers on PAFP and youth-friendly services, along with community engagement and referrals, helped remove common barriers to care: a) the project addressed the issue of providers' hesitation to counsel and administer LARCs to adolescent or youth users through training and mentoring; b) each facility had separate procedure rooms for conducting medical vacuum



aspiration to ensure adequate space and user confidentiality; c) implemented training and mentorship programs for mid-level providers on safe abortion, post-abortion care, and post-abortion contraception, with a focus on youth-friendly services; d) implemented a quality assessment tool based on the WHO's seven pillars of health systems strengthening and the structures and processes in place to support the domains of quality (safety, timeliness, effectiveness, efficiency, equity, and people-centeredness); e) created health facility registers to track post-abortion contraceptive uptake; f) monitored mentees' clinical competencies through logbooks, tracking their proficiency in LARC insertion and removal and manual vacuum aspiration; g) monitored the increase in couple counseling for SRH, indicating a rise in joint decision-making on SRH needs; h) community engagement and referral activities to raise awareness and reduce the stigma associated with abortion and contraception; i) incorporation of strong SRH rights advocacy partners into the project to build networks and enhance access to contraception and safe abortion; j) prioritization of youth-led SRH rights advocacy groups in projects to empower young people to make informed decisions about their sexual and reproductive health; k) establishment of quality improvement teams at each facility to assess progress and address gaps, fostering a culture of quality and motivation; l) education of teachers on SRH rights, recognizing the impact of teachers on students' contraceptive use; and m) implemented strategies to increase male involvement in SRH rights, including holding forums for men and mentoring male champions. The increase in couple counseling for SRH indicated a rising normalization of joint decision-making on SRH needs.

The gender-specific strategies (12-35) focused on informed contraceptive decision-making while recognizing the potential impact of gender dynamics on contraceptive use without actively challenging the underlying gender norms or power relations. The most common gender-specific strategies to expand post-pregnancy contraception included: a) enhancing provider-patient interactions through tailored, structured, or balanced counseling, such as in Nigeria (12), the United States (20, 28), Kenya (21), Tanzania (25); b) increasing the variety of contraceptive choices to better respond to women's needs, such as in China (14) and Somalia (16); c) integrating contraceptive education with health services like antenatal visits, such as in Ethiopia where they used a modified Integrated Maternal and Child Health to record the woman's method choice during those visits (23), Nepal (24, 27, 32), Spain (30), Ethiopia (31), Rwanda (33), and Malawi (34); right after admission for delivery services, such as in Tanzania (29); during infant vaccinations, such as in Rwanda (18), Ghana (19), and Nigeria (35); during the first year postpartum, such as in China (14); and other maternal and child health services, such as in the United States (20, 26); d) introducing competency-based training for healthcare providers, such as in Benin, Chad, Côte d'Ivoire, Niger, Senegal, and Togo (15), Somalia (16), Ghana (17); e) employing storytelling with relatable content, such as in Bangladesh (13) and Ghana (19) or showing informational videos in hospital waiting areas, such as in Sri Lanka (22) and Nepal (24). Community engagement, particularly involving religious and community leaders, was critical in addressing cultural and gender norms, as shown in Bangladesh (13), Benin, Chad, Côte d'Ivoire, Niger, Senegal, Togo (15), and Somalia (16).

Finally, the gender-sensitive strategies reported in one study conducted in the United States (36) and one in Tanzania and South Africa (37) were aimed at either increasing the uptake and choice of postpartum contraception through education and training of midwives, nurses, and non-specialist doctors on counseling and provision or by enhancing an existing nurse home visiting program. These studies did not address informed contraceptive decision-making nor how gender dynamics might be impacting women's experiences with the program.

Among the studies reviewed, we did not find indicators that measure and evaluate gender-related barriers and promote bodily autonomy. **Annex 1** shows a list of indicators based on the strategies that measure and evaluate the rights-based counseling and provision of post-pregnancy contraception.

### 3. Key components of gender-transformative strategies that can be adapted to different contexts or scaled up

Based on the four gender-transformative studies (8-11), **Table 2** shows the multifaceted approach needed to successfully implement gender-transformative interventions in post-pregnancy contraception across different cultural and regional contexts. Details from all the studies are available in **Annex 2**.

*Table 2: Key components of gender-transformative strategies that can be adapted to different contexts or scaled up*

A. Training, Capacity Building, Continuity of Care, and Supply Chain Management
1. Training healthcare staff in user-centered and rights-based contraception counseling during pregnancy and post-pregnancy to ensure informed decision-making and improve service delivery.
2. Training and mentorship programs for healthcare providers on safe abortion, post-abortion care, and contraception, with a focus on youth-friendly services, to sustain quality improvement strategies that support gender-transformative strategies.
3. Monitoring and tracking the clinical competencies of providers providing gender-transformative care, e.g. through logbooks.
4. Creating private spaces for counseling to ensure privacy and comfort.
5. Ensuring follow-up to address each woman's specific reproductive intentions and contraceptive needs, e.g. through follow up calls to answer questions about contraceptive methods.
6. Implementing supply chain management approaches that consider local challenges to sustain quality improvement plans that support gender-transformative strategies.
7. Monitoring of health systems quality issues, including procurement and supply chains, to ensure women have access to a wide range of contraceptive methods to increase the proportion of post-pregnancy women leaving the facility with their method of choice.
B. Counseling and Empowerment
1. Tailoring the counseling process to address each woman's specific reproductive intentions and contraceptive needs and prioritize autonomous decision-making.
2. Involving key family members, such as partners and mothers-in-law, in counseling sessions to challenge existing power dynamics and promote women's reproductive autonomy and equality in reproductive health decision-making.
3. Increasing couple's counseling for SRH to normalize joint decision-making on SRH needs.
C. Community Engagement, Mobilization, and Advocacy
1. Engaging men and women in local discussions about rights, fertility, and contraception to address and transform cultural and gender-based norms and relations.
2. Community mobilization, engagement, and referral activities through radio, participatory theater, and group dialogues, involving religious leaders across faiths and local leaders to discuss the role of community norms related to gender and fertility and raise awareness and support women's rights to access health services.
3. Increasing male involvement in discussions about sexual and reproductive health and rights, fertility, and contraception, including holding forums for men and mentoring male champions, to address and transform cultural and gender-based norms and relations at the local level.
4. Prioritization of youth-led SRH rights advocacy groups in projects to address and transform cultural and gender-based norms and relations.
5. Incorporation of strong SRH rights advocacy partners in the interventions to build networks and enhance access to contraception and safe abortion.
6. Engaging in dialogue with local community groups, especially women's organizations to better understand local gender norms and women's reproductive needs.

*Note:* Prepared by the authors based on gender-transformative studies (8-11).

### 4. Factors associated with using digital tools to promote post-pregnancy contraceptive use

Of the 30 studies included in the review, ten referred to the use of digital tools to promote post-pregnancy contraceptive use (15, 18, 23, 25-30, 32). Two studies conducted in Nepal and one in Spain reported that digital technology in healthcare improved service integration and facilitated the dissemination of contraceptive information (27, 30, 32). In Tanzania, mobile applications and digital counseling tools were instrumental in training healthcare workers, giving them access to the latest information and techniques to meet users' needs effectively (25). Digital data collection and management systems enhanced the continuity and follow-up of women users of contraception. For example, in Benin, Chad, Côte d'Ivoire, Niger, Senegal, and Togo, digital tools helped identify PPIUD complications (15). In Ethiopia, digital health records helped improve service integration and continuity of care, track PFP, and facilitate decision-making and timely adoption of contraceptives (23). In Spain, women received reminders to consult contraception information during pregnancy (30). In Nepal, community health workers used mobile applications to individualize the counseling based on women's responses to the balanced counseling strategy questions and to follow up at subsequent visits on initiation, barriers to access, and continuation (32). Finally, tablets and software for data input were used to collect data in Rwanda (18), Tanzania (29), the United States (26, 28), Spain (30), and Nepal (32).

## DISCUSSION

We reviewed 30 studies that met all the inclusion criteria. All but two studies emphasized the need to provide a supportive environment that respects and enhances women's bodily autonomy. The gender-transformative strategies reported in four of the studies shared a commitment to empowering women with the autonomy to make informed decisions about post-pregnancy contraception. They achieved this through quality improvement strategies that seamlessly integrated contraception services into the continuum of postpartum and post-abortion care. Providers delivered personalized counseling that respected each woman's reproductive goals and ensured privacy during these discussions. The initiatives included follow-up support to offer women continuous access to information and reinforce contraceptive practices. These efforts promoted a rights-based approach, prioritizing informed consent and defending women's reproductive rights. Women often chose long-acting reversible contraceptives, reflecting a preference for more reliable and enduring contraception methods when they received comprehensive information. These programs acknowledged and integrated the cultural and family context of women's health decisions by involving husbands and mothers-in-law, not as vehicles to increase post-pregnancy contraception but to challenge existing power dynamics and promote women's reproductive autonomy and equality in reproductive health decision-making. The implementation of these strategies, carefully adapted to each context, underscores the effectiveness of a comprehensive, multi-layered approach in transforming health interventions related to gender dynamics. By tackling both service delivery and broader social elements like gender norms and community standards, the findings from the gender-transformative strategies illustrate the potential for significant advancements in expanding post-pregnancy contraception.

The gender-specific studies focused on informed contraceptive decision-making. The studies demonstrated that multiple counseling sessions during antenatal care, tailored counseling during the third trimester, the free provision of contraception integrated into childbirth and abortion care, a range of options in contraceptive methods, and follow-up visits were crucial in ensuring the informed contraceptive decision-making process. Training healthcare providers in insertion techniques and providing informed consent and rights-based counseling was pivotal in increasing the informed uptake of post-pregnancy LARC methods and, specifically, PPIUD. Community engagement strategies that included the participation of religious and community leaders were instrumental in facilitating women's access to health facilities for post-abortion care services. The strategies also acknowledged the need for comprehensive monitoring and evaluation frameworks for data-driven programmatic decisions, engagement of key stakeholders from the initial stages, and the importance of youth-friendly services. While they recognized the potential impact of gender dynamics on contraceptive use and some interventions involved key family members in the counseling process, some strategies did not actively challenge the underlying gender norms or power relations.

Although several scoping and systematic reviews have addressed strategies that increase post-pregnancy uptake of contraception around the world (38-46), this is the first scoping review of interventions that promote bodily autonomy when addressing gender-related barriers to scaling up and sustaining post-pregnancy contraception.

The review has allowed us to identify key elements of gender-transformative post-pregnancy contraceptive strategies, with some important components that can be adapted to diverse contexts. Personal counseling emerged as a common thread across many interventions within a framework of a user-centered approach. Similarly, community dialogue and involvement of advocacy organizations, especially women's organizations, are crucial to understanding local gender norms and women's reproductive needs. Both are fundamental to delivering the individualized care required to promote bodily autonomy. However, for bodily autonomy to be meaningfully realized, other elements are also vital. With some notable exceptions, few strategies noted the wider health system context, such as procurement and supply chains to enable providers to meet women's contraceptive needs, institutionalized quality assurance mechanisms stressing gender transformative approaches, and ensuring an enabling health policy and legal environment.

Understanding cultural and familial contexts is another key element. The precise intent of strategies tailored to context is significant to ensure that they do not inadvertently reinforce unequal power relations, rather than advancing women's bodily autonomy. It is, therefore, especially significant that although the review allowed us to generate a list of results indicators to measure and evaluate the rights-based counseling and provision of post-pregnancy contraception, we did not find indicators that measure and evaluate gender-related barriers and promote bodily autonomy. Whilst success is still only defined according to contraceptive access and uptake, it cannot be explicitly guaranteed that any intervention will be delivered in such a manner that it will meaningfully promote bodily autonomy,

We recommend that future research urgently focus on developing and validating indicators that specifically address gender-related barriers and promote bodily autonomy so that contraceptive uptake is not mistakenly understood as a sole proxy for bodily autonomy, but rather as a dual goal. There is a need for more comprehensive strategies that integrate these indicators—and thus approach and goals—into practice, ensuring that post-pregnancy contraception programs empower women and girls, with the tools, knowledge, and resources to make informed decisions that align with their life goals.

## FUNDING SOURCE

This work received funding from the UNDP-UNFPA-UNICEF-WHO-World Bank Special Programme of Research, Development and Research Training in Human Reproduction (HRP), a cosponsored program executed by the World Health Organization (WHO) (WHO-SRH/HRP-CFC grant number 2022/1240770-0). WHO hired Arachu Castro as a consultant to conduct the scoping review. None of the authors had competing interests.

## REFERENCES

1. World Health Organization. WHO recommendations on maternal and newborn care for a positive postnatal experience. Licence: CC BY-NC-SA 3.0 IGO. Geneva: World Health Organization; 2022.
2. Coates A, Allotey P. Global health, sexual and reproductive health and rights, and gender: square pegs, round holes. *BMJ Glob Health*. 2023;8(1).
3. Page MJ, McKenzie JE, Bossuyt PM, Boutron I, Hoffmann TC, Mulrow CD, et al. The PRISMA 2020 statement: An updated guideline for reporting systematic reviews. *J Clin Epidemiol*. 2021;134:178-89.
4. Tricco AC, Lillie E, Zarin W, O'Brien KK, Colquhoun H, Levac D, et al. PRISMA Extension for Scoping Reviews (PRISMA-ScR): Checklist and Explanation. *Ann Intern Med*. 2018;169(7):467-73.
5. Levac D, Colquhoun H, O'Brien KK. Scoping studies: advancing the methodology. *Implement Sci*. 2010;5:69.
6. Castro A, Kabra R, Coates A, Kiarie J. Gender-related barriers to postpartum and post-abortion contraception: A scoping review. [Scoping review protocol registration]. 2023.
7. World Health Organization. Gender mainstreaming for health managers: a practical approach. Geneva: World Health Organization; 2011.
8. Tawfik Y, Rahimzai M, Ahmadzai M, Clark PA, Kamgang E. Integrating family planning into postpartum care through modern quality improvement: experience from Afghanistan. *Global Health: Science and Practice*. 2014;2(2):226-33.
9. Curry DW, Rattan J, Nzau JJ, Giri K. Delivering high-quality family planning services in crisis-affected settings I: program implementation. *Glob Health Sci Pract*. 2015;3(1):14-24.
10. Samuel M, Fetters T, Desta D. Strengthening Postabortion Family Planning Services in Ethiopia: Expanding Contraceptive Choice and Improving Access to Long-Acting Reversible Contraception. *Global Health: Science and Practice*. 2016;4(S2):S60-S72.

11. Mbehero F, Momanyi R, Hesel K. Facilitating Uptake of Post-abortion Contraception for Young People in Kenya. *Front Glob Womens Health*. 2021;2:733957.
12. Adanikin AI, Onwudiegwu U, Loto OM. Influence of multiple antenatal counselling sessions on modern contraceptive uptake in Nigeria. *Eur J Contracept Reprod Health Care*. 2013;18(5):381-7.
13. Cooper CM, Ahmed S, Winch PJ, Pfitzer A, McKaig C, Baqui AH. Findings from the use of a narrative story and leaflet to influence shifts along the behavior change continuum toward postpartum contraceptive uptake in Sylhet District, Bangladesh. *Patient Educ Couns*. 2014;97(3):376-82.
14. Huang Y, Merkatz R, Zhu H, Roberts K, Sitruk-Ware R, Cheng L, et al. The free perinatal/postpartum contraceptive services project for migrant women in Shanghai: effects on the incidence of unintended pregnancy. *Contraception*. 2014;89(6):521-7.
15. Pleah T, Hyjazi Y, Austin S, Diallo A, Dao B, Waxman R, et al. Increasing Use of Postpartum Family Planning and the Postpartum IUD: Early Experiences in West and Central Africa. *Glob Health Sci Pract*. 2016;4 Suppl 2(Suppl 2):S140-52.
16. Chukwumalu K, Gallagher MC, Baunach S, Cannon A. Uptake of postabortion care services and acceptance of postabortion contraception in Puntland, Somalia. *Reprod Health Matters*. 2017;25(51):48-57.
17. Gbagbo FY, Elzohry AAM. Post Abortion Contraception Model: A Comprehensive Package for Improving Safe Abortion Care in Developing Countries. *Journal of Family Medicine*. 2018;1(1):12-21.
18. Ingabire R, Nyombayire J, Hoagland A, Da Costa V, Mazzei A, Haddad L, et al. Evaluation of a multi-level intervention to improve postpartum intrauterine device services in Rwanda. *Gates Open Res*. 2018;2:38.
19. Lori JR, Chuey M, Munro-Kramer ML, Ofosu-Darkwah H, Adanu RMK. Increasing postpartum family planning uptake through group antenatal care: a longitudinal prospective cohort design. *Reprod Health*. 2018;15(1):208.
20. Torres LN, Turok DK, Clark EAS, Sanders JN, Godfrey EM. Increasing IUD and Implant Use Among Those at Risk of a Subsequent Preterm Birth: A Randomized Controlled Trial of Postpartum Contraceptive Counseling. *Womens Health Issues*. 2018;28(5):393-400.
21. Wendot S, Scott RH, Nafula I, Theuri I, Ikiugu E, Footman K. Evaluating the impact of a quality management intervention on post-abortion contraceptive uptake in private sector clinics in western Kenya: a pre- and post-intervention study. *Reprod Health*. 2018;15(1):10.
22. Karra M, Pearson E, Pradhan E, de Silva R, Samarasekera A, Canning D, et al. The effect of a postpartum IUD intervention on counseling and choice: Evidence from a cluster-randomized stepped-wedge trial in Sri Lanka. *Trials*. 2019;20(1):407.
23. Mossie MY, Pfitzer A, Yusuf Y, Wondimu C, Bazant E, Bansal V, et al. Counseling at all contacts for postpartum contraceptive use: can paper-based tools help community health workers improve continuity of care? A qualitative study from Ethiopia. *Gates Open Res*. 2019;3:1652.
24. Pradhan E, Canning D, Shah IH, Puri M, Pearson E, Thapa K, et al. Integrating postpartum contraceptive counseling and IUD insertion services into maternity care in Nepal: results from stepped-wedge randomized controlled trial. *Reprod Health*. 2019;16(1):69.
25. Stephens B, Mwandalima IJ, Samma A, Lyatuu J, Mimno K, Komwihangiro J. Reducing Barriers to Postabortion Contraception: The Role of Expanding Coverage of Postabortion Care in Dar es Salaam, Tanzania. *Glob Health Sci Pract*. 2019;7(Suppl 2):S258-s70.
26. Haider S, Stoffel C, Rankin K, Uesugi K, Handler A, Caskey R. A Novel Approach to Postpartum Contraception Provision Combined with Infant Care: A Randomized, Controlled Trial. *Womens Health Issues*. 2020;30(2):83-92.
27. Huber-Krum S, Khadka A, Pradhan E, Rohr J, Puri M, Maharjan D, et al. The effect of antenatal counseling and intrauterine device insertion services on postpartum contraceptive use in Nepal: Results from a stepped-wedge randomized controlled trial. *Contraception*. 2020;101(6):384-92.

28. Lacy MM, McMurtry Baird S, Scott TA, Barker B, Zite NB. Statewide quality improvement initiative to implement immediate postpartum long-acting reversible contraception. *Am J Obstet Gynecol*. 2020;222(4S):S910 e1-S e8.
29. Pearson E, Senderowicz L, Pradhan E, Francis J, Muganyizi P, Shah I, et al. Effect of a postpartum family planning intervention on postpartum intrauterine device counseling and choice: evidence from a cluster-randomized trial in Tanzania. *BMC Womens Health*. 2020;20(1):102.
30. Reyes-Lacalle A, Montero-Pons L, Manresa-Dominguez JM, Cabedo-Ferreiro R, Seguranyes G, Falguera-Puig G. Perinatal contraceptive counselling: Effectiveness of a reinforcement intervention on top of standard clinical practice. *Midwifery*. 2020;83:102631.
31. Sitrin D, Jima GH, Pfitzer A, Wondimu C, Belete TW, Pleah T, et al. Effect of integrating postpartum family planning into the health extension program in Ethiopia on postpartum adoption of modern contraception. *Journal of Global Health Reports*. 2020;4.
32. Wu WJ, Tiwari A, Choudhury N, Basnett I, Bhatt R, Citrin D, et al. Community-based postpartum contraceptive counselling in rural Nepal: a mixed-methods evaluation. *Sex Reprod Health Matters*. 2020;28(2):1765646.
33. Espey J, Ingabire R, Nyombayire J, Hoagland A, Da Costa V, Mazzei A, et al. Postpartum long-acting contraception uptake and service delivery outcomes after a multilevel intervention in Kigali, Rwanda. *BMJ Sex Reprod Health*. 2021;47(3):173-8.
34. Karra M, Maggio D, Guo M, Ngwira B, Canning D. The causal effect of a family planning intervention on women's contraceptive use and birth spacing. *Proc Natl Acad Sci U S A*. 2022;119(22):e2200279119.
35. Ijarotimi O, Ijarotimi I, Ubom A, Sowemimo O, Orji E. The effect of integration of family planning education with infant vaccination visits on the uptake of postpartum family planning in Ile-Ife, Nigeria. *J Obstet Gynaecol*. 2023;43(1):2186774.
36. Melnick AL, Rdesinski RE, Marino M, Jacob-Files E, Gipson T, Kuyl M, et al. Randomized Controlled Trial of Home-Based Hormonal Contraceptive Dispensing for Women At Risk of Unintended Pregnancy. *Perspect Sex Reprod Health*. 2016;48(2):93-9.
37. Furey A, Fiander A. Postpartum contraception - the experience and findings from the Leading Safe Choices programme in Tanzania and South Africa. *BJOG*. 2019;126(11):1301-4.
38. Shah IH, Santhya KG, Cleland J. Postpartum and Post-Abortion Contraception: From Research to Programs. *Stud Fam Plann*. 2015;46(4):343-53.
39. Blazer C, Prata N. Postpartum family planning: current evidence on successful interventions. *Open Access J Contracept*. 2016;7:53-67.
40. Rogers C, Dantas JAR. Access to contraception and sexual and reproductive health information post-abortion: a systematic review of literature from low- and middle-income countries. *J Fam Plann Reprod Health Care*. 2017;43(4):309-18.
41. Wakuma B, Mosisa G, Etafa W, Mulisa D, Tolossa T, Fetensa G, et al. Postpartum modern contraception utilization and its determinants in Ethiopia: A systematic review and meta-analysis. *PLoS One*. 2020;15(12):e0243776.
42. Yemane TT, Bogale GG, Egata G, Tefera TK. Postpartum Family Planning Use and Its Determinants among Women of the Reproductive Age Group in Low-Income Countries of Sub-Saharan Africa: A Systematic Review and Meta-Analysis. *Int J Reprod Med*. 2021;2021:5580490.
43. Nelson HD, Cantor A, Jungbauer RM, Eden KB, Darney B, Ahrens K, et al. Effectiveness and Harms of Contraceptive Counseling and Provision Interventions for Women : A Systematic Review and Meta-analysis. *Ann Intern Med*. 2022;175(7):980-93.
44. Sothornwit J, Kaewrudee S, Lumbiganon P, Pattanittum P, Averbach SH. Immediate versus delayed postpartum insertion of contraceptive implant and IUD for contraception. *Cochrane Database Syst Rev*. 2022;10(10):CD011913.
45. Hu D, Tang Y, Pei K. Strategies for Improving Postpartum Contraception Compared With Routine Maternal Care: A Systematic Review and Meta-Analysis. *Int J Public Health*. 2023;68:1605564.

46. Nabhan A, Kabra R, Allam N, Ibrahim E, Abd-Elmonem N, Wagih N, et al. Implementation strategies, facilitators, and barriers to scaling up and sustaining post pregnancy family planning, a mixed-methods systematic review. *BMC Womens Health*. 2023;23(1):379.