

Supplementary material




Figure S1: DEGREE centre locations within countries*

Table S1: Definitions of variables and any differences between studies

Centre	Hypertension definition	Diabetes definition	Heavy proteinuria definition	eGFR calculation	Other
Chile	Self-report of diagnosis or mean (of 2) SBP \geq 140mmHg or DBP \geq 90mmHg or use of antihypertensive drugs in past 2 weeks	Self-report or glycemia \geq 126mg/dL or use of hypoglycemic drugs	urine ACR \geq 300mg/g	CKD-EPI 2021	Tables provided
Ecuador	SBP \geq 140 or DBP \geq 90 mmHg in two different measurements	HbA1c \geq 6.5%.	urine dipstick protein \geq +++	CKD-EPI 2009 (with race) for eGFR _{creat}	Tables provided.
England	Self report of diagnosis, or mean (of 3) SBP \geq 140 mmHg or DBP \geq 90mmHg	Self report of diagnosis or HbA1c \geq 6.5%	urine ACR $>$ 300mg/g	CKD-EPI 2009 (with race) for eGFR _{creat} and CKD-EPI 2012 for eGFR _{cys}	Nationally representative nationwide survey. Age groups are one year out i.e. 18-29, 30-39, 40-49, 50-59.
Guatemala	Self report of diagnosis, or mean (of 3) SBP \geq 140mmHg or DBP \geq 90mmHg	Self report of diagnosis or HbA1c \geq 6.5%	urine ACR $>$ 300mg/g	CKD-EPI 2009 (non race) for eGFR _{creat}	
India 1 (CARRS)	Self-report of diagnosis or mean SBP \geq 140mmHg or DBP \geq 90 mmHg	Self report of diagnosis or HbA1c \geq 6.5% or fasting glucose \geq 126mg/dl	urine ACR $>$ 300mg/g	CKD-EPI 2009 (non race) for eGFR _{creat}	Age $<$ 20 years excluded. Pregnant, bedridden and participants who were unable to comprehend the questionnaires due cognitive deficiencies were excluded.
India 2 (ICMR)	Self-report of diagnosis or mean SBP \geq 140mmHg or DBP \geq 90mmHg	Self report of diagnosis or HbA1c \geq 6.5% or fasting glucose \geq 126mg/dl	urine ACR $>$ 300mg/g	CKD-EPI 2009 (non race) for eGFR _{creat}	Age $<$ 30 years excluded. Pregnant, bedridden and participants who were unable to comprehend the questionnaires due cognitive deficiencies were excluded.
India 3 (UDAY)	Self-report of diagnosis or mean SBP \geq 140mmHg or DBP \geq 90mmHg	Self report of diagnosis or HbA1c \geq 6.5% or fasting glucose \geq 126mg/dl	urine ACR $>$ 300mg/g	CKD-EPI 2009 (non race) for eGFR _{creat} and CKD-EPI 2012 for eGFR _{cys}	Age $<$ 30 years excluded. Pregnant, bedridden and participants who were unable to comprehend the questionnaires due cognitive deficiencies were excluded.
India 4 (Uddanam)	Self-report of $>$ 5 year hypertension	HbA1c \geq 6.5%	urine PCR $>$ 150mg/g	CKD-EPI 2012 (non race) for eGFR _{creat}	Tables provided.
India 5 (Prakasham)	SBP \geq 140mmHg or DBP \geq 90mmHg	HbA1c \geq 6.5%	urine ACR \geq 300mg/g	CKD-EPI 2021 (non-race) for eGFR _{creat}	Tables provided.
Italy	Self-report of diagnosis or SBP \geq 140mmHg or DBP \geq 90mmHg	Self-report of diagnosis	urine dipstick protein \geq ++	CKD-EPI 2009 (non race) for eGFR _{creat}	
Kenya	Self-report of diagnosis or medication for hypertension, mean SBP \geq 140mmHg or mean DBP \geq 90mmHg	Self-report of diagnosis or diabetic medication	urine dipstick protein \geq ++	CKD-EPI 2009 (non race) for eGFR _{creat} and CKD-EPI 2012 for eGFR _{cys}	

Malawi	Self-report of diagnosis, or hypertensive medication in last 2 weeks, or mean SBP \geq 140mmHg or DBP \geq 90mmHg	Glucose \geq 7mmol/L, or self-report previous diagnosis or diabetic medication	urine ACR $>$ 300mg/g	CKD-EPI 2009 (non race) for eGFR _{creat} and CKD-EPI 2012 for eGFR _{cys}	
Nepal	Diagnosis (verified through medical records provided by participant) or mean SBP \geq 140mmHg or DBP \geq 90mmHg	Taking diabetic medication, or fasting glucose \geq 126mg/dL or post prandial blood glucose levels \geq 200mg/dL	Urine protein \geq 300 mg/dL	CKD-MDRD for eGFR _{creat}	Nationally representative nationwide survey. Too frail participants and those who cannot provide written consent were excluded.
Nicaragua 1	Self-report of diagnosis, or SBP \geq 140mmHg or DBP \geq 90mmHg	Self-reported diagnosis or glucosuria \geq 100mg/dL using a urine stick	urine dipstick protein \geq +++	CKD-MDRD for eGFR _{creat}	Tables provided.
Nicaragua 2	Self-report of diagnosis or mean SBP \geq 140mmHg or DBP \geq 90mmHg	Self-reported diagnosis or serum glucose \geq 110mg/dL	urine dipstick protein \geq +++	CKD-EPI 2012 (with race) for eGFR _{creat}	Tables provided. Stratified by urban/rural to match the population rate
Peru	Self-report of diagnosis, or mean SBP \geq 140mmHg or DBP \geq 90mmHg	Self reported diagnosis, or taking diabetes medication, or fasting glucose \geq 126mg/dL	urine dipstick protein \geq ++	CKD-EPI 2009 for eGFR _{creat} and CKD-EPI 2012 for eGFR _{cys}	Stratified by urban/rural to 50% and sex. Female recruitment stopped after reaching 60% for each subgroup.
Sri Lanka	Self report of taking medication for hypertension, or mean SBP \geq 140mmHg or DBP \geq 90mmHg	Self-reported diagnosis, or current diabetic medication, or fasting glucose \geq 126mg/dL	urine ACR $>$ 300 mg/g	CKD-EPI 2009 for eGFR _{creat}	Stratified by urban/rural (almost 50% in each group) and by sex. Female recruitment stopped after reaching 60% for each subgroup.
Thailand	Self report of taking medication for hypertension, or mean SBP \geq 140mmHg or DBP \geq 90mmHg	Use of antihypoglycemic medications, or fasting plasma glucose \geq 126mg/dL	urine dipstick protein \geq ++	CKD-EPI 2009 (non race)	Tables provided. Stratified, nationally representative, nationwide survey.
USA	Self-report of diagnosis, or mean (of 3) SBP \geq 140mmHg or DBP \geq 90mmHg	Self report of diagnosis or HbA1c \geq 6.5%	urine ACR $>$ 300mg/g	CKD-EPI 2009 (non race) for eGFR _{creat}	Nationally representative nationwide survey. Urban/rural flag unavailable.

SBP=systolic blood pressure; DBP=diastolic blood pressure; ACR=albumin creatinine ratio; HbA1c=glycated haemoglobin; CKD-EPI=The Chronic Kidney Disease Epidemiology Collaboration; CKD-MDRD=Chronic Kidney Disease Modification of Diet in Renal Disease

Table S2: Age-standardised prevalence of eGFR<60 ml/min/1.73m² in total sample and complete data sample, by sex

Centre	Area	Rural / Urban	Total Sample men	Men with missing data	Total sample prevalence of eGFR<60 ^a in men % (95% CI)	Complete case prevalence of eGFR<60 ^a in men % (95% CI)	Total Sample women	Women with missing data	Total sample prevalence of eGFR<60 ^a in women % (95% CI)	Complete case prevalence of eGFR<60 ^a in women % (95% CI)
Chile	Molina	Rural	42	3	0.0 (N/A)	0.0 (N/A)	18	1	0.0 (N/A)	0.0 (N/A)
Chile	Molina	Urban	167	11	0.0 (N/A)	0.0 (N/A)	283	19	1.5 (0.1, 2.9)	1.6 (0.1, 3.1)
Ecuador	Miguelillo	Rural	312	0	2.2 (0.7, 3.8)	2.2 (0.7, 3.8)	442	0	6.4 (4.3, 8.5)	6.4 (4.3, 8.5)
England	all	Rural	167	6	0.4 (0, 1.0)	0.4 (0, 1.0)	233	10	2.1 (0.7, 3.5)	2.2 (0.8, 3.7)
England	all	Urban	789	45	0.6 (0.1, 1.0)	0.5 (0.1, 0.9)	1063	56	1.0 (0.6, 1.5)	1.0 (0.6, 1.5)
Guatemala	San Antonio Suchitepequez	Rural	119	4	3.6 (0.4, 6.8)	3.7 (0.4, 7.0)	224	3	1.3 (0, 2.8)	1.3 (0, 2.8)
Guatemala	Tecpan	Rural	116	7	0.8 (0, 2.4)	0.9 (0, 2.6)	236	27	0.0 (N/A)	0.0 (N/A)
India 1 (CARRS)	Chennai	Urban	2374	41	0.9 (0.5, 1.2)	0.9 (0.5, 1.3)	3108	75	0.6 (0.3, 0.9)	0.6 (0.3, 0.9)
India 1 (CARRS)	Delhi	Urban	1791	58	1.0 (0.6, 1.5)	1.0 (0.6, 1.5)	1890	59	1.2 (0.7, 1.6)	1.2 (0.8, 1.6)
India 2 (ICMR)	Delhi	Urban	864	27	1.2 (0.6, 1.8)	1.2 (0.6, 1.9)	1102	51	2.3 (1.5, 3.1)	2.3 (1.5, 3.1)
India 2 (ICMR)	Faridabad	Rural	670	41	1.6 (0.9, 2.4)	1.6 (0.8, 2.4)	845	61	1.6 (0.9, 2.4)	1.6 (0.9, 2.4)
India 3 (UDAY)	Sonipat	Rural	778	10	0.6 (0.2, 1.1)	0.6 (0.2, 1.1)	1184	48	0.6 (0.2, 1.0)	0.6 (0.2, 1.0)
India 3 (UDAY)	Sonipat	Urban	1046	8	0.9 (0.4, 1.3)	0.9 (0.4, 1.3)	1206	22	0.6 (0.2, 0.9)	0.6 (0.2, 0.9)
India 3 (UDAY)	Vizag	Rural	1019	85	6.1 (3.0, 9.1)	6.7 (2.9, 10.4)	1349	107	3.3 (2.2, 4.3)	3.1 (2.2, 4.1)
India 3 (UDAY)	Vizag	Urban	935	32	1.2 (0.6, 1.8)	1.2 (0.6, 1.8)	1193	63	0.7 (0.3, 1.1)	0.7 (0.3, 1.1)
India 4 (Uddanam)	Kanchili	Rural	148	0	4.8 (1.1, 8.6)	4.8 (1.1, 8.6)	169	0	7.2 (3.2, 11.3)	7.2 (3.2, 11.3)
India 4 (Uddanam)	Kaviti	Rural	102	0	12.2 (7.7, 16.7)	12.2 (7.7, 16.7)	110	0	8.0 (3.5, 12.4)	8.0 (3.5, 12.4)
India 4 (Uddanam)	Mandasa	Rural	95	0	18.4 (10.7, 26.1)	18.4 (10.7, 26.1)	105	0	10.6 (6.0, 15.1)	10.6 (6.0, 15.1)
India 4 (Uddanam)	Palasa	Rural	186	0	11.2 (5.9, 16.5)	11.2 (5.9, 16.5)	176	0	11.0 (6.1, 15.9)	11.0 (6.1, 15.9)
India 4 (Uddanam)	Sompeta	Rural	202	0	9.2 (4.9, 13.4)	9.2 (4.9, 13.4)	241	0	2.7 (0.9, 4.5)	2.7 (0.9, 4.5)
India 4 (Uddanam)	V_kothuru	Rural	250	0	5.3 (3.0, 7.5)	5.3 (3.0, 7.5)	281	0	4.9 (2.9, 7.0)	4.9 (2.9, 7.0)
India 5 (Prakasham)	Kanigiri	Rural	420	0	5.3 (3.4, 7.2)	5.3 (3.4, 7.2)	632	0	3.4 (2.0, 4.7)	3.4 (2.0, 4.7)
Italy	Barga	Rural	130	2	0.9 (0, 2.3)	0.9 (0, 2.3)	176	3	0.7 (0, 1.7)	0.7 (0, 1.7)
Kenya	Muhoroni East	Urban	138	0	0.0 (N/A)	0.0 (N/A)	122	0	0.9 (0, 2.7)	0.9 (0, 2.7)
Kenya	Owaga	Rural	138	25	0.0 (N/A)	0.0 (N/A)	151	22	2.1 (0, 4.5)	2.2 (0, 4.8)
Kenya	Tonde	Rural	114	0	0.0 (N/A)	0.0 (N/A)	119	0	0.0 (N/A)	0.0 (N/A)
Malawi	Karonga	Rural	286	15	3.0 (0.6, 5.3)	3.0 (0.7, 5.4)	382	7	3.0 (1.0, 4.9)	3.0 (1.0, 5.1)

Malawi	Lilongwe	Urban	96	0	2.2 (0, 6.2)	2.2 (0, 6.2)	216	0	4.9 (1.4, 8.4)	4.9 (1.4, 8.4)
Nepal	all	Rural	1796	106	0.9 (0.5, 1.3)	0.9 (0.6, 1.3)	3151	361	2.1 (1.6, 2.6)	2.1 (1.6, 2.6)
Nepal	all	Urban	1683	81	0.8 (0.4, 1.1)	0.8 (0.4, 1.1)	3160	326	3.0 (2.4, 3.5)	3.0 (2.4, 3.6)
Nicaragua 1	Chinandega (banana/sugarcane)	Rural	155	0	19.0 (12.5, 25.5)	19.0 (12.5, 25.5)	176	0	3.5 (0.6, 6.4)	3.5 (0.6, 6.4)
Nicaragua 1	Chinandega (service)	Rural	50	0	0.0 (N/A)	0.0 (N/A)	90	0	0.0 (N/A)	0.0 (N/A)
Nicaragua 1	Leon (coffee)	Rural	40	0	6.3 (0, 14.4)	6.3 (0, 14.4)	37	0	0.0 (N/A)	0.0 (N/A)
Nicaragua 1	Leon (fishing)	Rural	76	0	10.2 (3.2, 17.3)	10.2 (3.2, 17.3)	90	0	2.1 (0, 5.9)	2.1 (0, 5.9)
Nicaragua 1	Leon (mining/subsistence farming)	Rural	158	0	16.2 (10.4, 22.0)	16.2 (10.4, 22.0)	224	0	4.8 (1.7, 7.9)	4.8 (1.7, 7.9)
Nicaragua 2	Leon municipalities	Rural	247	0	15.3 (10.9, 19.7)	15.3 (10.9, 19.7)	329	0	3.1 (1.2, 5.0)	3.1 (1.2, 5.0)
Nicaragua 2	Leon municipalities	Urban	400	0	10.0 (7.2, 12.7)	10.0 (7.2, 12.7)	696	0	3.6 (2.3, 4.8)	3.6 (2.3, 4.8)
Peru	Tumbes	Rural	281	3	0.5 (0, 1.3)	0.5 (0, 1.3)	349	5	0.3 (0, 0.9)	0.3 (0, 0.9)
Peru	Tumbes	Urban	261	4	0.5 (0, 1.2)	0.5 (0, 1.2)	360	1	0.5 (0, 1.1)	0.5 (0, 1.1)
Sri Lanka	Halambagaswewa	Rural	246	4	8.2 (5.5, 10.9)	8.3 (5.5, 11.0)	504	7	3.8 (2.2, 5.3)	3.8 (2.2, 5.3)
Sri Lanka	Lolugaswewa	Rural	221	0	5.8 (3.5, 8.1)	5.8 (3.5, 8.1)	576	7	4.5 (3.1, 6.0)	4.5 (3.1, 5.9)
Sri Lanka	Pothana	Rural	201	7	4.9 (2.6, 7.2)	5.0 (2.6, 7.3)	504	7	1.6 (0.7, 2.6)	1.7 (0.7, 2.6)
Sri Lanka	Puhudivula	Rural	224	2	7.6 (5.2, 10.1)	7.7 (5.2, 10.2)	580	4	3.6 (2.3, 4.9)	3.6 (2.3, 5.0)
Sri Lanka	Sangilikandarawa	Rural	276	6	6.2 (3.9, 8.5)	6.1 (3.8, 8.4)	558	10	3.1 (1.8, 4.4)	2.9 (1.6, 4.3)
Thailand	Bangkok	Urban	381	0	1.1 (0.1, 2.2)	1.1 (0.1, 2.2)	1223	0	0.7 (0.0, 1.3)	0.7 (0.0, 1.3)
Thailand	Central	Rural	606	0	0.9 (0.3, 1.4)	0.9 (0.3, 1.4)	795	0	1.1 (0.4, 1.9)	1.1 (0.4, 1.9)
Thailand	Central	Urban	519	0	0.6 (0.1, 1.1)	0.6 (0.1, 1.1)	832	0	0.4 (0.1, 0.7)	0.4 (0.1, 0.7)
Thailand	North	Rural	668	0	1.1 (0.6, 1.7)	1.1 (0.6, 1.7)	730	0	1.0 (0.1, 1.8)	1.0 (0.1, 1.8)
Thailand	North	Urban	445	0	1.1 (0, 2.3)	1.1 (0, 2.3)	604	0	1.0 (0.4, 1.6)	1.0 (0.4, 1.6)
Thailand	North East	Rural	577	0	1.3 (0.5, 2.1)	1.3 (0.5, 2.1)	637	0	0.8 (0.3, 1.3)	0.8 (0.3, 1.3)
Thailand	North East	Urban	492	0	0.7 (0, 1.4)	0.7 (0, 1.4)	609	0	0.6 (0.2, 1.0)	0.6 (0.2, 1.0)
Thailand	South	Rural	549	0	0.5 (0.1, 1.0)	0.5 (0.1, 1.0)	678	0	0.2 (0, 0.5)	0.2 (0, 0.5)
Thailand	South	Urban	303	0	0.7 (0.0, 1.4)	0.7 (0.0, 1.4)	489	0	1.1 (0.3, 2.0)	1.1 (0.3, 2.0)
USA	all	All	1727	62	2.0 (1.4, 2.6)	2.0 (1.4, 2.6)	1924	72	1.2 (0.8, 1.6)	1.2 (0.8, 1.7)

eGFR=estimated glomerular filtration rate; CI=confidence interval using normal approximation ^a age-standardised prevalence using WHO global population age weights;

Table S3: Age-standardised prevalence of eGFR<90 ml/min/1.73m² by sex (for age 18-60)

Centre	Area	Rural / Urban	Sample with complete data				Sample of people without hypertension, diabetes and heavy proteinuria			
			Men		Women		Men		Women	
			n	eGFR<90 ^a % (95% CI)	n	eGFR<90 ^a % (95% CI)	n	eGFR<90 ^a % (95% CI)	n	eGFR<90 ^a % (95% CI)
Chile	Molina	Rural	39	36.2 (17.3, 55.2)	17	59.1 (35.7, 82.4)	16	35.9 (9.0, 62.8)	6	100 (NA, NA)
Chile	Molina	Urban	156	42.1 (34.2, 50.0)	264	49.0 (42.8, 55.8)	66	47.5 (36.0, 59.0)	137	46.5 (37.7, 55.2)
Ecuador	Miguelillo	Rural	312	34.1 (29.2, 39.1)	442	46.3 (41.9, 50.6)	180	33.4 (26.6, 40.3)	235	45.2 (38.6, 51.9)
England	all	Rural	161	35.7 (27.7, 43.8)	223	36.1 (29.7, 42.5)	98	35.8 (26.8, 44.7)	169	37.3 (29.9, 44.8)
England	all	Urban	744	37.5 (33.9, 41.2)	1007	31.7 (28.9, 34.5)	515	38.1 (34.0, 42.2)	759	32.9 (29.8, 35.9)
Guatemala	San Antonio Suchitepequez	Rural	115	17.2 (10.4, 24.1)	221	11.0 (6.8, 15.1)	86	14.8 (6.7, 22.8)	171	10.3 (5.1, 15.6)
Guatemala	Tecpan	Rural	109	26.6 (19.0, 34.3)	209	14.1 (9.7, 18.5)	83	24.7 (15.4, 33.9)	152	15.0 (8.9, 21.1)
India 1 (CARRS)	Chennai	Urban	2333	8.2 (7.1, 9.2)	3033	5.8 (5.0, 6.7)	1161	6.0 (4.6, 7.5)	1915	5.4 (4.1, 6.7)
India 1 (CARRS)	Delhi	Urban	1733	11.4 (10.1, 12.7)	1831	11.9 (10.6, 13.1)	770	10.8 (8.6, 12.9)	935	11.1 (8.9, 13.2)
India 2 (ICMR)	Delhi	Urban	837	33.3 (27.8, 38.8)	1051	37.0 (33.9, 40.0)	399	33.2 (25.8, 40.7)	571	36.1 (32.2, 39.9)
India 2 (ICMR)	Faridabad	Rural	629	18.6 (16.4, 20.8)	784	18.1 (12.8, 23.4)	380	18.6 (15.6, 21.5)	520	17.2 (11.0, 23.5)
India 3 (UDAY)	Sonipat	Rural	768	15.3 (9.1, 21.4)	1136	12.3 (10.0, 14.5)	530	13.7 (7.7, 19.8)	847	12.5 (10.1, 15.0)
India 3 (UDAY)	Sonipat	Urban	1038	12.0 (10.5, 13.4)	1184	11.2 (9.2, 13.2)	586	10.9 (9.0, 12.8)	768	9.9 (7.8, 12.0)
India 3 (UDAY)	Vizag	Rural	934	30.6 (24.6, 36.6)	1242	20.1 (17.6, 22.6)	696	30.2 (23.4, 37.0)	933	19.1 (16.4, 21.8)
India 3 (UDAY)	Vizag	Urban	903	28.3 (23.1, 33.5)	1130	14.6 (12.8, 16.5)	469	27.8 (21.5, 34.0)	692	12.8 (10.2, 15.4)
India 4 (Uddanam)	Kanchili	Rural	148	11.3 (6.3, 16.2)	169	15.8 (10.2, 21.4)	71	13.0 (5.6, 20.5)	83	13.9 (6.7, 21.1)
India 4 (Uddanam)	Kaviti	Rural	102	19.7 (14.4, 24.9)	110	20.0 (13.1, 26.9)	52	12.5 (5.2, 19.7)	65	14.0 (7.5, 20.5)
India 4 (Uddanam)	Mandasa	Rural	95	24.2 (16.1, 32.3)	105	18.5 (13.1, 24.0)	57	18.3 (8.5, 28.0)	54	12.5 (6.7, 18.3)
India 4 (Uddanam)	Palasa	Rural	186	21.4 (15.5, 27.4)	176	21.9 (15.5, 28.3)	84	17.4 (9.2, 25.6)	81	15.5 (8.5, 22.5)
India 4 (Uddanam)	Sompeta	Rural	202	16.2 (11.1, 21.4)	241	12.7 (8.4, 17.0)	99	14.0 (6.9, 21.0)	127	17.1 (10.3, 23.9)
India 4 (Uddanam)	V_kothuru	Rural	250	14.7 (10.9, 18.5)	281	13.7 (10.6, 16.8)	101	11.4 (5.6, 17.3)	121	11.6 (6.5, 16.7)
India 5 (Prakasham)	Kanigiri	Rural	420	27.8 (23.7, 31.8)	632	14.4 (11.8, 17.0)	221	25.5 (20.0, 31.1)	432	11.7 (8.3, 15.1)
Italy	Barga	Rural	128	31.5 (23.9, 39.2)	173	38.0 (31.2, 44.8)	73	31.5 (21.8, 41.2)	149	36.1 (29.3, 43.0)
Kenya	Muhoroni East	Urban	138	8.1 (4.0, 12.3)	122	20.5 (12.1, 28.9)	113	8.6 (3.6, 13.6)	104	21.6 (17.5, 25.6)
Kenya	Owaga	Rural	113	10.3 (4.4, 16.1)	129	16.1 (10.5, 21.7)	88	8.5 (1.3, 15.8)	98	17.9 (11.6, 24.2)
Kenya	Tonde	Rural	114	3.6 (0.5, 6.7)	119	7.2 (2.6, 11.9)	94	3.5 (0.2, 6.7)	100	8.2 (2.5, 13.8)
Malawi	Karonga	Rural	271	17.6 (12.9, 22.3)	375	14.5 (10.3, 18.8)	214	15.0 (9.5, 20.4)	309	13.8 (9.1, 18.5)

Malawi	Lilongwe	Urban	96	24.4 (13.9, 34.9)	216	24.7 (18.1, 31.3)	74	27.4 (16.2, 38.6)	159	31.3 (18.0, 44.6)
Nepal	all	Rural	1690	31.4 (29.2, 33.7)	2790	44.1 (42.4, 45.9)	1002	29.3 (26.6, 32.0)	1981	43.2 (41.2, 45.2)
Nepal	all	Urban	1602	40.0 (37.5, 42.6)	2834	54.9 (53.0, 56.8)	822	38.5 (35.3, 41.7)	1777	53.8 (51.6, 56.1)
Nicaragua 1	Chinandega (banana/sugarcane)	Rural	155	32.9 (25.5, 40.3)	176	6.5 (2.8, 10.3)	104	25.1 (16.4, 33.8)	111	2.2 (0, 4.6)
Nicaragua 1	Chinandega (service)	Rural	50	9.3 (1.3, 17.2)	90	2.4 (0, 5.5)	34	9.2 (0, 20.2)	58	2.4 (0, 6.8)
Nicaragua 1	Leon (coffee)	Rural	40	8.6 (0, 17.8)	37	5.6 (0, 14.4)	30	11.2 (0, 22.8)	26	0.0 (N/A)
Nicaragua 1	Leon (fishing)	Rural	76	31.2 (21.4, 40.9)	90	11.6 (4.5, 18.8)	55	24.5 (12.0, 36.9)	73	6.4 (1.2, 11.7)
Nicaragua 1	Leon (mining/subsistence farming)	Rural	158	41.2 (34.2, 48.2)	224	13.5 (8.8, 18.2)	106	35.9 (26.1, 45.6)	144	7.3 (1.7, 12.8)
Nicaragua 2	Leon municipalities	Rural	247	28.3 (23.0, 33.6)	329	11.5 (8.0, 14.9)	145	23.5 (16.8, 30.2)	211	9.3 (4.3, 14.3)
Nicaragua 2	Leon municipalities	Urban	400	21.9 (18.4, 25.4)	696	13.0 (10.8, 15.2)	256	15.9 (11.2, 20.6)	436	9.1 (6.5, 11.8)
Peru	Tumbes	Rural	278	7.5 (5.0, 10.0)	344	4.0 (2.0, 6.0)	210	4.5 (2.1, 7.0)	285	4.9 (2.0, 7.7)
Peru	Tumbes	Urban	257	15.5 (11.7, 19.3)	359	5.2 (3.2, 7.3)	186	14.4 (9.8, 19.0)	305	4.6 (2.3, 6.8)
Sri Lanka	Halambagaswewa	Rural	242	33.6 (28.6, 38.6)	497	29.1 (25.9, 32.4)	136	31.7 (25.1, 38.2)	336	26.5 (22.6, 30.4)
Sri Lanka	Lolugaswewa	Rural	221	39.5 (33.2, 45.8)	569	34.5 (31.4, 37.6)	138	38.1 (31.3, 44.9)	372	34.0 (30.3, 37.6)
Sri Lanka	Pothana	Rural	194	26.5 (22.0, 31.0)	497	30.3 (27.1, 33.6)	115	26.3 (20.1, 32.5)	330	29.6 (25.6, 33.6)
Sri Lanka	Puhudivula	Rural	222	46.3 (40.6, 51.9)	576	46.0 (42.5, 49.5)	112	49.0 (42.7, 55.4)	378	44.3 (39.9, 48.6)
Sri Lanka	Sangilikandarawa	Rural	270	30.4 (25.5, 35.3)	548	29.5 (26.2, 32.7)	157	27.7 (21.6, 33.8)	346	27.9 (23.8, 31.9)
Thailand	Bangkok	Urban	381	18.8 (15.0, 22.6)	1223	6.6 (5.4, 7.9)	233	20.3 (15.4, 25.2)	943	6.4 (5.0, 7.8)
Thailand	Central	Rural	606	11.6 (9.4, 13.9)	795	10.6 (8.6, 12.6)	424	11.4 (8.7, 14.2)	576	10.0 (7.7, 12.3)
Thailand	Central	Urban	519	15.9 (12.9, 18.8)	832	8.2 (6.6, 9.7)	336	15.0 (11.5, 18.5)	587	7.7 (6.0, 9.5)
Thailand	North	Rural	668	11.0 (8.8, 13.2)	730	7.4 (5.7, 9.2)	407	10.1 (7.4, 12.7)	498	7.2 (5.2, 9.2)
Thailand	North	Urban	445	16.3 (12.5, 20.1)	604	9.1 (7.0, 11.1)	267	15.2 (10.8, 19.7)	409	8.6 (6.2, 11.0)
Thailand	North East	Rural	577	12.5 (10.5, 14.6)	637	33.2 (29.0, 37.5)	428	11.9 (9.6, 14.2)	479	33.1 (28.5, 37.7)
Thailand	North East	Urban	492	16.4 (14.0, 18.8)	609	42.1 (37.2, 46.9)	351	14.7 (12.0, 17.5)	450	38.8 (33.7, 43.9)
Thailand	South	Rural	549	9.9 (7.7, 12.0)	678	4.3 (2.9, 5.6)	366	10.0 (7.2, 12.7)	493	4.0 (2.5, 5.6)
Thailand	South	Urban	303	14.5 (11.1, 17.9)	489	7.7 (5.6, 9.8)	193	13.7 (9.5, 17.9)	356	7.3 (4.8, 9.7)
USA	all	All	1665	25.7 (23.7, 27.6)	1852	17.2 (15.6, 18.7)	944	27.7 (25.0, 30.4)	1163	16.5 (14.5, 18.6)

^a age-standardised prevalence using WHO global population age weights; ^b only 41-60 years included; eGFR=creatinine-based estimated glomerular filtration rate; CI=confidence interval using normal approximation; N/A=no CI available due to zero estimate

Table S4: Age-standardised prevalence of creatinine- and cystatin C-based eGFR<90 ml/min/1.73m², for age 18-60 years, by sex, for participants with both measurements

Centre	Area	Rural / Urban	Sample with complete data						Sample of people without hypertension, diabetes, or heavy proteinuria					
			Men			Women			Men			Women		
			n	Creatinine eGFR<90 ^a % (95% CI)	Cystatin C eGFR<90 ^a % (95% CI)	n	Creatinine eGFR<90 ^a % (95% CI)	Cystatin C eGFR<90 ^a % (95% CI)	n	Creatinine eGFR<90 ^a % (95% CI)	Cystatin C eGFR<90 ^a % (95% CI)	n	Creatinine eGFR<90 ^a % (95% CI)	Cystatin C eGFR<90 ^a % (95% CI)
England	all	Rural	161	35.7 (27.7, 43.8)	32.3 (23.3, 41.4)	223	36.1 (29.7, 42.5)	18.3 (14.2, 22.4)	98	35.8 (26.8, 44.7)	32.4 (22.4, 42.4)	169	37.3 (29.9, 44.8)	15.6 (11.2, 20.0)
England	all	Urban	744	37.5 (33.9, 41.2)	25.9 (22.9, 29.0)	1007	31.7 (28.9, 34.5)	23.0 (20.4, 25.6)	515	38.1 (34.0, 42.2)	23.9 (20.5, 27.3)	759	32.9 (29.8, 35.9)	20.0 (17.2, 22.8)
India 3 (UDAY) ^b	Sonipat	Rural	-	-	-	-	-	-	177	25.3 (9.2, 41.3)	69.2 (51.0, 87.5)	253	11.5 (7.4, 15.6)	59.7 (51.5, 67.9)
India 3 (UDAY) ^b	Sonipat	Urban	-	-	-	-	-	-	199	10.1 (6.9, 13.4)	70.0 (58.1, 81.9)	273	8.7 (4.3, 13.2)	57.4 (48.8, 65.9)
India 3 (UDAY) ^b	Vizag	Rural	-	-	-	-	-	-	269	30.9 (19.4, 42.5)	78.6 (67.3, 90.0)	325	19.1 (14.4, 23.7)	60.4 (53.9, 66.8)
India 3 (UDAY) ^b	Vizag	Urban	-	-	-	-	-	-	152	29.2 (16.9, 41.5)	66.4 (53.1, 79.8)	244	13.2 (8.8, 17.6)	56.9 (50.5, 63.3)
Kenya	Muhoroni East	Urban	4	0.0 (N/A)	0.0 (N/A)	10	7.2 (0, 18.6)	14.3 (2.9, 25.8)	4	0.0 (N/A)	0.0 (N/A)	9	0.0 (N/A)	10.8 (0, 25.7)
Kenya	Owaga	Rural	44	7.2 (0, 15.3)	38.4 (23.3, 53.4)	50	21.0 (11.2, 30.8)	31.6 (21.4, 41.7)	37	7.2 (0, 15.3)	27.9 (15.4, 40.3)	43	19.7 (9.1, 30.3)	29.5 (18.7, 40.4)
Kenya	Tonde	Rural	26	11.4 (0, 23.1)	37.4 (26.3, 48.4)	39	2.8 (0, 7.7)	23.2 (14.6, 31.9)	19	7.8 (0, 17.7)	23.4 (10.1, 36.7)	29	5.6 (0, 14.4)	27.0 (14.1, 39.8)
Malawi	Karonga	Rural	271	17.6 (12.9, 22.3)	15.9 (11.4, 20.5)	375	14.5 (10.3, 18.8)	19.9 (15.6, 24.2)	214	15.0 (9.5, 20.4)	15.3 (9.7, 20.8)	309	13.8 (9.1, 18.5)	18.2 (13.1, 23.2)
Malawi	Lilongwe	Urban	96	24.4 (13.9, 34.9)	8.8 (0.8, 16.9)	216	24.7 (18.1, 31.3)	6.8 (2.3, 11.4)	74	27.4 (16.2, 38.6)	11.0 (0.8, 21.2)	159	31.3 (18.0, 44.6)	2.9 (0, 5.8)
Peru	Tumbes	Rural	278	7.5 (5.0, 10.0)	12.8 (9.1, 16.5)	343	4.0 (2.0, 6.0)	15.0 (11.3, 18.7)	210	4.5 (2.1, 7.0)	14.1 (9.7, 18.5)	284	4.9 (2.0, 7.7)	15.4 (11.1, 19.8)
Peru	Tumbes	Urban	257	15.5 (11.7, 19.3)	14.1 (10.1, 18.1)	358	5.2 (3.2, 7.3)	11.6 (8.4, 14.7)	186	14.4 (9.8, 19.0)	14.3 (9.5, 19.0)	304	4.6 (2.3, 6.8)	11.6 (8.1, 15.1)

eGFR=estimated glomerular filtration rate; ^a age-standardised prevalence using WHO global population age weights; ^b complete data not available as only participants without hypertension, diabetes, or heavy proteinuria had cystatin C measured; N/A=no CI available due to zero estimate