

Supplementary Table 1. Prevalence ratios and 95% confidence intervals of preventive dental visits and health literacy after including participants who reported to have diabetes in the past but not currently

	N	Events	Unadjusted model		Adjusted model*	
			PR (95% CI)	P value	PR (95% CI)	P value
All participants						
Low (<4 points)	1,110	557 (50.2)	Reference		Reference	
High (\geq 4 points)	589	341 (57.9)	1.15 (1.05 to 1.27)	0.002	1.09 (0.99 to 1.19)	0.064
1 point increase (continuous)			1.11 (1.05 to 1.20)	0.001	1.07 (1.00 to 1.15)	0.047
Aged <65 years						
Low (<4 points)	513	216 (42.1)	Reference		Reference	
High (\geq 4 points)	273	142 (52.0)	1.24 (1.06 to 1.44)	0.007	1.17 (1.00 to 1.36)	0.055
1 point increase (continuous)			1.13 (1.01 to 1.26)	0.029	1.09 (0.97 to 1.22)	0.15
Aged \geq65 years						
Low (<4 points)	597	341 (57.1)	Reference		Reference	
High (\geq 4 points)	316	199 (63.0)	1.10 (0.99 to 1.23)	0.081	1.06 (0.95 to 1.19)	0.27
1 point increase (continuous)			1.09 (1.00 to 1.19)	0.039	1.06 (0.97 to 1.16)	0.17

Abbreviations: CI, confidence interval; PR, prevalence ratio.

*Adjusted for sex, age, education, marital status, living arrangement, working status, annual household income, self-rated health, and body-mass index.