

1 **MAGNITUDE OF NEONATAL HYPOGLYCEMIA AND ITS ASSOCIATED FACTORS**
2 **AMONG NEONATES ADMITTED TO NEONATAL INTENSIVE CARE UNIT AT**
3 **HAWASSA CITY PUBLIC HOSPITALS, ETHIOPIA, 2023.**

4
5

6 **Selam Tadele^{1*}, Wegene Jembere², Mastewal Aschale², Tewodros Mulugeta¹, Samuel**
7 **Jigso¹, Mequanint Ayehu¹, Migbar Sibhat¹**

8
9
10
11

12 ¹ Department of Nursing, College of Medicine and College of Health Science, Dilla University,
13 Dilla, Ethiopia

14 ² School of nursing, College of medicine and health Sciences, Hawassa University, Hawassa,
15 Ethiopia

16
17
18
19
20
21
22
23

24 *Corresponding author
25 E-mail: selmesty1216@gmail.com

26
27
28
29

30

Abstract

31

32 **Background:** Neonatal hypoglycemia is one of the most common metabolic abnormalities seen
33 in newborns. It is a major contributing factor to neonatal morbidity and mortality. Globally, it affects
34 around 5–15% of all babies and approximately 50% of at-risk babies.. In Ethiopia, neonatal
35 hypoglycemia is frequently diagnosed and one of the commonest causes of admission to the
36 neonatal intensive care unit. Nevertheless, documented records regarding its magnitude and
37 factors associated with hypoglycemia are scarce in the study area. Therefore we aimed to assess
38 the magnitude of neonatal hypoglycemia and its associated factors among neonates admitted to
39 the neonatal intensive care unit at Hawassa City Public Hospitals, Ethiopia.

40 **Method:** Institution-based cross-sectional study was conducted from April 20 – June 20,
41 2023 among 293 neonates. A systematic random sampling technique was used to reach the study
42 subjects. The data was collected through face-to-face interviews and card review by using
43 structured pretested questionnaire and analyzed using SPSS software version 25. A multivariable
44 logistic regression model was used to determine factors significantly associated with neonatal
45 hypoglycemia with adjusted odds ratio, p-values <0.05 at 95% confidence interval (CI).

46 **Result:** The magnitude of neonatal hypoglycemia was found 16.6%. Variables significantly
47 associated with the occurrence of neonatal hypoglycemia were: Diabetes mellitus [AOR=9.8,
48 95%CI (3.08-31.37)], perinatal asphyxia [AOR=2.87, 95%CI (1.07-7.72)], delayed initiation of
49 breastfeeding [AOR=2.63, 95%CI (1.04-6.6)] and hypothermia [AOR=3.8, 95%CI (1.6-9.1)].

50 **Conclusion:** In this study the magnitude of neonatal hypoglycemia among neonates was high.
51 Neonates with hypothermia, perinatal asphyxia, and delayed initiation of breastfeeding and
52 maternal history of diabetes mellitus have an increased risk of developing hypoglycemia. Hence,
53 Health care providers who are working on delivery and neonatal care should focus on early
54 identification and management of these identified factors.

55 **Keywords:** neonate, diabetes mellitus, hypoglycemia, Blood glucose

1. Introduction

56

57 Neonatal hypoglycemia is defined as an abnormally low blood glucose level (1). There has been
58 much controversy about the ‘numerical’ definition of neonatal hypoglycemia. Although the most
59 often used definition of newborn hypoglycemia is a glucose concentration of 47 mg/dl (2.6

60 mmol/l), (2, 3), in Ethiopia it is clinically defined as a blood glucose level less than 40 mg/dl
61 (4)..

62 It is the most prevalent metabolic abnormality in the newborn period (5), which is a major
63 contributing factor to neonatal morbidity and mortality(6). Globally, it affects around 5–15% of
64 all babies (7, 8) and approximately 50% of at-risk babies and is associated with a range of
65 adverse consequences (9)..

66 In resource-limited settings, evidence shows there is widespread neonatal hypoglycemia, even
67 though in some countries there is a shortage of advanced techniques for hypoglycemia detection
68 and an absence of routine assessment for hypoglycemia. In African countries Nigeria and
69 Uganda have an incidence of 32.7% and 2.2 % respectively (6, 10). In Ethiopia, the magnitude
70 ranges from 15%-25% (11, 12).

71 If left untreated, severe and prolonged hypoglycemia can have a devastating neurologic and
72 developmental outcome which includes cerebral palsy, long-term neurodevelopmental
73 disabilities, mental retardation, epilepsy, personality disorders, and even death (13).

74 Despite these devastating health impacts of neonatal hypoglycemia, organized data regarding the
75 magnitude of hypoglycemia and the factors that predict its occurrence were scarce in Ethiopia.
76 Furthermore, we couldn't find a single study to investigate this issue in our study area. Hence,
77 this study aims to determine the magnitude of neonatal hypoglycemia, and factors associated
78 with it among neonates admitted to neonatal intensive care units at Hawassa City Public
79 Hospitals, Sidama Regional State, Ethiopia.

80 **3. Methods and Materials**

81 **3.1. Study design and area**

82 An institutional-based cross-sectional study was conducted in Hawassa City from April 20- June
83 20, 2023. Hawassa city is the capital city of the newly emerged Sidama National Regional State
84 and it is located 275 km south of far from Addis Ababa, Ethiopia. There are 9 hospitals (4 public
85 and 5 private) in Hawassa City, as well as 12 public health centers and 18 health posts. The four
86 public hospitals are Hawassa University Comprehensive Specialized Hospital (14), Adare

87 General Hospital (15), Motite Furra Primary Hospital and Hawela Tula Primary Hospital
88 (HTPH). Only the three hospitals have NICU. All three hospitals follow Ethiopia's National
89 Standard Treatment Guideline for Hospitals(16).

90 **3.2. Source population and Study population**

91 **Source population**

- 92 • All neonates who were admitted to the neonatal intensive care units of Hawassa city
93 public hospitals are the source population

94 **Study population**

- 95 • All randomly selected neonates admitted to the neonatal intensive care unit of selected
96 public hospitals during the study period were the study population

99 **3.3. Inclusion and Exclusion Criteria**

100 All neonates who were admitted to NICU in the selected hospitals during the data collection
101 period were included and neonates admitted without mothers or caregivers were excluded from
102 the study.

103 **3.4. Sample size calculation**

104 The sample size was determined by using single population proportion formula by taking the
105 prevalence of neonatal hypoglycemia from the previous study done in Hiwot Fana Hospital, East
106 Harerge, Ethiopia which is 21.2% (11). Considering the parameters of the single population
107 proportion formula which are a marginal error of 0.05, a 95% confidence interval, and a p-value
108 of 0.21 assuming a 15% non-response rate.

109 $P = \text{proportion hypoglycemia} = 21.2\%$

110 $d = \text{margin of error } 5\%$

111 $Z_{\alpha/2} = \text{the corresponding } Z \text{ score of } 95\% \text{ CI}$

112 $N = \text{Sample size}$

113 $N = \frac{Z^2 (a/2)^2 P (1-P)}{d^2}$

114 $d^2 = 0.0025$

115 $= (1.96)^2 * 0.21 * (1 - 0.21) = 255$

116 By adding a 15 % non-response rate, a total of 293 participants were included in the study.

117 **3.5. Sampling technique**

118 Three public hospitals (Hawassa University Comprehensive Specialized Hospital, Adare General
119 Hospital, and Tulla Primary Hospital) are found in Hawassa City which had NICU. Motite Furra
120 Primary Hospital was excluded because it does not provide NICU service. To select the study
121 participants from each hospital, the proportional allocation formula was used based on caseloads
122 by fixing two months of admission before the survey. Simple random sampling was used to
123 select the first participant and systematic random sampling was used for the next consecutive
124 samples.

125 **3.6. Operational definition**

126 **Magnitude of hypoglycemia:** The number of neonates who develop hypoglycemia at admission
127 to NICU during the study period.

128 **Neonatal hypoglycemia:** - A measure of blood glucose level less than 40 mg/dl(4).

129 **ANC follow up:** any history of visit or follow up during current pregnancy at any health
130 institution for check-up of pregnancy recorded on the chart.

131 **Hypothermia:** A neonate's axillary body temperature below 36.5 o C(4).

132 **Perinatal asphyxia:** was considered when the 5th minute APGAR score is less than 7 or a
133 neonate did not cry immediately at birth or resuscitation needed at birth and documented on the
134 chart(4).

135 **3.7. Data collection tools and procedures**

136 The data was collected using the tool adopted from the previous studies done at Addis Abeba
137 (Tikur Anbessa and St., Paulos), and Harar, Hiwot Fana Hospital (Fikirte Kasaye, 2021). A
138 structured interviewer-based questionnaire and structured checklist was used to collect data for
139 maternal socio-demographic factors and obstetric-related characteristics, and neonatal related
140 factors was taken using a structured checklist from the medical records of a neonate during the
141 data collection period. The data was collected from 20th April, 2023 to June 20, 2023.

142 Regarding the blood glucose level of the neonate, a blood sample was taken from the neonate,
143 and their blood glucose level was measured using a glucometer. Blood glucose was determined

144 using the Precisa Active Blood Glucose Meter at the admission. Blood glucose results were
145 provided in mg/dl. Neonates who were found to be hypoglycemic were managed as per the
146 NICU protocol. For the data collection, three data collectors and one supervisor who were BSC
147 nurses and who had research experience were chosen. Before data collection one-day training
148 was given for data collectors and supervisors.

149 **3.8. Data management and analysis**

150 Data was checked for completeness and consistency and then it was cleaned, coded and entered
151 using Epi data version 3.1 and it was exported to SPSS software version 25 for analysis. Cross-
152 tabulation was done among dependent variables and independent variables. Frequencies,
153 Percentage, mean and standard deviation were used to summarize descriptive statistics. Binary
154 logistic regression and multiple logistic regression were done at p-value <0.25 and <0.05
155 respectively with a 95% Confidence interval. Hosmer and Lomeshow goodness of fit test was
156 done to check model fitness and the model was fitted. In addition, a multi-collinearity or
157 tolerance test was done with all variables to solve the issue of confounding effects among
158 independent variables. Finally tables and charts were used for data presentation.

159 **3.9. Quality assurance technique**

160 The questionnaire was prepared in English and translated to Amharic and sidammu-afu and then
161 retranslated to English to check for inconsistency. The data collection tool was pretested on 5%
162 two weeks before the actual data collection period and modifications were made accordingly.
163 The trained data collectors were assigned to selected hospitals and they collected the data.
164 During the data collection time, close supervision and monitoring was carried out by supervisors
165 and investigator to ensure the quality of the data. Daily evaluations of the data for completeness
166 and encountered difficulties at the time of data collection were attended accordingly. Finally, all
167 the collected data was checked by the supervisor and investigator for completeness and
168 consistency during the data management, storage, and analysis.

169 **3.10. Ethical considerations**

170 Ethical clearance was obtained from the Hawassa University College of Medicine and Health
171 Science institutional review board (Ref Number: IRB/313/15). Then officials at different levels
172 in the Hawassa city administration health office and hospitals were communicated through
173 letters. Since neonates could not speak any information, the neonates' mothers or guardian were

174 informed about the purpose of the study and written informed consent was obtained to confirm
175 willingness. They were notified that they had the right to refuse or terminate at any point of the
176 interview. The study was conducted as per the declaration of Helsinki. Confidentiality of the
177 information was secured throughout the study process.

178 4. Result

179 4.1. Descriptive statistics of the result

180 4.1.1. Socio-demographic characteristics of the respondents

181 A total of 271 neonates were included with an overall response rate of 92.5%. The mean age of
182 the mothers was 29.1 (± 5.29) years. The majority of participants were living in urban areas
183 179(66.6%). Regarding the educational status of the mothers, a higher proportion of mothers
184 134(49.5%) completed secondary educational level. Concerning the marital status, 244 (90%)
185 mothers were married. Whereas, five (1.8%) and thirteen (4.8%) mothers were widowed and
186 divorced respectively. (Table 1)

187 **Table 1:-** Socio-demographic characteristics of the mothers of neonates on admission to the
188 neonatal intensive care unit in Hawassa City Public Hospitals, Ethiopia 2023. (n=271)

Variable	Category	Frequency (n=271)	Percentage (%)
Age group (in a year)	≤ 24	58	21.4
	25-34	163	60.1
	≥ 35	50	18.5
Residence	Urban	179	66.1
	Rural	92	33.9
Educational status	No formal education	67	24.7
	Primary education	70	25.8
	Secondary education	75	27.7
	College and above	59	21.8
Occupation	Government employee	57	21.0
	private employee	51	18.8
	Merchant	50	18.5
	Daily Laborer	17	6.3
	Housewife	86	31.7
	Other*	10	3.7
Marital status	Single	7	2.6

Married	246	90.8
Widowed	5	1.8
Divorced	13	4.8

189 **Key:** Others *= student, street women

190 **4.1.2. Description of maternal and obstetric characteristics**

191 In this study, the majority of 234(86.3%) of the mothers had ANC follow-up during the current
192 pregnancy. Of these, 115(45.8%) had at least four ANC follow up. About 194 (71.9%) mothers
193 had more than one child. Regarding the place of delivery, the majority of the neonates were
194 delivered at health institutions. of which 221(72.3%) were delivered at the hospital followed by
195 59(21.8%) at health centers. Concerning obstetrics history, 14(5.2%) of the mother had
196 prolonged labor and higher proportions of the neonate 166(61.6%) were delivered through
197 spontaneous vaginal delivery. Moreover, 121(44.6%) and 122(45%) of the delivery process were
198 conducted by midwives and medical Doctors (general practitioners, residents and specialists)
199 respectively (Table 2).

200 As shown in Table 2, Around 26(9.6%) of mothers had diabetes mellitus. Among those 13
201 (4.8%) of the mothers had gestational Diabetes mellitus and 13 (4.8%) of them had pre-
202 gestational diabetes mellitus. Regarding pregnancy-induced hypertension, 41(15.1%) of mothers
203 had developed it and 20(7.4%) of the mothers were preeclamptic. Regarding other chronic
204 illnesses, 15(5.5%) of the mothers were hypertensive and 10(3.7%) were HIV/AIDs clients
205 (Table 2).

206

207 **Table 2:** Obstetric-related factors of neonatal hypoglycemia among neonates on admission to
 208 neonatal intensive care unit in Hawassa City public Hospitals, Ethiopia, 2023. (n=271)

Variable	Category	Frequency (n=271)	Percentage
ANC	Had no ANC follow-up	37	13.7
	Had 1-3 ANC follow-up	115	42.4
	Had >4 ANC follow-up	119	43.9
Number of children(parity)	1 child(Primi-parity)	77	28.4
	More than 1 children (Multi-parity)	194	71.9
Prolonged labor	Yes	14	5.2
	No	257	94.8
Place of delivery	Home	16	5.9
	Health Center	59	21.8
	Hospital	196	72.3
Position	Cephalic	221	81.5
	Non-cephalic	50	18.5
Mode of delivery	Spontaneous vaginal delivery	166	61.3
	Assisted vaginal delivery	31	11.4
	Cesarean Section	74	27.3
Premature rupture of membrane	Yes	58	21.4
	No	213	78.6
Type of pregnancy	Single	251	92.6
	Twin and above	20	7.4
Conducted by	Nurse	12	4.4
	Midwives	121	44.6
	Medical Doctor	122	45
	Traditional birth attendant	16	5.9
Diabetes Mellitus	Yes	26	9.6
	No	245	90.4
Pre-pregnancy DM	Yes	13	4.8
	No	258	95.2
Gestational DM	Yes	13	4.8
	No	258	95.2
Pregnancy-induced hypertension	Yes	41	15.1
	No	230	84.9
Preeclampsia	Yes	20	7.4
	No	251	92.6
Eclampsia	Yes	31	11.4
	No	240	88.6
Chronic Hypertension	Yes	15	5.5
	No	256	94.5
HIV/AIDs	Yes	10	3.7
	No	261	96.3

209

210 **4.1.3. Neonatal-related variables**

211 Among admitted neonates around 156 (57.6%) were male and more than half 171 (63.1%) were
212 gestational age between 37-42 weeks. About 112 (41.3%) of them had birth weights between
213 2500gm - 4000gm. The majority of the neonates 216 (79.7%) were admitted within 24hr after
214 delivery. Around, 119 (43.9%) of the neonates had hypothermia. Regarding feeding initiation
215 time 99 (36.5%) of neonates started feeding within one hour (Table 3).

216 Concerning the neonatal-related comorbid illness, 241(88.9%) of neonates were admitted with
217 comorbid illness. Among those 130(48%) had neonatal sepsis, 52(19.2%) had birth asphyxia,
218 38(14%) had respiratory distress syndrome, 30(11.1%) had Meconium Aspiration Syndrome and
219 53(19.6%) had jaundice.

220

221

222

223

224

225

226

227

228

229

230

231

232 **Table 3.** Neonatal-related predictors of hypoglycemia among neonates on admission to neonatal
 233 intensive care unit in Hawassa City public Hospitals, Ethiopia 2023. (n=271)

Variable	Category	Frequency (n=271)	Percentage
Sex	male	156	57.6
	female	115	42.4
GA	<37 weeks	96	35.4
	37-42 week	171	63.1
	>42 weeks	4	1.5
Birth weight	2500-3999	118	43.5
	<2500	112	41.3
	≥4000	28	10.3
Age at admission	<24 hours	213	78.6
	≥24 hours	58	21.4
Hypothermia	Yes	119	43.9
	No	152	56.1
Hypoglycemia	Yes	45	16.6
	No	226	83.4
Time of initiation of breast feeding	≤1 Hours	106	39.1
	>1 Hours	165	60.9
Is there any comorbid illness	Yes	241	88.9
	No	30	11.1
Neonatal sepsis	Yes	130	48.0
	No	141	52.0
Birth asphyxia	Yes	53	19.6
	No	218	80.4
Respiratory distress syndrome	Yes	38	14.0
	No	233	86.0
Meconium Aspiration Syndrome	Yes	30	11.1
	No	241	88.9
Jaundice	Yes	53	19.6
	No	218	80.4
Birth Trauma	Yes	25	9.2
	No	246	90.8
congenital anomaly	Yes	22	8.1
	No	249	91.9

234 **4.2. The Magnitude of neonatal hypoglycemia**

235 In this study, the overall magnitude of neonatal hypoglycemia was observed to be 45(16.6%)
 236 (Figure 1).

237

4.3. Factors associated with Neonatal Hypoglycemia

238 The bivariate logistic regression showed that diabetes mellitus, parity, antenatal care visits,
 239 prolonged labor, and preeclampsia were selected among maternal-related factors. In the same
 240 way, the age of the neonate, birth weight, neonatal sepsis, birth asphyxia, hypothermia, and time
 241 of initiation of breastfeeding were selected from neonatal-related risk factors.

242 The multivariable logistic regression result showed that those mothers who have had Diabetes
 243 mellitus were about ten times to have newborns suffering from neonatal hypoglycemia compared
 244 with those who had not [AOR=9.8, 95% CI(3.08-31.37)]. Neonates born with Birth asphyxia
 245 were nearly three times more likely to be hypoglycemic than their counterpart [AOR=2.87, 95%
 246 CI (1.07-7.72)]. Regarding the time of initiation of breastfeeding, neonates who start
 247 breastfeeding after one hour of delivery were approximately three times more likely to develop
 248 neonatal hypoglycemia than those who start within one hour [AOR=2.63, 95% CI(1.04-6.6)].
 249 Similarly, neonates who develop neonatal hypothermia were nearly four times to be
 250 hypoglycemic than their counterparts [AOR=3.8, 95%CI (1.6-9.1)]. (Table-4)

251 Antenatal care visits, parity, prolonged labor, preeclampsia, birth weight, neonatal sepsis, and
 252 age at admission were not found to have statistically significant association with neonatal
 253 hypoglycemia in this study.

254 **Table 4.** Factors associated with Hypoglycemia among neonates on admission to neonatal
 255 intensive care unit in Hawassa public Hospitals, Ethiopia 2023. (n=271)

Variable	Category	Hypoglycemia (n=271)		COR	AOR
		Yes	No		
DM	1. Yes	14	12	8.05(3.4-18.99)	9.8(3.08-31.37)*
	2. No	31	214	1	1
Age of the neonate	1. < 24hr	32	181	1	1
	2. >24 hr	13	45	1.63(0.79-3.36)	2.407(0.92-6.30)
ANC follow up	1. No ANC followup	10	27	1.95(0.8-4.68)	2.35(0.73-7.62)
	2. 1-3 ANC follow up	16	99	0.85(0.4-1.75)	1.06(0.44-2.53)
	3. > 4 ANC follow up	19	100	1	1

Hypothermia	1. Yes	29	90	2.74(1.41-5.33)	3.8(1.6-9.1)*
	2. No	16	136	1	1
Birth weight	1. 2500-3999	16	102	1	1
	2. <2500	18	94	1.22(0.59-2.53)	0.82(0.32-2.1)
	3. ≥4000	9	19	3.02(1.16-7.83)	1.9(0.52-6.7)
Prolonged labor	1. Yes	4	10	0.48(0.14-1.59)	1.65(0.35-7.56)
	2. No	41	216	1	1
Number of children (parity)	1. 1 child	17	60	1.68(0.86-3.29)	1.34(0.56-3.2)
	2. More than 1 children	28	166	1	1
Preeclampsia	1. Yes	7	13	3.02(1.13-8.05)	2.2(0.64-7.3)
	2. No	38	213	1	1
Perinatal asphyxia	1. Yes	14	39	2.17(1.06-4.45)	2.87(1.07-7.72)*
	2. No	32	187	1	1
Sepsis	1. Yes	26	104	1.6(0.84-3.07)	1.76(0.77-4.05)
	2. No	19	122	1	1
Delayed initiation of breast feeding	1. Yes	37	128	3.54(1.58-7.9)	2.63(1.04-6.6)*
	2. No	8	98	1	1

256 -*p-value<0.05 significantly associated with neonatal hypoglycemia

257 5. Discussion

258 Hypoglycemia is one of a metabolic disturbance which occurs during the neonatal period.
 259 Screening at-risk infants and the management of low blood glucose levels in the first hours to
 260 days of life is a frequent issue in the care of newborn infants. If it is persistent and not detected
 261 early, it could lead to adverse neurological complications and poor prognosis. So this study was
 262 targeted to identify factors associated with neonatal hypoglycemia.

263 In this study, the prevalence of neonatal hypoglycemia at the admission point was 16.6% among
 264 neonates in the Neonatal intensive care unit, in Hawassa city. This study is in line with a study
 265 conducted in West Bengal and India which was 16.3% and 15.2% respectively (18, 19). On the

266 other hand the finding was lower than the study carried out in Ethiopia St. Paul at 25% and
267 Nigeria at 32.7 % (10, 12). This might be due to that these two studies used a higher cut-off point
268 to diagnose neonatal hypoglycemia which is 47mg/dl.

269 In contrast, the magnitude of neonatal hypoglycemia was higher than the study conducted in
270 USA(3.2%)(20), India(8.2%) (21), Nigeria (11%) (22), and Madagascar (3.1%) (23). The
271 possible reasons may be, because our study includes infants of diabetic mothers, neonates born
272 from hypertensive mothers, and newborns with severe congenital malformation which were
273 excluded in the above study areas. Another study done in Uganda showed a 2.2% prevalence,
274 which is much lower compared to the current study (6). The possible explanation for this may be
275 that the study conducted in Uganda was community-based which is different from institutional
276 based and large number of study participants involved in the study.

277 Regarding factors associated with neonatal Hypoglycemia, Diabetes mellitus (gestational
278 Diabetes mellitus Pre-diabetes mellitus), perinatal asphyxia, delayed initiation of breastfeeding
279 and the presence of Hypothermia were independently associated with it.

280 The current study finding revealed that neonates born from mothers with diabetes mellitus were
281 nearly ten times more likely to have hypoglycemia compared with non-diabetic mothers. This
282 study finding was supported by the study done in Ethiopia (Addis Ababa Black Loin Hospital
283 and Harer Hiwot Fana Hospital), South Africa, Bangladesh, and China (11, 17, 24-26). This
284 finding was can be justified by the fact that infants of diabetic mothers have hyper-insulinemia
285 caused by high maternal glucose levels and after birth when maternal glucose is withdrawn the
286 baby may develop hypoglycemia(27).

287 In contrast, the study carried out in St. Paul's, Ethiopia, Indonesia and Israel, maternal DM was
288 not found to be a contributing factor to neonatal hypoglycemia (12, 28, 29). The possible reason
289 for maternal DM to be non-significant may be due to the intervention of early management of
290 maternal DM. Another evidence for this variation may be due to absence or a small number of a
291 mother with DM was found as a study participant.

292 Another factor that was found significantly associated with hypoglycemia was found to be
293 perinatal asphyxia. Neonates born with perinatal asphyxia were nearly three times more likely to

294 have hypoglycemia compared with non-asphyxia neonates. A similar result was also reported in
295 Indonesia which showed that those who have perinatal asphyxia have a three times risk of having
296 hypoglycemia(28). Evidence from china and Colombia suggests that hypoglycemia was
297 significantly related to perinatal asphyxia (30)(Cristo Colmenares et al., 2021). This may be due
298 to the reason in perinatal asphyxia, increasing metabolic and energy demands are achieved
299 transiently by increasing glucose production through glycogenolysis. Due to hypoxia, the infant
300 switches to anaerobic metabolism, resulting in rapid exhaustion of glycogen stores,
301 hypoglycemia, and lactic acedemia(31).

302 Hypothermia was found to be one of the significant risk factors associated with hypoglycemia in
303 this study. Neonates with hypothermia were four times more likely to have hypoglycemia than
304 non-hypothermic. Various evidence show that hypothermia is one of the major risk factor for
305 hypoglycemia (11, 22, 24, 26). In a cross-sectional study in Ethiopia, Hiwot Fana Hospital and
306 St. Paulo's Hospital the odds of developing risk of hypoglycemia in hypothermic neonates were
307 three and two respectively (11, 12). After birth maintaining newborn body temperature is very
308 important. When the baby gets cold the newborn uses up more glycogen to keep warm. Then
309 the newborn must utilize glucose stores to keep warm, then the blood sugar drops and they
310 become hypothermic along with hypoglycemic(32).

311 Delayed initiation of breastfeeding is a major contributing factor to neonatal hypoglycemia.
312 Newborns that started initial breastfeeding after one hour had a three times higher chance of
313 developing neonatal hypoglycemia compared with those who started within one hour. Similarly,
314 research conducted in China, Uganda and Ethiopia Harer, late initiation of breastfeeding has
315 shown that three times higher risk of developing hypoglycemia (6, 11, 24). This similarity can
316 be due to the reason that neonates with delayed initiation of breastfeeding lacks the advantage
317 early initiation of breastfeeding i.e. colostrum increases the blood glucose level within one hour
318 and continuing breast feeding maintain infants at eu-glycemic state(33).

319

320 **6. Strengths and Limitations of the Study**

321 This study used primary data for blood glucose measurement, so it reduces information bias.
322 The limitation of this study is an institution based study it does not address the general
323 community and the study doesn't incorporate laboratorial investigations like RFT, LFT,
324 Bilirubin count etc.

325 **7. Conclusion**

326 In this study, the magnitude of hypoglycemia was 16.6% which is high. The result of this study
327 proved that maternal DM, delayed initiation feeding, hypothermia and perinatal asphyxia were
328 associated with the increasing risk of neonatal hypoglycemia. Therefore, early initiation of
329 feeding is an important factor to prevent hypoglycemia for all newborns. On the other hand,
330 preventing, early detection and management of hypoglycemia should be done in neonates with
331 hypothermia and perinatal asphyxia.

332 **8. Acknowledgment**

333 We the authors acknowledged study participants, data collectors, supervisors, Hawassa City
334 Administration Health Bureau, and public hospitals for feeding the necessary information and
335 their unwavering efforts in the research process. Moreover, we appreciate Hawassa University to
336 pursue this chance to conduct this research and Dilla University for covering the data collection
337 cost.

338

339 9. Reference

- 340 1. stanford m. Hypoglycemia in a Newborn Baby 2023 [cited 2023 march, 2]. Available
341 from: [https://www.stanfordchildrens.org/en/topic/default?id=hypoglycemia-in-the-](https://www.stanfordchildrens.org/en/topic/default?id=hypoglycemia-in-the-newborn-90-P01961)
342 [newborn-90-P01961](https://www.stanfordchildrens.org/en/topic/default?id=hypoglycemia-in-the-newborn-90-P01961).
- 343 2. Dixon KC, Ferris RL, Marikar D, Chong M, Mittal A, Manikam L, et al. Definition and
344 monitoring of neonatal hypoglycaemia: a nationwide survey of NHS England Neonatal
345 Units. Archives of Disease in Childhood-Fetal and Neonatal Edition. 2017;102(1):F92-F3.
- 346 3. Harris DL, Weston PJ, Battin MR, Harding JE. A survey of the management of neonatal
347 hypoglycaemia within the Australian and New Zealand Neonatal Network. Journal of
348 Paediatrics and Child Health. 2014;50(10):E55-E62.
- 349 4. EMOH. neonatal intensive care unit management protocol2021.
- 350 5. Stanescu A, Stoicescu S. Neonatal hypoglycemia screening in newborns from diabetic
351 mothers-Arguments and controversies. Journal of medicine and life. 2014;7(Spec Iss
352 3):51.
- 353 6. Mukunya D, Odongkara B, Piloya T, Nankabirwa V, Achora V, Batte C, et al. Prevalence
354 and factors associated with neonatal hypoglycemia in Northern Uganda: a community-
355 based cross-sectional study. Tropical Medicine and Health. 2020;48(1):1-8.
- 356 7. Hay WW, Raju TN, Higgins RD, Kalhan SC, Devaskar SU. Knowledge gaps and research
357 needs for understanding and treating neonatal hypoglycemia: workshop report from
358 Eunice Kennedy Shriver National Institute of Child Health and Human Development. The
359 Journal of pediatrics. 2009;155(5):612-7.
- 360 8. Harding JE, Harris DL, Hegarty JE, Alsweiler JM, McKinlay CJ. An emerging evidence
361 base for the management of neonatal hypoglycaemia. Early human development.
362 2017;104:51-6.
- 363 9. Shah R, Harding J, Brown J, McKinlay C. Neonatal Glycaemia and Neurodevelopmental
364 Outcomes: A Systematic Review and Meta-Analysis. Neonatology. 2019;115(2):116-26.
- 365 10. Dedeke I, Okeniyi J, Owa J, Oyedeji G. Point-of-admission neonatal hypoglycaemia in a
366 Nigerian tertiary hospital: incidence, risk factors and outcome. Nigerian Journal of
367 Paediatrics. 2011;38(2):90-4.
- 368 11. Sertsu A, Nigussie K, Eyeberu A, Tibebe A, Negash A, Getachew T, et al. Determinants
369 of neonatal hypoglycemia among neonates admitted at Hiwot Fana Comprehensive

- 370 Specialized University Hospital, Eastern Ethiopia: A retrospective cross-sectional study.
371 SAGE Open Medicine. 2022;10:20503121221141801.
- 372 12. Fantahun BY, Nurussen I. Prevalence and Risk factors of Hypoglycaemia in Neonates at
373 St. Paul's Hospital Millennium Medical College Neonatal Intensive Care Unit, Ethiopia:
374 A Cross Sectional Study. 2020.
- 375 13. Amponsah G, Hagan O, Okai E. Neonatal Hypoglycaemia at Cape Coast Teaching
376 Hospital. *Journal of the West African College of Surgeons*. 2015;5(2):100.
- 377 14. HUCSHHR hucshhr. Number of serving population and staff profile. 2023.
- 378 15. AGHHR aghhr. Number of serving population and staff profile. 2023.
- 379 16. TPHHR TpH. Number of serving population and staff profile. 2023.
- 380 17. Fikirte Kasaye RM, Kerebih Abere. Prevalence and predictors of hypoglycemia among
381 neonates on admission to NICU in Addis Ababa public Hospitals, Addis Ababa, Ethiopia,
382 2021. Addis abeba university repository 2021.
- 383 18. Meyur R, Sadhu A, Bhakta A, Bandyopadhyay M, Kundu B, Bhaumik S, et al.
384 INCIDENCE & CAUSES OF NEONATAL HYPOGLYCEMIA AFTER CESAREAN
385 SECTION IN A RURAL SETUP OF WEST BENGAL. *Journal of Evolution of medical
386 and Dental Sciences*. 2014;3:1191-4.
- 387 19. Singh YP, Devi TR, Gangte D, Devi TI, Singh NN, Singh MA. Hypoglycemia in newborn
388 in Manipur. *Journal of Medical Society*. 2014;28(2):108-11.
- 389 20. Hosagasi NH, Aydin M, Zenciroglu A, Ustun N, Beken S. Incidence of hypoglycemia in
390 newborns at risk and an audit of the 2011 American academy of pediatrics guideline for
391 hypoglycemia. *Pediatrics & Neonatology*. 2018;59(4):368-74.
- 392 21. Babu M, D'Souza J, Susheela C. Study of incidence, clinical profile and risk factors of
393 neonatal hypoglycemia in a tertiary care hospital. *Int J Pediatr Res*. 2016;3(10):753.
- 394 22. Ochoga MO, Aondoaseer M, Abah RO, Ogbu O, Ejeliogu EU, Tolough GI. Prevalence of
395 Hypoglycaemia in Newborn at Benue State University Teaching Hospital, Makurdi,
396 Benue State, Nigeria. *Open Journal of Pediatrics*. 2018;8(2):189-98.
- 397 23. Sambany E, Pussard E, Rajaonarivo C, Raobijaona H, Barennes H. Childhood
398 dysglycemia: prevalence and outcome in a referral hospital. *PloS one*. 2013;8(5):e65193.
- 399 24. Zhao T, Liu Q, Zhou M, Dai W, Xu Y, Kuang L, et al. Identifying risk effectors involved
400 in neonatal hypoglycemia occurrence. *Bioscience Reports*. 2020;40(3).

- 401 25. Magadla Y, Velaphi S, Moosa F. Incidence of hypoglycaemia in late preterm and term
402 infants born to women with diabetes mellitus. *South African Journal of Child Health*.
403 2019;13(2):78-83.
- 404 26. Hassan M, Pervez A, Biswas R, Debnath S, Syfullah K. Incidence and Risk Factors of
405 Neonatal Hypoglycemia During the First 48 Hours of Life in a Tertiary Level Hospital.
406 *Faridpur Medical College Journal*. 2020;15:12-5.
- 407 27. Bamehrez M. Hypoglycemia and associated comorbidities among newborns of mothers
408 with diabetes in an academic tertiary care center. *Frontiers in Pediatrics*. 2023;11.
- 409 28. Yunarto Y, Sarosa GI. Risk factors of neonatal hypoglycemia. *Paediatrica Indonesiana*.
410 2019;59(5):252-6.
- 411 29. Bromiker R, Perry A, Kasirer Y, Einav S, Klinger G, Levy-Khademi F. Early neonatal
412 hypoglycemia: incidence of and risk factors. A cohort study using universal point of care
413 screening. *The Journal of Maternal-Fetal & Neonatal Medicine*. 2019;32(5):786-92.
- 414 30. Zhou W, Yu J, Wu Y, Zhang H. Hypoglycemia incidence and risk factors assessment in
415 hospitalized neonates. *The Journal of Maternal-Fetal & Neonatal Medicine*.
416 2015;28(4):422-5.
- 417 31. Chandran S, Samuel Rajadurai V, Alim Abdul Haium A, Hussain K. Current perspectives
418 on neonatal hypoglycemia, its management, and cerebral injury risk. *Research and reports*
419 *in Neonatology*. 2015:17-30.
- 420 32. Senadhi V, Dutta S. Recurrent Hypothermia and Hypoglycemia as the Initial Presentation
421 of Hepatitis C: 724. *Official journal of the American College of Gastroenterology | ACG*.
422 2010;105:S262.
- 423 33. Cordero L, Stenger MR, Landon MB, Nankervis CA. Early feeding, hypoglycemia and
424 breastfeeding initiation in infants born to women with pregestational diabetes mellitus. *J*
425 *Neonatal Perinatal Med*. 2018;11(4):357-64.

426

HYPOGLYCEMIA




Figure 1. The magnitude of hypoglycemia among neonates on admission to neonatal intensive care units in Hawassa City Public Hospitals, Ethiopia 2023. (n=271)

Figure