

Supplementary material for “White matter hyperintensity spatial patterns: risk factors and clinical correlates”

Frauke Beyer^{a,b}; Ami Tsuchida^c; Aicha Soumaré^a; Hema Sekhar Reddy Rajula^a; Aniket Mishra^a; Fabrice Crivello^c; Markus Loeffler^f; Christophe Tzourio^a; Philippe Amouyel^g; Arno Villringer^{b,e}; Markus Scholz^f; Hélène Jacqmin-Gadda^a; Marc Joliot^c; A. Veronica Witte^{b,d,e}; Carole Dufouil^a; ; Stéphanie Debette^a

^a Bordeaux Population Health Research Center, University of Bordeaux, Inserm, UMR 1219, Bordeaux, France

^b Department of Neurology, Max Planck Institute for Human Cognitive and Brain Sciences, Leipzig, Germany

^c University of Bordeaux, CEA, CNRS, Institute of Neurodegenerative Diseases, UMR5293, Neurofunctional imaging group, Bordeaux, France

^d CRC 1052 “Obesity Mechanisms”, Subproject A1, University of Leipzig, Leipzig, Germany

^e Clinic of Cognitive Neurology, University Clinic Leipzig, Leipzig, Germany

^f Institute for Medical Informatics, Statistics, and Epidemiology (IMISE), University of Leipzig, Leipzig, Germany

^gINSERM U1167, University of Lille, Institut Pasteur de Lille, Lille, France

Corresponding author: Stéphanie Debette, University of Bordeaux, INSERM, Bordeaux Population Health Center, UMR1219, 146 rue Léo Saignat, 33076 Bordeaux, France, stephanie.debette@u-bordeaux.fr

1. Using LIFE-Adult as reference cohort for PCA decomposition

In order to check whether the spatial components were stable across cohorts, we also used LIFE-Adult as reference cohort to derive the spatial components with PCA.




Figure 1: WMH components derived from LIFE-Adult using 7 components and oblimin rotation




Figure 2: Spatial correlation of the components, calculated as correlation of loadings over the 36 Bullseye regions. Anatomical labeling on x-axis refers to components based on 3C-Dijon and described in the manuscript. Numerical labels (TC) on y-axis refer to components based on LIFE-Adult.

For the PCA in LIFE-Adult, the optimal number of components was 5, but we used 7 components to be able to compare to 3C-Dijon. We then calculated the correlation of component loadings between the original components derived from 3C-Dijon (Figure 2, labeled with anatomical descriptions) and the components derived from LIFE-Adult (labeled with numbers, compare Figure 2). The resulting components were very highly correlated (Pearson's $r > 0.75$) with slight changes in the order. We then calculated the correlation of the original and the projected component scores over participants for each sample and component. The results for each component are shown in Figure 3. For all components, both approaches yielded very consistent component scores ($R^2 > 0.75$). We therefore believe that it was appropriate to use 3C-Dijon as the reference in this

study, and that we could expect similar results when deriving components from LIFE-Adult.




Figure 3: Comparison of component scores derived from 3C and LIFE-Adult and their correlations across participants from both cohorts

2 Characterization of WMH spatial patterns

In the following, we aimed to characterize the WMH components derived from 3C-Dijon based on their intercorrelations, and their associations with age and sex.

WMH spatial components were correlated with one another (see Figure 4), as expected when using oblimin rotation in PCA. In general, correlations were weaker in 3C-Dijon.

For both cohorts, strongest correlations were between fronto-parietal, anterior PV and temporal WMH. Anterior PV WMH were most strongly associated with total WMH volume ($r=0.91$ in LIFE-Adult). Left and right posterior horn WMH were strongly correlated in LIFE-Adult and less so in 3C-Dijon. The correlations of juxtacortical-deep occipital and deep basal ganglia WMH with the other components and total WMH were weaker in both studies.



Figure 4: Correlation of component scores and total WMH volume in 3C-Dijon (upper triangle) and LIFE-Adult (lower triangle)

3 Age- and sex association of WMH spatial components

In 3C-Dijon, only fronto-parietal (C1) and anterior PV WMH (C3) were positively associated with age. All other components except for left posterior horn WMH (C2) showed slight negative associations (Figure 5). In LIFE-Adult, all components were significantly positively associated with age (Figure 6). Age explained least variance in juxtacortical-deep occipital WMH (C5, 1%) and most in anterior PV WMH (C3, 11%). In 3C, males had higher scores in fronto-parietal (C1), right posterior horn (C4) and deep peri basal ganglia WMH (C7) (Figure 7). Similarly, in LIFE, males had higher scores in left and right posterior horn (C2, C4) and temporal WMH (C6) (Figure 8).




Figure 5: Association of age with WMH components in 3C-Dijon. R^2 and P-values of simple regression are shown.




Figure 6: Association of age with WMH components in LIFE-Adult. R^2 and P -values of simple regression are shown.




Figure 7: Association of sex with WMH components in 3C-Dijon. P -values of two-sample t -tests are shown.




Figure 8: Association of sex with WMH components in LIFE-Adult- P-values of two-sample t-tests are shown

4. Flowcharts




Figure 9: Flowchart of sample selection for 3C-Dijon




Figure 10: Flowchart of sample selection for LIFE-Adult

5. Details on cognitive assessments in 3C-Dijon and LIFE-Adult

A description of number of participants at baseline and throughout waves in 3C-Dijon is found in Tables 1 and 2. The sample for the baseline cognitive assessment in LIFE-Adult is described in Table 3.

Table 1: Characteristics of baseline cognitive sample in 3C-Dijon

Variable	N	Mean/%	Std. Dev.
Age (y)	1668	72	4.1
Gender(women)	1668 (1011)	61%	
Education			
no formal education	85	5%	
primary education with diploma	179	11%	
secondary education short	733	44%	
secondary education long	313	19%	
higher education	358	21%	

Table 2: Characteristics of cognitive assessment across waves in 3C-Dijon. Shown are average, standard deviation, minimal and maximal values of raw test scores as well as average time since baseline per time point. N indicates number of participants who participated in each of the assessments at timepoints T0, T1, T2, T4, T5 and T6.

Variable	N	Mean	SD	Min	Max
T0 (Time since baseline in years)	1668	0	0	0	0
BENTON	1655	11.8	1.88	0	15
ISA	1659	34.2	6.73	11	60
TMTA	1651	52.6	20.1	15	300
TMTB	1621	101	41.1	11	300
T1	1583	1.85	0.172	0.751	3.66
BENTON	1577	11.7	2.02	0	15
ISA	1583	36.4	6.78	12	59
TMTA	0				
TMTB	0				
T2	1502	3.58	0.206	2.83	5.4
BENTON	1495	11.4	1.99	1	15
ISA	1500	35.5	6.66	4	66
TMTA	1482	53.1	19.9	18	210
TMTB	1462	104	43.7	18	300
T4	1241	7.04	0.379	5.35	8.61
BENTON	1237	11.6	2.21	0	15
ISA	1240	34.6	6.78	9	58
TMTA	1228	48.6	18.8	16	200
TMTB	1219	119	48.3	16	300
T5	1016	8.8	0.413	7.35	10.6
BENTON	1013	11.9	2.09	0	15
ISA	1016	35.2	6.88	13	64
TMTA	1006	48.6	20.3	20	206
TMTB	997	118	47.8	22	300

Variable	N	Mean	SD	Min	Max
T6	808	10.8	0.387	9.67	12.2
BENTON	796	12.1	1.9	0	15
ISA	805	35	6.46	13	55
TMTA	801	48.4	17.6	16	154
TMTB	792	117	47.5	43	300




Figure 11: Overview of trajectories of neuropsychological test scores in 3C-Dijon. Blue line shows local polynomial regression.

Table 3: Descriptive summary of the subsample with cognitive data in LIFE-Adult

Variable	N	Mean/%	Median
Age (y)	1039	70.5	70
Gender (women)	1039 (475)	45.7%	
Education			
no secondary school degree,	7	0.7%	
secondary school degree,	86	8.3%	
advanced secondary school degree	576	55.4%	
university entrance degree	370	35.6%	
MMST	1020	28.5	29

5. Vascular risk factor results

Table 4: Associations of VR factors with WMH spatial components in the 3C-Dijon and LIFE-Adult samples. Shown are all associations with $pFDR < 0.05$ in at least one of the cohorts. Estimate refers to standardized regression estimates from regression models adjusting for age, sex and TIV. FDR-p values are across cohorts and estimates. β , SE, and p-value from the meta-analysis. APOE-e2: dominant model, APOE-e2_homo: comparing homozygotes to non-carriers, APOE-e2_33: comparing homozygotes to 33 carriers, HTN: hypertension, SBP: systolic blood pressure, DBP: diastolic blood pressure, WHR: waist-to-hip ratio, WMH: white matter hyperintensities, FDR: false-discovery rate, SE: standard error.

		3C-Dijon			LIFE			Meta-analysis		
WMH Measure	Predictor	β	SE	pFDR	β	SE	pFDR	β	SE	p
Fronto-Parietal	HTN	0.173	0.058	0.7	0.251	0.062	8.08e-04	0.210	0.042	8.18e-07
Fronto-Parietal	SBP	0.022	0.025	0.2	0.171	0.030	6.98e-07	0.084	0.019	1.11e-05
Fronto-Parietal	DBP	0.096	0.024	0.26	0.157	0.030	6.24e-06	0.121	0.019	1.84e-10
Fronto-Parietal	BMI	0.014	0.024	0.035	0.073	0.029	0.073	0.038	0.019	0.041
Fronto-Parietal	WHR	0.007	0.024	0.91	0.085	0.029	0.031	0.039	0.019	0.037
Fronto-Parietal	WHR binary	0.036	0.056	0.011	0.183	0.088	0.15	0.079	0.047	0.096
Fronto-Parietal	Diabetes	0.068	0.077	0.011	0.131	0.073	0.23	0.101	0.053	0.057
Left Posterior	SBP	-0.003	0.026	0.2	0.125	0.030	6.25e-04	0.051	0.019	0.0095

		3C-Dijon			LIFE			Meta-analysis		
WMH Measure	Predictor	β	SE	pFDR	β	SE	pFDR	β	SE	p
Horn										
Left Posterior Horn	DBP	0.020	0.025	0.43	0.103	0.030	0.008	0.054	0.019	0.0055
Left Posterior Horn	Smoking	-0.046	0.062	0.86	0.011	0.069	0.0077	-0.021	0.046	0.65
Anterior PV	HTN	0.238	0.058	0.95	0.298	0.062	4.49e-05	0.266	0.042	3.33e-10
Anterior PV	DBP	0.149	0.024	0.74	0.154	0.030	1.13e-05	0.151	0.019	1.33e-15
Anterior PV	WHR	0.061	0.024	0.82	0.094	0.029	0.011	0.075	0.019	6.06e-05
Anterior PV	Smoking	0.057	0.061	0.76	0.067	0.069	7.20e-04	0.062	0.046	0.18
Right posterior horn	SBP	-0.038	0.025	0.41	0.110	0.030	0.0032	0.023	0.019	0.23
Right posterior horn	DBP	-0.002	0.025	0.57	0.116	0.030	0.0018	0.046	0.019	0.018
Right posterior horn	Smoking	0.046	0.062	0.38	-0.099	0.069	4.49e-05	-0.019	0.046	0.69
Juxtacortical-deep occipital	Diabetes	0.135	0.078	0.7	-0.059	0.077	8.08e-04	0.037	0.055	0.5
Juxtacortical-deep occipital	Smoking	0.078	0.062	0.2	-0.033	0.072	6.98e-07	0.031	0.047	0.51

		3C-Dijon			LIFE			Meta-analysis		
WMH Measure	Predictor	β	SE	pFDR	β	SE	pFDR	β	SE	p
Juxtacortical-deep occipital	APOE-e2	0.056	0.070	0.035	-0.263	0.092	0.073	-0.061	0.056	0.28
Juxtacortical-deep occipital	APOE-e2_homo	0.201	0.335	0.011	-1.163	0.358	0.15	-0.436	0.245	0.075
Juxtacortical-deep occipital	APOE-e2_33	0.208	0.334	0.011	-1.142	0.353	0.23	-0.429	0.242	0.076
Temporal	HTN	0.120	0.059	0.86	0.214	0.063	0.0077	0.164	0.043	1.42e-04
Temporal	BMI	0.045	0.025	0.9	0.110	0.029	0.0021	0.072	0.019	1.37e-04
Temporal	Smoking	-0.057	0.062	0.2	0.051	0.069	6.25e-04	-0.008	0.046	0.86
Deep peri basal ganglia	HTN	0.103	0.059	0.76	0.263	0.064	7.20e-04	0.176	0.043	5.35e-05
Deep peri basal ganglia	Diabetes	0.170	0.078	0.95	0.007	0.076	4.49e-05	0.087	0.054	0.11
Deep peri basal ganglia	Smoking	0.064	0.061	0.82	0.004	0.070	3.54e-07	0.038	0.046	0.41
WMH volume	HTN	0.241	0.057	0.38	0.292	0.061	4.49e-05	0.265	0.042	1.96e-10
WMH volume	SBP	0.071	0.025	0.87	0.183	0.029	1.21e-07	0.117	0.019	4.89e-10
WMH volume	DBP	0.133	0.024	0.51	0.161	0.030	2.34e-06	0.144	0.019	9.48e-15
WMH volume	WHR	0.035	0.024	0.7	0.094	0.029	0.01	0.059	0.018	0.0012
WMH volume	Smoking	0.048	0.060	0.41	0.009	0.067	0.0032	0.031	0.045	0.49

6. APOE results

Table 5: Associations of APOE genotypes (APOE-e4/APOE-e2: dominant model including e24 alleles and APOE-e4_homo/APOE-e2_homo: model comparing homozygotes to non-carriers) with WMH spatial components in the 3C and LIFE cohorts. Shown are regression estimates from regression models adjusting for age, sex and TIV, FDR-corrected p-values for both cohorts and estimate, p-value and FDR-p from the meta-analysis. APOE: apolipoprotein E, WMH: white matter hyperintensities, FDR: false-discovery rate, SE: standard error.

WMH Measure	Model	3C				LIFE				Meta-analysis			
		β	SE	p	pFDR	β	SE	p	pFDR	β	SE	p	pFDR
Fronto-Parietal	APOE-e4	-0.03	0.06	0.65	0.81	-0.02	0.07	0.79	0.89	-0.02	0.04	0.61	0.78
Left Posterior Horn	APOE-e4	-0.05	0.06	0.44	0.73	-0.06	0.07	0.40	0.61	-0.05	0.05	0.25	0.48
Anterior PV	APOE-e4	-0.05	0.06	0.35	0.63	-0.10	0.07	0.15	0.31	-0.07	0.04	0.10	0.25
Right posterior horn	APOE-e4	0.02	0.06	0.73	0.85	0.02	0.07	0.76	0.88	0.02	0.05	0.65	0.80
Juxtacortical-deep occipital	APOE-e4	-0.03	0.06	0.63	0.81	0.12	0.07	0.10	0.22	0.03	0.05	0.52	0.70
Temporal	APOE-e4	0.03	0.06	0.66	0.81	0.09	0.07	0.23	0.40	0.05	0.05	0.27	0.48
Deep peri basal ganglia	APOE-e4	-0.13	0.06	0.02	0.14	-0.05	0.07	0.52	0.73	-0.10	0.05	0.03	0.09
WMH volume	APOE-e4	-0.04	0.06	0.51	0.76	-0.07	0.07	0.33	0.52	-0.05	0.04	0.26	0.48
Fronto-Parietal	APOE-e2	0.14	0.07	0.04	0.23	-0.20	0.09	0.02	0.08	0.01	0.05	0.88	0.92
Left Posterior Horn	APOE-e2	-0.03	0.07	0.71	0.84	-0.12	0.09	0.19	0.36	-0.06	0.06	0.27	0.48
Anterior PV	APOE-e2	0.08	0.07	0.25	0.49	-0.12	0.09	0.19	0.36	0.01	0.05	0.91	0.94
Right posterior horn	APOE-e2	0.05	0.07	0.52	0.76	-0.09	0.09	0.29	0.48	-0.01	0.06	0.88	0.92
Juxtacortical-deep occipital	APOE-e2	0.06	0.07	0.43	0.72	-0.26	0.09	0.00	0.02	-0.06	0.06	0.28	0.48
Temporal	APOE-e2	0.09	0.07	0.18	0.39	-0.02	0.09	0.87	0.90	0.05	0.06	0.34	0.56
Deep peri basal ganglia	APOE-e2	0.04	0.07	0.59	0.78	-0.02	0.09	0.85	0.90	0.02	0.06	0.76	0.87
WMH volume	APOE-e2	0.13	0.07	0.06	0.26	-0.15	0.09	0.09	0.21	0.02	0.05	0.65	0.80
Fronto-Parietal	APOE-e4_homo	0.49	0.21	0.02	0.14	0.27	0.24	0.26	0.45	0.40	0.16	0.01	0.05
Left Posterior Horn	APOE-e4_homo	0.13	0.22	0.54	0.76	-0.05	0.24	0.83	0.89	0.05	0.16	0.76	0.87
Anterior PV	APOE-e4_homo	0.09	0.21	0.68	0.82	-0.13	0.24	0.59	0.78	-0.01	0.16	0.96	0.97

		3C				LIFE				Meta-analysis			
WMH Measure	Model	β	SE	p	pFDR	β	SE	p	pFDR	β	SE	p	pFDR
Right posterior horn	APOE-e4_homo	0.08	0.22	0.72	0.85	0.47	0.24	0.05	0.15	0.25	0.16	0.12	0.27
Juxtacortical-deep occipital	APOE-e4_homo	0.33	0.22	0.13	0.35	0.05	0.25	0.84	0.90	0.21	0.16	0.20	0.41
Temporal	APOE-e4_homo	0.31	0.22	0.16	0.37	0.34	0.25	0.17	0.33	0.32	0.16	0.05	0.13
Deep peri basal ganglia	APOE-e4_homo	-0.09	0.22	0.69	0.82	-0.11	0.26	0.68	0.83	-0.10	0.16	0.56	0.75
WMH volume	APOE-e4_homo	0.49	0.21	0.02	0.14	0.11	0.24	0.65	0.81	0.32	0.16	0.04	0.12
Fronto-Parietal	APOE-e2_homo	0.23	0.33	0.49	0.76	-0.26	0.34	0.44	0.66	-0.01	0.24	0.97	0.97
Left Posterior Horn	APOE-e2_homo	-0.49	0.33	0.14	0.36	0.05	0.34	0.88	0.91	-0.22	0.24	0.35	0.56
Anterior PV	APOE-e2_homo	-0.08	0.33	0.80	0.88	-0.35	0.35	0.32	0.52	-0.21	0.24	0.38	0.58
Right posterior horn	APOE-e2_homo	-0.01	0.33	0.97	0.98	-0.26	0.34	0.45	0.66	-0.13	0.24	0.58	0.77
Juxtacortical-deep occipital	APOE-e2_homo	0.20	0.34	0.55	0.76	-1.16	0.36	0.00	0.01	-0.44	0.24	0.07	0.19
Temporal	APOE-e2_homo	0.17	0.33	0.61	0.80	-0.11	0.35	0.74	0.87	0.03	0.24	0.88	0.92
Deep peri basal ganglia	APOE-e2_homo	0.06	0.33	0.85	0.91	-0.17	0.36	0.64	0.81	-0.04	0.25	0.86	0.92
WMH volume	APOE-e2_homo	0.20	0.32	0.53	0.76	-0.38	0.34	0.26	0.45	-0.07	0.23	0.75	0.87

7 Executive function results

Table 6: Associations of WMH spatial components with baseline and change in executive function in 3C-Dijon and LIFE. Regression estimates are from the latent class mixed-effect models with WMH components and total WMH volume as predictors for intercept and change of general cognitive function, adjusting for age, sex and education. FDR-corrected p-values were calculated separately per cohort. Also shown are estimates and p-values from the meta-analysis. WMH: white matter hyperintensities, FDR: false-discovery rate, SE: standard error.

WMH Measure	Model	3C				LIFE				Meta-analysis			
		β	SE	p	pFDR	β	SE	p	pFDR	β	SE	p	pFDR
Fronto-Parietal	Intercept	-0.105	0.033	0.00176	0.111	-0.163	0.045	3.00e-04	0.0255	-0.125	0.027	3.02e-06	2.42e-05
Fronto-Parietal	Change	-0.012	0.044	0.789	1.000	-0.070	0.080	0.38	1	-0.026	0.039	0.509	0.627
Left Posterior Horn	Intercept	0.018	0.032	0.583	1.000	-0.090	0.044	0.0414	1	-0.020	0.026	0.442	0.627
Left Posterior Horn	Change	0.004	0.044	0.931	1.000	-0.164	0.076	0.03	1	-0.039	0.038	0.31	0.495
Anterior PV	Intercept	-0.100	0.034	0.00295	0.180	-0.121	0.045	0.00675	0.479	-0.108	0.027	6.19e-05	3.30e-04
Anterior PV	Change	-0.119	0.042	0.00506	0.301	-0.076	0.078	0.33	1	-0.109	0.037	0.00342	0.0109
Right posterior horn	Intercept	0.032	0.033	0.332	1.000	-0.050	0.044	0.251	1	0.002	0.026	0.933	0.933

		3C				LIFE				Meta-analysis			
WMH Measure	Model	β	SE	p	pFDR	β	SE	p	pFDR	β	SE	p	pFDR
Right posterior horn	Change	0.013	0.041	0.751	1.000	-0.149	0.075	0.0482	1	-0.025	0.036	0.499	0.627
Juxtacortical-deep occipital	Intercept	0.039	0.033	0.239	1.000	-0.070	0.042	0.0963	1	-0.003	0.026	0.917	0.933
Juxtacortical-deep occipital	Change	0.005	0.043	0.901	1.000	-0.090	0.077	0.24	1	-0.018	0.038	0.64	0.731
Temporal	Intercept	-0.017	0.033	0.608	1.000	-0.179	0.045	6.00e-05	0.00522	-0.075	0.027	0.00521	0.0139
Temporal	Change	0.059	0.043	0.169	1.000	0.013	0.081	0.874	1	0.049	0.038	0.197	0.351
Deep peri basal ganglia	Intercept	-0.067	0.033	0.0422	1.000	-0.149	0.043	5.40e-04	0.0443	-0.098	0.026	2.01e-04	8.04e-04
Deep peri basal ganglia	Change	0.080	0.042	0.0575	1.000	0.063	0.085	0.455	1	0.077	0.038	0.0419	0.0959
WMH volume	Intercept	-0.124	0.034	2.60e-04	0.017	-0.154	0.045	7.30e-04	0.0584	-0.135	0.027	7.41e-07	1.19e-05
WMH volume	Change	-0.051	0.044	0.243	1.000	-0.120	0.079	0.129	1	-0.068	0.038	0.0788	0.158

8 Memory function

Table 7: Associations of WMH spatial components with baseline and change in memory function in 3C-Dijon. Regression estimates are from the latent class mixed-effect models with WMH components and total WMH volume as predictors for intercept and (linear and quadratic) change of visual memory function, adjusting for age, sex and education. WMH: white matter hyperintensities, FDR: false-discovery rate, SE: standard error.

WMH Measure	Effect	β	SE	p	pFDR
Fronto-Parietal	Intercept	-0.032	0.028	0.259	1.000
Fronto-Parietal	Linear	0.057	0.118	0.629	1.000
Fronto-Parietal	Quadratic	-0.708	1.122	0.528	1.000
Left posterior horn	Intercept	0.024	0.034	0.477	1.000
Left posterior horn	Linear	-0.005	0.196	0.981	1.000
Left posterior horn	Quadratic	0.182	1.826	0.921	1.000
Anterior PV	Intercept	-0.054	0.028	0.054	1.000
Anterior PV	Linear	-0.076	0.114	0.505	1.000
Anterior PV	Quadratic	-0.262	1.070	0.806	1.000
Right posterior horn	Intercept	0.013	0.028	0.640	1.000
Right posterior horn	Linear	-0.011	0.128	0.933	1.000
Right posterior horn	Quadratic	-0.182	1.207	0.880	1.000
Juxtacortical-deep occipital	Intercept	-0.023	0.028	0.412	1.000
Juxtacortical-deep occipital	Linear	0.097	0.117	0.406	1.000
Juxtacortical-deep occipital	Quadratic	-0.904	1.107	0.414	1.000
Temporal	Intercept	0.026	0.027	0.321	1.000
Temporal	Linear	-0.073	0.097	0.449	1.000

WMH Measure	Effect	β	SE	p	pFDR
Temporal	Quadratic	-0.042	0.892	0.962	1.000
Deep peri basal ganglia	Intercept	-0.085	0.028	0.002	0.132
Deep peri basal ganglia	Linear	0.296	0.116	0.011	0.624
Deep peri basal ganglia	Quadratic	-2.479	1.083	0.022	1.000
WMH volume	Intercept	-0.072	0.038	0.057	1.000
WMH volume	Linear	0.026	0.154	0.867	1.000
WMH volume	Quadratic	-1.303	1.457	0.371	1.000

Table 8: Associations of WMH spatial components with baseline and change in verbal memory function in LIFE. Standardized regression estimates are from the latent class mixed-effect models with WMH components and total WMH volume as predictors for intercept and change of memory, adjusting for age, sex and education. FDR-corrected p-values are shown. WMH: white matter hyperintensities, FDR: false-discovery rate, SE: standard error.

WMH Measure	Model	β	SE	p	pFDR
Fronto-Parietal	Intercept	-0.156	0.038	5.00e-05	0.004
Fronto-Parietal	Change	-0.075	0.054	0.167	1.000
Left posterior horn	Intercept	-0.116	0.038	0.00194	0.147
Left posterior horn	Change	-0.067	0.051	0.194	1.000
Anterior PV	Intercept	-0.112	0.038	0.00293	0.220
Anterior PV	Change	-0.069	0.052	0.182	1.000
Right posterior horn	Intercept	-0.136	0.038	3.00e-04	0.025
Right posterior horn	Change	-0.068	0.051	0.186	1.000
Juxtacortical-deep occipital	Intercept	-0.025	0.036	0.494	1.000

WMH Measure	Model	β	SE	p	pFDR
Juxtacortical-deep occipital	Change	-0.081	0.051	0.113	1.000
Temporal	Intercept	-0.103	0.038	0.00647	0.466
Temporal	Change	-0.111	0.053	0.0378	1.000
Deep peri basal ganglia	Intercept	-0.097	0.036	0.00753	0.527
Deep peri basal ganglia	Change	0.015	0.058	0.793	1.000
WMH volume	Intercept	-0.150	0.039	1.10e-04	0.009
WMH volume	Change	-0.100	0.054	0.0635	1.000

9 Dementia and stroke results

Table 9: Associations of WMH spatial components with all-cause dementia incidence in 3C-Dijon from Cox proportional hazard models adjusting for age, sex, education and TIV.

WMH measure	HR	2.5% CI	97.5% CI	p	pFDR
Fronto-Parietal	1.11	0.94	1.33	0.224	0.90
Left posterior horn	0.86	0.72	1.02	0.088	0.44
Anterior PV	1.27	1.05	1.53	0.012	0.08
Right posterior horn	1.10	0.92	1.33	0.293	0.90
Juxtacortical-deep occipital	1.22	1.00	1.47	0.045	0.27
Temporal	0.97	0.81	1.16	0.716	1.00
Deep peri basal ganglia	0.99	0.83	1.18	0.910	1.00
WMH volume	1.37	1.08	1.74	0.010	0.08

Table 10: Associations of WMH spatial components with probable or possible AD dementia incidence in 3C-Dijon from Cox proportional hazard models adjusting for age, sex, education and TIV.

WMH Measure	HR	2.5% CI	97.5% CI	P	pFDR
Fronto-Parietal	1.11	0.91	1.35	0.32	0.78
Left posterior horn	0.87	0.70	1.07	0.18	0.78
Anterior PV	1.20	0.97	1.49	0.10	0.78
Right posterior horn	1.10	0.88	1.36	0.40	0.78
Juxtacortical-deep occipital	1.21	0.97	1.51	0.10	0.78
Temporal	0.86	0.70	1.05	0.14	0.78
Deep peri basal ganglia	0.89	0.72	1.09	0.25	0.78
WMH volume	1.25	0.95	1.65	0.11	0.78

Table 11: Associations of WMH spatial components with any stroke incidence in 3C-Dijon (N=60) from Cox proportional hazard models adjusting for age, sex and TIV.

WMH Measure	HR	2.5% CI	97.5% CI	P	pFDR
Fronto-Parietal	1.39	1.09	1.78	0.0073	0.12
Left posterior horn	1.03	0.80	1.32	0.81	1
Anterior PV	1.67	1.28	2.18	1.50e-04	0.0036
Right posterior horn	1.20	0.93	1.54	0.17	1
Juxtacortical-deep occipital	1.32	1.02	1.72	0.038	0.56
Temporal	1.22	0.95	1.56	0.11	1
Deep peri basal ganglia	1.16	0.91	1.46	0.23	1
WMH volume	1.80	1.32	2.45	2.10e-04	0.0048

Table 12: Associations of WMH spatial components with ischemic stroke incidence in 3C-Dijon (N=50) from Cox proportional hazard models adjusting for age, sex, education and TIV.

WMH Measure	HR	2.5% CI	97.5% CI	P	pFDR
Fronto-Parietal	1.25	0.94	1.64	0.12	1
Left posterior horn	1.03	0.77	1.36	0.86	1
Anterior PV	1.54	1.14	2.08	0.004	0.076
Right posterior horn	1.26	0.94	1.68	0.12	1
Juxtacortical-deep occipital	1.07	0.82	1.41	0.62	1
Temporal	1.21	0.92	1.61	0.18	1
Deep peri basal ganglia	0.96	0.73	1.26	0.76	1
WMH volume	1.36	1.04	1.78	0.023	0.36

Table 13: Associations of WMH spatial components with hemorrhagic stroke incidence in 3C-Dijon (N=10) from Cox proportional hazard models adjusting for age, sex, education and TIV.

WMH Measure	HR	2.5% CI	97.5% CI	p	pFDR
Fronto-Parietal	1.88	1.02	3.45	0.044	0.62
Left posterior horn	1.06	0.56	2.00	0.86	1
Anterior PV	3.39	1.58	7.29	0.002	0.04
Right posterior horn	1.32	0.68	2.57	0.41	1
Juxtacortical-deep occipital	3.69	1.51	9.05	0.004	0.076
Temporal	1.42	0.77	2.62	0.27	1
Deep peri basal ganglia	2.39	1.43	3.99	0.001	0.021
WMH volume	2.51	1.46	4.32	8.91e-04	0.02