

Radiobiological meta-analysis of the response of prostate cancer to high dose rate brachytherapy

Eva G Kölmel^{1,2}, Miguel Pombar², Juan Pardo-Montero^{1,2,*}

1. Group of Medical Physics and Biomathematics, Instituto de Investigación Sanitaria de Santiago (IDIS), Santiago de Compostela, Spain.

2. Department of Medical Physics, Complejo Hospitalario Universitario de Santiago de Compostela, Spain.

***Correspondence:** Juan Pardo-Montero, Grupo de Física Médica e Biomatemáticas, Instituto de Investigación Sanitaria de Santiago (IDIS), Servizo de Radiofísica e Protección Radiolóxica, Hospital Clínico Universitario de Santiago, Trav. Choupana s/n, 15706, Santiago de Compostela (Spain); E-mail: juan.pardo.montero@sergas.es; Phone: +34 981955604

Abstract

Background and purpose: Clinical studies have shown a marked reduction in tumor control in prostate cancer treated with radically hypofractionated high-dose-rate brachytherapy (HDR-BT). The purpose of this study was to analyze the dose-response of prostate cancer treated with HDR-BT, specifically aiming at investigating the potential failure of the linear-quadratic (LQ) model to describe the response at large doses-per-fraction.

Materials and methods: We collated a dataset of dose-response to HDR-BT (3258 patients). The analysis was conducted separately for low and intermediate risk, resulting in 21 schedules (1643 patients) and 22 schedules (1615 patients), respectively. Data were fitted to tumor control probability models based on the LQ model, the linear-quadratic-linear (LQL), and a modification of the LQ model to include the effect of reoxygenation during treatment.

Results: The LQ cannot fit the data unless the α/β is allowed to be very high ($\sim[20-100]$ Gy, 95% confidence interval). If the α/β is constrained to be low (< 8 Gy) the LQ model cannot reproduce the clinical results, and the LQL model, which includes a moderation of radiation damage with increasing dose, significantly improves the fitting. On the other hand, the reoxygenation model does not match the results obtained with the LQL.

Conclusion: The clinically observed reduction in tumor control in prostate cancer treated with radical HDR-BT is better described by the LQL model. Using the best-fitting parameters, the BED for a $20 \text{ Gy} \times 1$ treatment (95 Gy) is far less than that of a conventional $2 \text{ Gy} \times 37$ fractionation (184 Gy). These results may assist in the design of radical HDR-BT treatment.

NOTE: This preprint reports new research that has not been certified by peer review and should not be used to guide clinical practice.

Introduction

Radiotherapy is widely used to treat prostate cancer. Radiotherapy options include external beam radiotherapy [1], proton therapy [2, 3], and brachytherapy, either low-dose-rate or high-dose-rate (HDR-BT) [4, 5]. HDR-BT is nowadays used as monotherapy for many patients [6], achieving good clinical outcomes, and because it is delivered with hypofractionated schedules, reducing treatment time and increasing patient comfort [5].

The response of prostate cancer to radiotherapy has been extensively studied [7, 8, 9]. The consensus is that the α/β ratio of prostate cancer is low (typically in the 1-4 Gy range), and therefore this tumor is very sensitive to fractionation. Nonetheless, alternatives to the low α/β have been suggested, like tumor hypoxia [10]. In recent years, Stereotactic Body RadioTherapy (SBRT) has become widely used to treat prostate cancer [11], with doses per fraction reaching up to 10 Gy. The response of prostate cancer to hypofractionated SBRT protocols has been recently analyzed [12, 13, 14]. All three studies reported low α/β ratios, in agreement with values obtained from lower doses per fraction. HDR-BT is delivered with hypofractionated protocols that are even more radical, reaching >20 Gy in a single fraction. However, several HDR-BT clinical trials have shown a marked reduction in tumor control (<70%) when delivering single fraction treatments with >20 Gy. This loss in tumor control is not supported by a low α/β ratio. Guirado *et al.* have recently analyzed the response of prostate cancer to HDR-BT, suggesting a large α/β ratio (~23 Gy) to explain the poor control achieved with HDR-BT single-fraction treatments [15]. They also argued that the linear-quadratic (LQ) model may not be adequate to describe the response to very large doses per fraction.

The validity of the LQ model for large doses per fraction has long been questioned [16], with different studies suggesting either a moderation or a boost of the cell killing effect predicted by the LQ with increasing dose per fraction [17, 18]. The moderation of the cell killing effect predicted by the LQ with increasing doses might explain the clinical results obtained with HDR-BT single-fraction treatments. This effect can be modeled with the linear-quadratic-linear (LQL) [19]. In fact, some evidence of a LQL-like response in the dose-response curves of prostate cancer treated with external radiotherapy was recently discussed in Refs. [13, 14].

The poor control obtained with HDR-BT single-fraction treatments could also be rationally explained because of hypoxia and reoxygenation, as originally suggested by Nahum *et al.* [10]. If tumors are hypoxic and reoxygenate during treatment, short protocols delivering larger doses per

fraction may be suboptimal.

In this work we have collated and analyzed a dataset of dose-response for HDR-BT of prostate cancer. We have performed a radiobiological analysis of the dose-response, considering not only the LQ model but also more advanced models including damage saturation at large doses and re-oxygenation, aiming at advancing the understanding of the response of prostate cancer to very large doses per fraction.

Materials and methods

Clinical dataset

The clinical dataset was created following a two-step process: initially, we expanded upon previously compiled datasets reported in [5, 20, 6]; subsequently, we performed a systematic search in Pubmed (in July 2023) for articles published after 2018 (the publication year of [5]). From each study, we extracted the number of patients, the distribution of patients according to the risk level, the number/percentage of patients receiving androgen deprivation therapy (ADT), the dose per fraction, the total dose, the overall treatment time and schedule details (fractions per day, time intervals between fractions), and the 5 year control rate, with control defined as freedom from clinical or biochemical failure (PSA nadir + 2 ng/mL). Studies that did not report any of these variables were excluded. Some studies included slightly different fractionations, and in those cases, the most commonly used fractionation was included. If different studies reported on the same (or similar) cohort with different follow-up, only the most recent publication was considered.

Overall, the collated dataset contained data from 19 studies (3258 patients). Our analysis was conducted separately for low risk (LR) and intermediate risk (IR), resulting in 21 schedules (1643 patients) for LR and 22 schedules (1615 patients) for IR. A limited number of data were found for high risk, but they were ignored because the number of schedules was not large enough to conduct the analysis. Some studies stratified patients in more than three groups (e.g. “favorable intermediate risk”, “unfavorable intermediate risk” and “very low risk”). In such cases, these results were merged into a single group. The percentage of patients receiving ADT was included in the dataset, even though this variable was not used in this analysis.

An overview of the schedules included in the analysis is presented in Table 1, with further detailed

information is presented in Supplementary Table 1.

Radiobiological models

We used three different models to investigate the dose-response of prostate cancer to HDR-BT:

1. The LQ model: the surviving fraction of tumor cells after a dose d is [40],

$$-\log SF_{LQ} = \alpha d + \beta d^2 \quad (1)$$

with α and β being the linear and quadratic parameters of the LQ model.

2. The LQL model: this model [19] includes a moderation of the quadratic term of the LQ model with increasing dose, which is controlled by the parameter δ :

$$-\log SF_{LQL} = \alpha d + \frac{2\beta}{\delta^2} (\delta d + e^{-\delta d} - 1) \quad (2)$$

3. Stavrev's model of reoxygenation: this model [41] relies on the LQ model, but with time-dependent α and β parameters to account for reoxygenation during treatment:

$$\alpha(t) = \alpha_0 \exp\left(\frac{-bt^2}{2}\right) + \alpha_0(1+\phi) \left(1 - \exp\left(\frac{-bt^2}{2}\right)\right) \quad (3)$$

$$\beta(t) = \beta_0 \left(\frac{\alpha(t)}{\alpha_0}\right)^2 \quad (4)$$

with parameters b and ϕ controlling the evolution of $\alpha(t)$ and $\beta(t)$. This results in a time dependent α/β ratio, which due to the quadratic dependence of β increases with time.

Incomplete repair

Several schedules in the dataset delivered multiple fractions per day. In this situation, incomplete repair between consecutive fractions may play a role in the response to treatment. Therefore, we also investigated the possible contribution of incomplete repair by including it in the modeling. We used the LQ with incomplete repair correction [42, 40]. The surviving fraction of cells following the

i-th radiation fraction is given by:

$$\log SF_i = -\alpha d_i - \beta d_i^2 - 2\beta d_i \sum_{p=1}^{i-1} d_p \prod_{q=p}^{i-1} \theta_q \quad (5)$$

$$\theta_q = \exp(-\nu \Delta t_q); \Delta t_q = t_{q+1} - t_q \quad (6)$$

where ν is the repair rate of sub-lethal damage (we will refer instead to the half-life of damage, defined as $T_{repair} = \log 2/\nu$).

Incomplete repair was also considered in the LQL and reoxygenation models, using an expression identical to Eq. (5), but with $\alpha(t_i)$ and $\beta(t_i)$ replacing α and β for the reoxygenation model (where t_i is the delivery time of the i-th fraction), and with an effective β -term replacing β for the LQL model, which can be obtained from Eq. (2):

$$\beta_{eff} = \frac{2\beta}{(d\delta)^2} (\delta d + e^{-\delta d} - 1) \quad (7)$$

Overall surviving fraction and proliferation

When delivering n fractions in an overall treatment time T , the surviving fraction is given by:

$$SF_{treatment} = \left(\prod_{i=1}^n SF_i \right) \exp(\lambda \max(0, T - T_k)) \quad (8)$$

where SF_i is the surviving fraction associated to each fraction, and proliferation is modeled as exponential with rate λ after a kick-off time T_k .

Tumor control probability and EQD2

The tumor control probability (TCP) was modeled using a logistic function [43],

$$TCP = \frac{1}{1 + \left(\frac{D_{50}}{EQD2} \right)^{4y}} \quad (9)$$

where D_{50} is the dose corresponding to 50% control (in 2 Gy fractions), and γ controls the slope of the dose-response curve. The *equivalent dose in 2 Gy fractions*, $EQD2$, of each schedule is model-dependent, and its explicit calculation for the models used in this work is shown in the Supplementary Materials.

Statistical methods

We fitted the models to the clinical data with the maximum likelihood methodology, assuming binomial statistics for the reported control. We used an in-house developed algorithm [14] based on the simulated annealing method to perform the optimization (minimization of $-\log L$, where L is the likelihood). Confidence intervals (CI) for the best-fitting parameters were obtained using the profile likelihood method [44].

Fits with the LQ model have five free parameters (α/β , $\lambda'=\lambda/\alpha$, T_k , γ , D_{50}). For the LQL model there is an extra free parameter, δ , and for Stavrev's reoxygenation model there are two extra parameters, φ and b . When including incomplete repair there is an extra parameter for each model, T_{repair} .

The space of parameter values was constrained to avoid reaching solutions that could be unphysical or not supported by biological data, and to speed up convergence. In particular, dose compensation due to accelerated proliferation was limited to $\lambda' \leq 2 \text{ Gy day}^{-1}$, a limit well higher than the proliferation found in [14], and the half-life of sublethal repair was limited to $T_{repair} \leq 6 \text{ h}$. For the α/β ratio, we employed two different constraints due to the discrepancies on the reported values from external radiotherapy and HDR-BT: on the one hand, a constraint $1 \leq \alpha/\beta \leq 100 \text{ Gy}$ to allow for large α/β ratios like those reported in [15]; on the other hand, a stronger constraint $1 \leq \alpha/\beta \leq 8 \text{ Gy}$ to force a low α/β ratio consistent with many reports from external radiotherapy.

The Akaike Information Criterion with sample size correction (AIC_c) was used to evaluate the performance of the models [45]. The ΔAIC_c of a given *model* compared to the *reference* model (the LQ model in this work) is:

$$\Delta AIC_c^{model} = AIC_c^{ref} - AIC_c^{model} \quad (10)$$

Models with lower AIC_c are preferred, *i.e.* positive ΔAIC_c .

The implementation of the methodology was performed in Matlab (Mathworks, Natick, MA).

Results

In Table 2, we present the best-fitting parameters and the goodness-of-fit ($-\log L$ and AIC_c) obtained from fitting the LQ, LQL, and Stavrev's reoxygenation models (with and without incomplete repair correction) to low and intermediate risk data. For these fits, the α/β ratio was allowed to lie in a large interval ($1 \leq \alpha/\beta \leq 100$ Gy). In Figure 1 we present the TCP versus EQD2 curves (experimental data and model best-fits) obtained from this fitting strategy for the LQ, LQL, and reoxygenation models without incomplete repair correction. 95% confidence intervals for the α/β and δ (LQL) are presented in Supplementary Table 2.

In Table 3, we present the best-fitting parameters and the goodness-of-fit for the same models fitted to the same data, but forcing the α/β ratio to be low ($1 \leq \alpha/\beta \leq 8$ Gy), as many radiobiological studies in external radiotherapy support a low α/β ratio. In Table 4 we report the 95% confidence intervals of the best-fitting parameters for the latter scenario. The analysis of the confidence intervals, being computationally demanding, was limited to the LQ model with and without incomplete repair correction and the LQL model. In Figure 2 we present TCP versus EQD2 curves (experimental data and model best-fits) for the LQ, LQL, and reoxygenation models without incomplete repair correction.

Discussion

The response of prostate cancer to radiotherapy has been extensively studied, and the consensus is that the α/β ratio of prostate cancer is low [7-9, 13, 14]. This makes this tumor very sensitive to fractionation. Many external radiotherapy hypofractionated protocols have been investigated for prostate cancer [11]. Several studies have suggested that the LQ model may fail to describe tumor response at large doses per fraction [16-18] (even though the why and the how are not entirely clear, with different studies suggesting that the LQ may underestimate/overestimate the damage at large doses per fraction). In fact, recent analyses of the response of prostate cancer to SBRT reported a slight moderation of the LQ-predicted response at large doses [13, 14].

The possible moderation of the damage with increasing dose is soft at the doses employed for SBRT, and clinical trials of SBRT for prostate cancer still reported high tumor control with doses per fraction up to 10 Gy [14]. This is not the case in HDR-BT, which is delivered with protocols that are more radical than those of SBRT, reaching >20 Gy in a single fraction. This makes HDR-BT an ideal scenario to investigate dose-response at large doses per fraction and the potential failure of the LQ model at such doses. Several HDR-BT clinical trials have shown a marked reduction in tumor control when delivering ~20 Gy in a single fraction [21–25], including very recent studies [46] (not analyzed in this work). This important loss in tumor control is not supported by a low α/β ratio and an LQ behaviour at large doses: for example, assuming $\alpha/\beta=3$ Gy and using the LQ model, a 20 Gy \times 1 treatment would be roughly isoeffective to 8 Gy \times 5, and more effective than a conventional 2 Gy \times 37 (BEDs of 153.3, 146.7, and 123.3 Gy₃, respectively, ignoring proliferation). This led Guirado *et al.* to suggest a large α/β ratio in a recent analysis of tumor response to HDR-BT [15]. However, such a large α/β ratio is not consistent with many radiobiological studies that found a low α/β ratio for prostate cancer. It may be that the LQ model is indeed not adequate to describe the response to very large doses per fraction [16].

In this study, we investigated the dose–response of prostate cancer to HDR-BT from a dataset containing 21 schedules (1643 patients) for LR and 22 schedules (1615 patients) for IR, with doses per fraction ranging from 6 to 20.5 Gy per fraction (Table 1, Supplementary Table 1). Our analysis specifically focused on investigating the LQ and alternative models to characterize dose-response at such large doses. Because the clinical data point to an overestimation of cell killing effect by the LQ at large doses, we investigated the LQL model [19], which includes a moderation of cell killing at large doses. This particular response at large doses could also be caused by the role of hypoxia/reoxygenation, as originally suggested by Nahum *et al.* [10]. If re-oxygenation plays an important role in the response to fractionated radiotherapy, extremely hypofractionated protocols might lose tumor control. To investigate the effect of reoxygenation, we have used the simple model proposed by Stavrev *et al.* [41], which accounts for reoxygenation through time-dependent α and β parameters. Because several schedules in the dataset delivered multiple fractions per day, we also investigated the role of incomplete repair on the modeling of response to treatment in each of the three models under investigation.

We followed two strategies for data fitting. First, we imposed broad constraints on the values of the best-fitting parameters. When following this strategy, the LQ proved superior to both the LQL and reoxygenation models to describe dose-response, with $\Delta AIC_c^{LQL}=-2$ and $\Delta AIC_c^{Sta}=-8$ (Table 2, Figure 1). However, in order to fit the data, the LQ required a large α/β value (95% confidence

intervals 28-100 Gy and 17-100 Gy for LR and IR, respectively; Supplementary Table 2). This agrees with [15], who found a large α/β ratio (~ 23 Gy) from the analysis of a smaller dataset. The inclusion of incomplete repair in the modeling of response to treatment does not improve data fitting. For prostate cancer, studies have shown sublethal repair rates characterized by half-lives $T_{repair} \sim 1.5-2$ h [47]. The effect of such a repair rate would be small for times between fractions ≥ 6 h, the typical time between fractions in the studies analyzed in this work.

The large α/β ratio obtained from this fitting strategy is not in agreement with many studies analyzing dose-response of prostate cancer treated with external radiotherapy. Therefore, we investigated a second fitting strategy where the value α/β ratio was constrained to be low, $1 \leq \alpha/\beta \leq 8$ Gy. These limits were qualitatively set to double the 95% confidence intervals reported in Ref. [14]. When forcing the α/β to be low, the results were quite different (Table 3, Figure 2), and the LQL model became superior to both the LQ and reoxygenation models, $\Delta AIC_c^{LQL} \sim 22$ for LR and 19 for IR. Analyses based on the AIC typically demand $\Delta AIC_c > 10$ to state the superiority of a given model [48]. Using the best fitting parameters of the LQL ($\alpha/\beta = 1$ Gy, $\delta = 0.48$ Gy⁻¹, for LR) the BED calculated with the LQL model for a 20 Gy \times 1 treatment (95 Gy₁) is far less than that of a conventional 2 Gy \times 37 fractionation (184 Gy₁).

Interestingly, while the superiority of the LQL over the LQ is clear, the re-oxygenation model does not improve the performance of the LQ model. This cannot be used to conclude that reoxygenation does not play a role in the response to HDR-BT, and may simply be due to the simplicity of the model considered. In particular, the implementation of time variation of α and β due to reoxygenation is not dose/treatment dependent. The study of other more complex models accounting for hypoxia and reoxygenation has not been addressed in this work. For example, the models proposed by Kuperman & Lubich [49] or Jeong *et al.* [50].

The possibility that the LQ model overestimates the effect on prostate cancer of large doses per fraction has already been discussed in the context of external radiotherapy. Refs. [13, 14] found some evidence of a moderation of the LQ predicted effect with increasing doses. HDR-BT is delivered with more radical protocols than SBRT, >20 Gy in a single fraction, therefore the moderation of the effect can be more significant. Such radical hypofractionations are currently under investigation in external radiotherapy, with Zilli *et al.* currently investigating a 19 Gy \times 1 fractionation [51]. The results of this and future clinical trials will shed more light on the response of prostate cancer to extreme hypofractionation.

In this study we have focused on studying potential radiobiological reasons for the decline of tumor control in extremely hypofractionated HDR-BT. A potential dosimetric origin of such low control (target coverage, dose homogeneity) was not investigated. Recently, Kuperman & Lubich [52] modeled the effect of target dose heterogeneities on the BED, and found that dose heterogeneities reduce the BED compared to the homogeneous dose scenario, especially for hypofractionated schedules. This effect might also explain the origin of the observed loss of tumor control for extreme hypofractionation, and merits further investigation.

Our study presents some limitations. In particular, the limited number of patients/schedules and the heterogeneity of the dataset may increase the uncertainties and potential sources of bias of the analysis by including different studies that may use different margins, different dose constraints, different dose calculation algorithms, etc. Another limitation was that we only analyzed a limited number of dose–response models, as discussed above.

Conclusions

This analysis showed that the dose–response curves of prostate cancer to hypofractionated HDR-BT (LR and IR) is well described by the LQ model if the α/β is large (~ 90 Gy), far beyond the values reported in several radiobiological studies of dose–response of prostate cancer treated with external radiotherapy. If the α/β is constrained to be low (≤ 8 Gy), the LQ model cannot fit the dose–response curves and the LQL proves the superior model. This is in agreement with recent studies in external radiotherapy that have found evidence of a moderation of the LQ-predicted effect with increasing dose per fraction. This moderation of the effect with increasing dose per fraction might affect dose and/fractionation prescription for prostate cancer.

The origin of the loss of control of radical single-fraction HDR-BT treatments merits further investigation: while in this work a reoxygenation model did not fit the data as well as the LQL model, more complex reoxygenation models might provide better fits to the clinical data; also, target dose heterogeneity may lead to patterns like those observed experimentally (loss of effectiveness for extreme hypofractionations), and should be further explored.

Acknowledgements: This project has received funding from Ministerio de Ciencia e Innovación,

Agencia Estatal de Investigación and FEDER, UE (grant PID2021-128984OB-I00), and Xunta de Galicia, Axencia Galega de Innovación (grant IN607D 2022/02).

Declaration of competing interest: The authors declare that they have no competing interests.

References

1. Arcangeli G, Saracino B, Arcangeli S, et al. Moderate Hypofractionation in High-Risk, Organ-Confined Prostate Cancer: Final Results of a Phase III Randomized Trial. *J Clin Oncol*. 2017;35(17):1891-1897. doi:10.1200/JCO.2016.70.4189
2. Bryant CM, Henderson RH, Nichols RC, et al. Consensus Statement on Proton Therapy for Prostate Cancer. *Int J Part Ther*. 2021;8(2):1-16. doi:10.14338/IJPT-20-00031.1
3. Wu YY, Fan KH. Proton therapy for prostate cancer: current state and future perspectives. *Br J Radiol*. 2022;95(1131):20210670. doi:10.1259/bjr.20210670
4. Stish BJ, Davis BJ, Mynderse LA, McLaren RH, Deufel CL, Choo R. Low dose rate prostate brachytherapy. *Transl Androl Urol*. 2018;7(3):341-356. doi:10.21037/tau.2017.12.15
5. Mendez LC, Morton GC. High dose-rate brachytherapy in the treatment of prostate cancer. *Transl Androl Urol*. 2018;7(3):357-370. doi:10.21037/tau.2017.12.08
6. Crook J, Marbán M, Batchelar D. HDR Prostate Brachytherapy. *Semin Radiat Oncol*. 2020;30(1):49-60. doi:10.1016/j.semradonc.2019.08.003
7. Miralbell R, Roberts SA, Zubizarreta E, Hendry JH. Dose-fractionation sensitivity of prostate cancer deduced from radiotherapy outcomes of 5,969 patients in seven international institutional datasets: $\alpha/\beta = 1.4$ (0.9-2.2) Gy. *Int J Radiat Oncol Biol Phys*. 2012;82(1):e17-24. doi:10.1016/j.ijrobp.2010.10.075
8. Fowler JF. The radiobiology of prostate cancer including new aspects of fractionated radiotherapy. *Acta Oncol*. 2005;44(3):265-276. doi:10.1080/02841860410002824
9. Bentzen SM, Ritter MA. The alpha/beta ratio for prostate cancer: what is it, really? *Radiother Oncol*. 2005;76(1):1-3. doi:10.1016/j.radonc.2005.06.009
10. Nahum AE, Movsas B, Horwitz EM, Stobbe CC, Chapman JD. Incorporating clinical measurements of hypoxia into tumor local control modeling of prostate cancer: implications for the alpha/beta ratio. *Int J Radiat Oncol Biol Phys*. 2003;57(2):391-401. doi:10.1016/s0360-3016(03)00534-0
11. Hannan R, Tumati V, Xie XJ, et al. Stereotactic body radiation therapy for low and intermediate risk prostate cancer-Results from a multi-institutional clinical trial. *Eur J Cancer*. 2016;59:142-151. doi:10.1016/j.ejca.2016.02.014
12. Datta NR, Stutz E, Rogers S, Bodis S. Clinical estimation of α/β values for prostate cancer from isoeffective phase III randomized trials with moderately hypofractionated radiotherapy. *Acta Oncol*. 2018;57(7):883-894. doi:10.1080/0284186X.2018.1433874
13. Vogelius IR, Bentzen SM. Diminishing Returns From Ultrahypofractionated Radiation Therapy for Prostate Cancer. *Int J Radiat Oncol Biol Phys*. 2020;107(2):299-304. doi:10.1016/j.ijrobp.2020.01.010

14. Pardo-Montero J, González-Crespo I, Gómez-Caamaño A, Gago-Arias A. Radiobiological Meta-Analysis of the Response of Prostate Cancer to Different Fractionations: Evaluation of the Linear–Quadratic Response at Large Doses and the Effect of Risk and ADT. *Cancers*. 2023;15(14):3659. doi:10.3390/cancers15143659
15. Guirado D, Ruiz-Arebola S, Tornero-López AM, de la Vega JM, Prada PJ, Lallena AM. A radiobiological study of the schemes with a low number of fractions in high-dose-rate brachytherapy as monotherapy for prostate cancer. *J Contemp Brachytherapy*. 2020;12(2):193-200. doi:10.5114/jcb.2020.94492
16. Kirkpatrick JP, Meyer JJ, Marks LB. The linear-quadratic model is inappropriate to model high dose per fraction effects in radiosurgery. *Semin Radiat Oncol*. 2008;18(4):240-243. doi:10.1016/j.semradonc.2008.04.005
17. Wang JZ, Huang Z, Lo SS, Yuh WTC, Mayr NA. A generalized linear-quadratic model for radiosurgery, stereotactic body radiation therapy, and high-dose rate brachytherapy. *Sci Transl Med*. 2010;2(39):39ra48. doi:10.1126/scitranslmed.3000864
18. Song CW, Lee YJ, Griffin RJ, et al. Indirect Tumor Cell Death After High-Dose Hypofractionated Irradiation: Implications for Stereotactic Body Radiation Therapy and Stereotactic Radiation Surgery. *Int J Radiat Oncol Biol Phys*. 2015;93(1):166-172. doi:10.1016/j.ijrobp.2015.05.016
19. Guerrero M, Li XA. Extending the linear-quadratic model for large fraction doses pertinent to stereotactic radiotherapy. *Phys Med Biol*. 2004;49(20):4825-4835. doi:10.1088/0031-9155/49/20/012
20. Zeng H, Dai J, Cao D, et al. Safety and efficacy associated with single-fraction high-dose-rate brachytherapy in localized prostate cancer: a systematic review and meta-analysis. *Strahlenther Onkol*. 2023;199(6):525-535. doi:10.1007/s00066-023-02063-z
21. Hauswald H, Kamrava MR, Fallon JM, et al. High-Dose-Rate Monotherapy for Localized Prostate Cancer: 10-Year Results. *Int J Radiat Oncol Biol Phys*. 2016;94(4):667-674. doi:10.1016/j.ijrobp.2015.07.2290
22. Barkati M, Williams SG, Foroudi F, et al. High-dose-rate brachytherapy as a monotherapy for favorable-risk prostate cancer: a Phase II trial. *Int J Radiat Oncol Biol Phys*. 2012;82(5):1889-1896. doi:10.1016/j.ijrobp.2010.09.006
23. Strouthos I, Tselis N, Chatzikonstantinou G, et al. High dose rate brachytherapy as monotherapy for localised prostate cancer. *Radiother Oncol*. 2018;126(2):270-277. doi:10.1016/j.radonc.2017.09.038
24. Kukielka AM, Dąbrowski T, Walasek T, Olchawa A, Kudzia R, Dybek D. High-dose-rate brachytherapy as a monotherapy for prostate cancer--Single-institution results of the extreme fractionation regimen. *Brachytherapy*. 2015;14(3):359-365. doi:10.1016/j.brachy.2015.01.004
25. Jawad MS, Dilworth JT, Gustafson GS, et al. Outcomes Associated With 3 Treatment Schedules of High-Dose-Rate Brachytherapy Monotherapy for Favorable-Risk Prostate Cancer. *Int J Radiat Oncol Biol Phys*. 2016;94(4):657-666. doi:10.1016/j.ijrobp.2015.10.011
26. Prada PJ, Cardenal J, Blanco AG, et al. High-dose-rate interstitial brachytherapy as monotherapy in one fraction for the treatment of favorable stage prostate cancer: Toxicity and long-term biochemical results. *Radiother Oncol*. 2016;119(3):411-416. doi:10.1016/j.radonc.2016.04.006
27. Behmueller M, Tselis N, Zamboglou N, et al. High-Dose-Rate Brachytherapy as Monotherapy for Low- and Intermediate-Risk Prostate Cancer. Oncological Outcomes After a Median 15-Year Follow-Up. *Front Oncol*. 2021;11:770959. doi:10.3389/fonc.2021.770959
28. Johansson B, Olsén JS, Karlsson L, Lundin E, Lennernäs B. High-dose-rate brachytherapy as monotherapy for low- and intermediate-risk prostate cancer: long-term experience of Swedish single-center. *J Contemp Brachytherapy*. 2021;13(3):245-253. doi:10.5114/jcb.2021.105846
29. Morton G, McGuffin M, Chung HT, et al. Prostate high dose-rate brachytherapy as monotherapy for low and intermediate risk prostate cancer: Efficacy results from a randomized phase II clinical trial of one fraction of 19 Gy

or two fractions of 13.5 Gy. *Radiother Oncol.* 2020;146:90-96. doi:10.1016/j.radonc.2020.02.009

30. Tselis N, Tunn UW, Chatzikonstantinou G, et al. High dose rate brachytherapy as monotherapy for localised prostate cancer: a hypofractionated two-implant approach in 351 consecutive patients. *Radiat Oncol.* 2013;8:115. doi:10.1186/1748-717X-8-115
31. Nagore G, Lopez Guerra JL, Krumina E, et al. High dose rate brachytherapy for prostate cancer: A prospective toxicity evaluation of a one day schedule including two 13.5 Gy fractions. *Radiother Oncol.* 2018;127(2):219-224. doi:10.1016/j.radonc.2018.03.022
32. Yamazaki H, Masui K, Suzuki G, et al. Comparison of three moderate fractionated schedules employed in high-dose-rate brachytherapy monotherapy for clinically localized prostate cancer. *Radiother Oncol.* 2018;129(2):370-376. doi:10.1016/j.radonc.2018.07.026
33. Prada PJ, Ferri M, Cardenal J, et al. High-dose-rate interstitial brachytherapy as monotherapy in one fraction of 20.5 Gy for the treatment of localized prostate cancer: Toxicity and 6-year biochemical results. *Brachytherapy.* 2018;17(6):845-851. doi:10.1016/j.brachy.2018.06.002
34. Hannoun-Levi JM, Chand-Fouche ME, Pace-Loscos T, et al. Single fraction of HDR brachytherapy for prostate cancer: Results of the SiFEPI phase II prospective trial. *Clin Transl Radiat Oncol.* 2022;37:64-70. doi:10.1016/j.ctro.2022.08.007
35. Siddiqui ZA, Gustafson GS, Ye H, et al. Five-Year Outcomes of a Single-Institution Prospective Trial of 19-Gy Single-Fraction High-Dose-Rate Brachytherapy for Low- and Intermediate-Risk Prostate Cancer. *Int J Radiat Oncol Biol Phys.* 2019;104(5):1038-1044. doi:10.1016/j.ijrobp.2019.02.010
36. Yoshioka Y, Suzuki O, Isohashi F, et al. High-Dose-Rate Brachytherapy as Monotherapy for Intermediate- and High-Risk Prostate Cancer: Clinical Results for a Median 8-Year Follow-Up. *Int J Radiat Oncol Biol Phys.* 2016;94(4):675-682. doi:10.1016/j.ijrobp.2015.05.044
37. Rogers CL, Alder SC, Rogers RL, et al. High dose brachytherapy as monotherapy for intermediate risk prostate cancer. *J Urol.* 2012;187(1):109-116. doi:10.1016/j.juro.2011.09.050
38. Patel S, Demanes DJ, Ragab O, et al. High-dose-rate brachytherapy monotherapy without androgen deprivation therapy for intermediate-risk prostate cancer. *Brachytherapy.* 2017;16(2):299-305. doi:10.1016/j.brachy.2016.11.002
39. Hoskin P, Rojas A, Ostler P, Hughes R, Alonzi R, Lowe G. Single-dose high-dose-rate brachytherapy compared to two and three fractions for locally advanced prostate cancer. *Radiother Oncol.* 2017;124(1):56-60. doi:10.1016/j.radonc.2017.06.014
40. Fowler JF. The linear-quadratic formula and progress in fractionated radiotherapy. *Br J Radiol.* 1989;62(740):679-694. doi:10.1259/0007-1285-62-740-679
41. Stavrev P, Stavreva N, Genova B, et al. The Impact of Different Timing Schedules on Prostate HDR-Mono-Brachytherapy. A TCP Modeling Investigation. *Cancers.* 2021;13(19):4899. doi:10.3390/cancers13194899
42. Wakisaka Y, Yagi M, Sumida I, Takashina M, Ogawa K, Koizumi M. Impact of time-related factors on biologically accurate radiotherapy treatment planning. *Radiat Oncol.* 2018;13(1):30. doi:10.1186/s13014-018-0973-6
43. Bentzen SM, Tucker SL. Quantifying the position and steepness of radiation dose-response curves. *Int J Radiat Biol.* 1997;71(5):531-542. doi:10.1080/095530097143860
44. Royston P. Profile Likelihood for Estimation and Confidence Intervals. *Stata J.* 2007;7(3):376-387. doi:10.1177/1536867X0700700305
45. Akaike H. A new look at the statistical model identification. *IEEE Trans Autom Control.* 1974;19(6):716-723.

doi:10.1109/TAC.1974.1100705

46. Hudson JM, Loblaw A, McGuffin M, et al. Prostate high dose-rate brachytherapy as monotherapy for low and intermediate-risk prostate cancer: Efficacy results from a randomized phase II clinical trial of one fraction of 19 Gy or two fractions of 13.5 Gy: A 9-year update. *Radiother Oncol.* 2024 IN PRESS. doi:10.1016/j.radonc.2024.110381
47. Nickers P, Hermesse J, Deneufbourg JM, Vanbelle S, Lartigau E. Which α/β ratio and half-time of repair are useful for predicting outcomes in prostate cancer? *Radiother Oncol.* 2010;97(3):462-466. doi:10.1016/j.radonc.2010.06.006
48. Burnham KP, Anderson DR, eds. *Model Selection and Multimodel Inference*. Springer; 2004. doi:10.1007/b97636
49. Kuperman VY, Lubich LM. Effect of reoxygenation on hypofractionated radiotherapy of prostate cancer. *Med Phys.* 2020;47(10):5383-5391. doi:10.1002/mp.14343
50. Jeong J, Shoghi KI, Deasy JO. Modelling the interplay between hypoxia and proliferation in radiotherapy tumour response. *Phys Med Biol.* 2013;58(14):4897-4919. doi:10.1088/0031-9155/58/14/4897
51. Zilli T, Franzese C, Guckenberger M, et al. ONE SHOT - single shot radiotherapy for localized prostate cancer: 18-month results of a single arm, multicenter phase I/II trial. *Radiother Oncol.* 2024;194:110181. doi:10.1016/j.radonc.2024.110181
52. Kuperman VY, Lubich LM. Impact of target dose inhomogeneity on BED and EUD in lung SBRT. *Phys Med Biol.* 2021;66(1):01NT02. doi:10.1088/1361-6560/abd0d1

TABLES

Table 1: Overview of the characteristics of the schedules included in the analysis.

Risk	Number of schedules	Total number of patients	Number of patients per schedule (range)	Dose per fraction (range)	Total dose (range)	Overall treatment time (range)	Control at 5 years (range)	References
LR	21	1643	2 – 288	6.0 – 20.5 Gy	19 – 54 Gy	0 – 27 days	0.66 – 1.00	[21–35]
IR	22	1615	16 – 284	6.0 – 20.5 Gy	19 – 54 Gy	0 – 41 days	0.63 – 0.996	[21, 23, 25–30, 32, 33, 36–39]

Table 2: Best fits obtained with the LQ, LQL, and Stavrev’s reoxygenation model, LQ_{ST}, without or with (subscript SD) sublethal damage incomplete repair correction, separated by risk (low, LR, and intermediate risk, IR). The table shows best fitting parameters, maximum likelihood, AIC_c, and ΔAIC_c (referred to the LQ model) values. The symbol * indicates that the best-fitting parameter reached the edge of the constraint window.

Risk	Model	Parameters									-log(L)	AIC _c	ΔAIC _c
		α/β [Gy]	λ' [Gy day ⁻¹]	T_k [day]	δ [Gy ⁻¹]	φ	b [h ²]	T_{repair} [h]	D_{50} [Gy]	γ_{50}			
LR	LQ	100*	0.05	21.8	-	-	-	-	16.17	1.00	48.19	110.37	-
	LQL	5.0	0.24	29.8	0.73	-	-	-	16.64	0.96	47.37	112.74	-2.37
	LQ _{ST}	100*	0.09	28.1	-	4.2E-2	2.5E-7	-	16.21	0.99	48.11	118.84	-8.47
	LQ _{SD}	100*	0.11	27.6	-	-	-	0.17	16.62	1.04	48.05	114.10	-3.73
	LQL _{SD}	5.0	0.24	29.8	0.73	-	-	0.00*	16.64	0.96	47.37	117.35	-6.98
	LQ _{ST,SD}	100*	0.11	27.6	-	0.00*	0.00*	0.17	16.62	1.04	48.05	124.10	-13.73
IR	LQ	80.4	0	-	-	-	-	-	15.75	0.75	55.71	125.17	-
	LQL	1.0*	0	-	0.51	-	-	-	24.72	0.78	55.40	128.39	-3.22
	LQ _{ST}	84.2	0.09	0.1	-	1.50	7E-7	-	15.13	0.69	55.40	132.81	-7.63
	LQ _{SD}	77.6	0	-	-	-	-	1.00	15.90	0.75	55.71	129.02	-3.85
	LQL _{SD}	1.0*	0	-	0.51	-	-	0.00*	24.84	0.78	55.40	132.81	-7.64
	LQ _{ST,SD}	65.1	0.16	0	-	1.47	8e-7	0.724	17.00	0.80	55.02	137.13	-11.95

Table 3: Best fits obtained with the LQ, LQL, and Stavrev's reoxygenation model, LQ_{ST}, without or with (subscript SD) sublethal damage incomplete repair correction, separated by risk (low, LR, and intermediate risk, IR). The values of α/β were constrained to $1 \leq \alpha/\beta \leq 8$ Gy, to take into account the low α/β values typically reported for prostate cancer. The table shows best fitting parameters, maximum likelihood, AIC_c, and Δ AIC_c (referred to the LQ model) values. The symbol * indicates that the best-fitting parameter reached the edge of the constraint window.

Risk	Model	Parameters									$-\log(L)$	AIC _c	Δ AIC _c
		α/β [Gy]	λ' [Gy day ⁻¹]	T_k [day]	δ [Gy ⁻¹]	φ	b [h ²]	T_{repair} [h]	D_{50} [Gy]	γ_{50}			
LR	LQ	8.0*	1.00	35.54	-	-	-	-	41.0	1.84	60.21	134.43	-
	LQL	1.0*	1.33	31.45	0.48	-	-	-	27.2	1.09	47.23	112.45	21.98
	LQ _{ST}	8.0*	0.60	28.93	-	8.5E-3	5.4E-5	-	41.0	1.84	60.15	142.91	-8.48
	LQ _{SD}	8.0*	0.00	-	-	-	-	2.75	42.8	1.72	55.29	128.59	5.84
	LQL _{SD}	1.0*	1.33	31.45	0.48	-	-	0.00*	27.2	1.09	47.23	117.07	17.36
	LQ _{ST,SD}	8.0*	0.98	37.78	-	0.14	3.7E-5	3.62	40.9	1.33	54.34	136.67	-2.24
IR	LQ	8.0*	0.00*	-	-	-	-	-	39.0	1.43	66.67	147.10	-
	LQL	1.0*	0.00*	-	0.51	-	-	-	24.6	0.78	55.40	128.39	18.71
	LQ _{ST}	8.0*	0.00*	-	-	0.00*	0.00*	-	39.0	1.43	66.67	155.39	-8.30
	LQ _{SD}	8.0*	0.00*	-	-	-	-	3.00	41.0	1.34	61.59	140.78	6.31
	LQL _{SD}	1.0*	0.00*	-	0.51	-	-	0.00*	24.7	0.78	55.40	132.79	14.30
	LQ _{ST,SD}	8.0*	0.00*	-	-	0.00*	0.00*	3.00	41.0	1.34	61.59	150.26	-3.16

Table 4: 95% confidence intervals of best fitting parameters for the LQ and LQL models without incomplete repair correction, and LQ model with incomplete repair correction (LQ_{SD}). Results are separated by risk, low (LR) and intermediate (IR). The symbol * indicates that the parameter value reached the edge of the constraint window.

Risk	Model	Parameters						
		α/β [Gy]	λ' [Gy day ⁻¹]	T_k [day]	δ [Gy ⁻¹]	T_{repair} [h]	D_{50} [Gy]	γ_{50}
LR	LQ	[7.1, 8*]	[0*, 2*]	[0*, 42*]	-	-	[36.0, 44.4]	[1.4, 2.3]
	LQL	[1*, 8*]	[0*, 2*]	[0*, 42*]	[0.10, 1*]	-	[12.3, 48.7]	[0.7, 1.6]
	LQ _{SD}	[6.6, 8*]	[0*, 2*]	[0*, 42*]	-	[1.6, 4.9]	[36.0, 47.3]	[1.2, 2.1]
IR	LQ	[7.4, 8*]	[0*, 2*]	[0*, 42*]	-	-	[32.1, 43.2]	[1.0, 1.9]
	LQL	[1*, 8*]	[0*, 2*]	[0*, 42*]	[0.06, 1*]	-	[10.2, 44.4]	[0.5, 1.3]
	LQ _{SD}	[6.8, 8*]	[0*, 2*]	[0*, 42*]	-	[1.6, 6*]	[31.5, 45.8]	[0.7, 1.8]

FIGURES




Figure 1: Best fits of LQ, LQL, and Stavrev's reoxygenation models to dose–response data for prostate cancer treated with HDR-BT (low risk, top panels; intermediate risk, bottom panels). The α/β ratio was allowed to lie in a large interval ($1 \leq \alpha/\beta \leq 100$ Gy). Clinical data (*) and 95% confidence intervals (bars), and modeled curves (solid lines). Single fraction schedules are highlighted as red circles.




Figure 2: Best fits of LQ, LQL, and Stavrev's reoxygenation models to dose–response data for prostate cancer treated with HDR-BT (low risk, top panels; intermediate risk, bottom panels). The α/β ratio was constrained to be low ($1 \leq \alpha/\beta \leq 8$ Gy). Clinical data (*) and 95% confidence intervals (bars), and modeled curves (solid lines). Single fraction schedules are highlighted as red circles.