




MR leave-one-out sensitivity analysis for 'Dimethylglycine levels' on 'Parkinson's disease || id:ieu-b-7'




MR leave-one-out sensitivity analysis for '1-(1-enyl-palmitoyl)-2-arachidonoyl-gpc (p-16:0/20:4) levels' on 'Parkinson's disease || id:ieu-b-7'




MR leave-one-out sensitivity analysis for '1-stearoyl-2-docosaheptaenoyl-gpc (18:0/22:6) levels' on 'Parkinson's disease || id:ieu-b-7'




MR leave-one-out sensitivity analysis for 'Gluconate levels' on 'Parkinson's disease || id:ieu-b-7'




MR leave-one-out sensitivity analysis for 'Glycerophosphoinositol levels' on 'Parkinson's disease || id:ieu-b-7'




MR leave-one-out sensitivity analysis for
'Oxalate (ethanedioate) levels' on 'Parkinson's disease || id:ieu-b-7'



MR leave-one-out sensitivity analysis for 'X-12015 levels' on 'Parkinson's disease || id:ieu-b-7'



MR leave-one-out sensitivity analysis for 'X-12411 levels' on 'Parkinson's disease || id:ieu-b-7'



MR leave-one-out sensitivity analysis for 'X-23587 levels' on 'Parkinson's disease || id:ieu-b-7'