

Comparing different approaches of modelling the effects of temperature and rainfall on malaria transmission in high and low transmission settings

Gladstone T. Madito^{1*} and Sheetal P. Silal^{1,2}

¹ Modelling And Simulations Hub, Africa (MASHA), Department of Statistical Sciences, University of Cape Town, Cape Town, South Africa.

² Centre for Global Health, Nuffield Department of Medicine, University of Oxford, Oxford, United Kingdom.

*Corresponding author(s). E-mail(s): mdt gla001@myuct.ac.za;

Appendix A Supplementary plots






Fig. A1: Mean monthly temperature and rainfall trends used to simulate the models for a period of 6 years from year 12 to 18.



(a) Daily mosquito biting rate. (b) Daily larval mortality rate.



(c) Daily mosquito mortality rate.

Fig. A2: Thermal response curves of temperature regulated mosquito life-cycle characteristics examined in Approach A.







Fig. A3: Thermal response curves of temperature regulated mosquito life-cycle characteristics examined in Approach B.



(a) Mosquito biting rate. (b) Larval mortality rate.



(c) Mosquito mortality rate. (d) Parasite development rate.



(e) Larval development period.

Fig. A4: Thermal response curves of temperature regulated mosquito life-cycle characteristics examined in Approach C.