

AP2S1 Variant	REVEL score	CADD score	Source
R10W	0.471	32.0	GeneDx
R10Q	0.273	28.5	GeneDx, DiscovEHR
K18E	0.699	30.0	Genomics England
K18N	0.673	27.0	GeneDx
R61H	0.447	29.8	GeneDx

**Supplementary Table 1.** Computational analysis of AP2 $\sigma$ 2 variants using Rare Exome Variant Ensemble Learner (REVEL) and Combined Annotation Dependent Deletion (CADD). REVEL scores are based on a combination of scores from 13 tools (MutPred, FATHMM v2.3, VEST 3.0, PolyPhen-2, SIFT, PROVEAN, Mutation Assessor, MutationTaster, LRT, GERP++, SiPhy, phyloP and phastCons) (Ioannidis et al. *Am. J. Hum. Genet.* 2016 99:877-885). Scores range from 0 to 1 with higher scores reflecting a greater likelihood that the variant is disease-causing. 75.4% of disease mutations but only 10.9% of neutral variants have a REVEL score >0.5. CADD scores are based on genomic features from surrounding sequence context, gene model annotations, evolutionary constraint, epigenetic measurements and functional predictions (Kircher et al. 2014 *Nat. Genet.* 46:310-315). The higher the CADD score the more the variant is predicted to be deleterious. Thus, variants with CADD scores >10, >20 and >30 are predicted to be among the 10%, 1% and 0.1% most deleterious possible substitutions in the human genome.