

Supplementary file

Article title: A comparison of prevalence estimates of smoking, alternative nicotine and alcohol use in Great Britain collected via telephone versus face-to-face: Smoking and Alcohol Toolkit surveys

Contents

1. Measures	1
2. Unweighted and weighted data (2022 and 2024)	5
3. Comparing 2024 data using telephone data from February 2024.....	8
4. Comparing 2024 data using telephone data from January to March 2024	12
5. Exploring differences in e-cigarette use	15

1. Measures

Age: categories into 16-24/25-34/35-44/45-54/55-64/65+ years [\[asked to all\]](#)

Gender: categorised into men, women, or non-binary [\[asked to all\]](#)

Social grade: as a measure of socioeconomic position, categorised into AB (high or intermediate managerial, administrative or professional), C1 (supervisory, clerical and junior managerial, administrative or professional), C2 (skilled manual workers), D (semi and unskilled manual workers), E (state pensioners, casual or lowest grade workers, unemployed with state benefits only).¹ [\[asked to all\]](#)

Smoking status: assessed with the following question, which was [asked to all](#),

“Which of the following best applies to you? Please note we are referring to cigarettes and other kinds of tobacco that you set light to and NOT electronic or ‘heat-not-burn’ cigarettes.

1. *I smoke cigarettes (including hand-rolled) every day*
2. *I smoke cigarettes (including hand-rolled), but not every day*
3. *I do not smoke cigarettes at all, but I do smoke tobacco of some kind (eg. Pipe, cigar or shisha)*
4. *I have stopped smoking completely in the last year*
5. *I stopped smoking completely more than a year ago*
6. *I have never been a smoker (i.e. smoked for a year or more)*
7. *Don’t know”*

Smoking in the past year: binary variable – yes, if one of answer options 1-4 selected above [\[asked to all\]](#)

Tried to stop smoking in past year: binary variable – assessed with the following question which was [asked to those who smoked in the past year](#) (see above),

“How many serious attempts to stop smoking have you made in the last 12 months? By serious attempt I mean you decided that you would try to make sure you never smoked again. Please include any attempt that you are currently making and please include any successful attempt made within the last year.”

¹ Collis, D. (2009). Social grade: A classification tool – Bite sized through piece (Ipsos MediaCT, Ipsos Mori). <https://www.ipsos.com/en-uk/social-grade>

Categorised as tried to stop smoking in the past year if answered with at least one attempt.

Quit smoking in past year among those who smoked in past year: binary variable – assessed with the following question, which was [asked to all](#),

“Which of the following best applies to you? Please note we are referring to cigarettes and other kinds of tobacco that you set light to and NOT electronic or 'heat-not-burn' cigarettes.”

and the answer response, *“I have stopped smoking completely in the last year”*.

Proportion calculated by dividing the number of people with that response by the number of people who smoked in the past year (see above).

NRT/HTP/e-cigarette/pouch use (in people smoking in the past year or in total population): assessed with the following questions,

- *“Which, if any, of the following are you currently using to help you cut down the amount you smoke?”* [\[asked to people who currently smoke\]](#)
- *“Do you regularly use any of the following in situations when you are not allowed to smoke?”* [\[asked to people who currently smoke\]](#)
- *“Can I check, are you using any of the following?”* [\[asked to people who have not smoke in past year\]](#)
- *“Can I check, are you using any of the following either to help you stop smoking, to help you cut down or for any other reason at all?”* [\[asked to people who smoked in past year\]](#)

Categorised as using nicotine replacement therapy (NRT) if answered *“Nicotine gum”, “Nicotine replacement lozenges/tablets”, “Nicotine replacement inhaler”, “Nicotine replacement nasal spray”, “Nicotine mouthspray”* or *“Nicotine patch”* to any of these questions.

Categorised as using heated tobacco products (HTP) if answered *“Heat-not-burn cigarette (e.g. IQOS with HEETS, heatsticks)”* to any of these questions.

Categorised as using e-cigarettes if answered *“Electronic cigarette or vaping device”* or *“Juil”* to any of these questions.

Categorised as using nicotine pouches if answered *“Tobacco-free nicotine pouch/pod or 'white pouches' that you place on your gum (e.g., Zyn, On!, Nordic Spirit, Velo, Lyft, Skruf)”* to any of these questions.

E-cigarette device used: assessed with the following question, which was [asked to those who reported using e-cigarettes](#) (see above),

“Which of the following do you mainly use...?”

1. *A disposable e-cigarette or vaping device (non –rechargeable)*
2. *An e-cigarette or vaping device that uses replaceable pre-filled cartridges (rechargeable)*
3. *An e-cigarette or vaping device with a tank that you refill with liquids (rechargeable)*
4. *A modular system that you refill with liquids (you use your own combination of separate devices: batteries, atomizers, etc.)*
5. *Don't know”*

AUDIT-C questions

“The next few questions form part of a study about consumption of alcohol. We understand that this is a highly sensitive topic and would therefore like to remind you that any information you give me is strictly confidential and will be used for research purposes only. Some questions asked may not necessarily apply to you.

These first few questions ask about the alcohol you have drunk in the last 6 months, including about how many standard drinks you have consumed. Please note that 1 standard drink equals 1 unit of alcohol. So, for example, a small glass of wine or a single measure of spirits is 1 standard drink, while a pint of regular beer or lager is equal to 2 standard drinks or 2 units, and a bottle of wine is equal to 9 units. If you are unsure, please ask me to help you work it out.

Please be aware that all your answers will be handled confidentially.

AUDIT 1. How often do you have a drink containing alcohol?

- i. Never*
- ii. Monthly or less*
- iii. 2 to 4 times a month*
- iv. 2 to 3 times a week*
- v. 4 to 5 times a week*
- vi. 6 or more times a week*
- vii. Don't know*
- viii. Refused*

AUDIT 2. How many standard drinks containing alcohol do you have on a typical day when you are drinking?

- i. 1 to 2*
- ii. 3 to 4*
- iii. 5 to 6*
- iv. 7 to 9*
- v. 10 to 12*
- vi. 13 to 15*
- vii. 16 or more*
- viii. Don't know*
- ix. Refused*

AUDIT 3. How often do you have six or more standard drinks on one occasion?

- i. Never*
- ii. Less than monthly*
- iii. Monthly*
- iv. Weekly*
- v. Daily or almost daily*
- vi. Don't know*
- vii. Refused”*

AUDIT 1 asked to all, AUDIT 2 and AUDIT 3 to those who responded with one of answer options ii to vi to AUDIT 1 (i.e., those who reported alcohol consumption).

Increasing or higher risk drinking: asked to all and operationalised as AUDIT-C score of 5 or above

AUDIT 1 – answer options and their respective value for the AUDIT-C score:

- Never: 0
- Monthly or less: 1
- 2 to 4 times a month: 2
- 2 to 3 times a week: 3
- 4 to 5 times a week or 6 or more times a week: 4

AUDIT 2 – answer options and their respective value for the AUDIT-C score:

- 1 to 2: 0
- 3 to 4: 1
- 5 to 6: 2
- 7 to 9: 3
- 10 to 12 or 13 to 15: 4

AUDIT 3 – answer options and their respective value for the AUDIT-C score:

- Never: 0
- Less than monthly: 1
- Monthly: 2
- Weekly: 3
- Daily or almost daily: 4

2. Unweighted and weighted data (2022 and 2024)

This section includes face-to-face data collected in March 2022 and January to March 2024 and telephone data collected in March 2022 and February 2024.

Table S1: Sociodemographic estimates from face-to-face (F2F) and telephone (Tel) data collected in 2022 and 2024 (N_{F2F}=4381 and N_{Tel}=4982).

Sociodemographic characteristics		Unweighted				Weighted			
		F2F, n	F2F, % (95% CI)	Tel, n	Tel, % (95% CI)	F2F, n	F2F, % (95% CI)	Tel, n	Tel, % (95% CI)
Age	16-24 years	602	13.7 (12.7-14.7)	507	10.2 (9.4-11.0)	574	13.1 (12.0-14.2)	657	13.2 (11.9-14.4)
	25-34 years	654	14.9 (13.8-16.0)	650	13.0 (12.1-13.9)	727	16.6 (15.4-17.8)	829	16.6 (15.3-18.0)
	35-44 years	687	15.7 (14.6-16.8)	630	12.6 (11.7-13.5)	687	15.7 (14.5-16.8)	778	15.6 (14.3-16.9)
	45-54 years	598	13.6 (12.6-14.6)	872	17.5 (16.4-18.6)	717	16.4 (15.1-17.6)	814	16.3 (15.2-17.5)
	55-64 years	699	16.0 (14.9-17.1)	855	17.2 (16.2-18.2)	675	15.4 (14.3-16.5)	762	15.3 (14.1-16.4)
	65+ years	1141	26.0 (24.7-27.3)	1465	29.4 (28.1-30.7)	1001	22.9 (21.6-24.1)	1138	22.9 (21.6-24.1)
	Refused	0	0 (0-0)	3	0.1 (0.0-0.2)	0	0 (0-0)	3	0.1 (0.0-0.1)
Gender	Men	2192	50.1 (48.6-51.6)	2493	50.3 (48.9-51.7)	2142	49.0 (47.4-50.5)	2417	48.8 (47.1-50.4)
	Women	2181	49.9 (48.4-51.4)	2440	49.2 (47.8-50.6)	2231	51.0 (49.5-52.6)	2516	50.7 (49.1-52.4)
	Non-binary	0	0 (0-0)	25	0.5 (0.3-0.7)	0	0 (0-0)	25	0.5 (0.3-0.7)
Social grade	AB	1014	23.1 (21.9-24.3)	1290	25.9 (24.7-27.1)	1155	26.4 (24.9-27.8)	1312	26.3 (24.9-27.8)
	C1	1586	36.2 (34.8-37.6)	2086	41.9 (40.5-43.3)	1298	29.6 (28.3-31.0)	1477	29.6 (28.3-31.0)
	C2	782	17.8 (16.7-18.9)	738	14.8 (13.8-15.8)	891	20.3 (19.0-21.6)	1010	20.3 (18.8-21.7)
	D	571	13.0 (12.0-14.0)	359	7.2 (6.5-7.9)	637	14.5 (13.4-15.7)	722	14.5 (13.0-16.0)
	E	428	9.8 (8.9-10.7)	509	10.2 (9.4-11.0)	400	9.1 (8.3-10.0)	460	9.2 (8.4-10.1)

Table S2: Estimates for smoking and use of other nicotine products from face-to-face (F2F) and telephone (Tel) data collected in 2022 and 2024 (N_{F2F}=4381 and N_{Tel}=4982).

Smoking and use of other nicotine products		Unweighted				Weighted			
		F2F, n	F2F, % (95% CI)	Tel, n	Tel, % 95% CI	F2F, n	F2F, % (95% CI)	Tel, n	Tel, % (95% CI)
Tobacco use in total population	Daily cigarette smoking	474	10.8 (9.9-11.7)	478	9.6 (8.8-10.4)	485	11.1 (10.1-12.1)	526	10.6 (9.5-11.7)
	Non-daily cigarette smoking	121	2.8 (2.3-3.3)	145	2.9 (2.4-3.4)	120	2.7 (2.2-3.3)	170	3.4 (2.8-4.1)
	Non-cigarette smoking	41	0.9 (0.6-1.2)	77	1.5 (1.2-1.8)	43	1.0 (0.7-1.3)	87	1.7 (1.3-2.2)
	Stopped last year	66	1.5 (1.1-1.9)	94	1.9 (1.5-2.3)	65	1.5 (1.1-1.9)	114	2.3 (1.7-2.8)
	Stopped 1+ year ago	785	17.9 (16.8-19.0)	1178	23.6 (22.4-24.8)	765	17.5 (16.3-18.6)	1114	22.4 (21.0-23.7)
	Never smoked	2883	65.8 (64.4-67.2)	2953	59.3 (57.9-60.7)	2892	66.0 (64.5-67.5)	2912	58.5 (56.8-60.1)
	Don't know	11	0.3 (0.2-0.4)	57	1.1 (0.8-1.4)	12	0.3 (0.1-0.4)	59	1.2 (0.8-1.6)
Other nicotine use in total population	NRT use	149	3.4 (2.9-3.9)	135	2.7 (2.2-3.2)	149	3.4 (2.8-4.0)	139	2.8 (2.2-3.4)
	HTP use	6	0.1 (0.0-0.2)	12	0.2 (0.1-0.3)	7	0.2 (0.0-0.3)	13	0.3 (0.1-0.5)
	E-cigarette use	259	5.9 (5.2-6.6)	417	8.4 (7.6-9.2)	279	6.4 (5.6-7.1)	520	10.4 (9.3-11.5)
	Pouch use	43	1.0 (0.7-1.3)	19	0.4 (0.2-0.6)	39	0.9 (0.6-1.2)	24	0.5 (0.2-0.7)
Quitting/other nicotine use in people smoking in past year (N _{F2F} =702, N _{Tel} =794)	Tried to stop in past year	232	35.7 (32.0-39.4)	263	34.9 (31.5-38.3)	241	36.2 (32.3-40.1)	292	34.7 (30.6-38.8)
	Quit in past year	66	9.4 (7.2-11.6)	94	11.8 (9.6-14.0)	65	9.1 (6.9-11.4)	114	12.7 (9.8-15.6)
	NRT use	123	17.5 (14.7-20.3)	102	12.8 (10.5-15.1)	122	17.1 (14.2-20.0)	109	12.1 (9.5-14.8)
	HTP use	5	0.7 (0.1-1.3)	10	1.3 (0.5-2.1)	6	0.8 (0.1-1.6)	11	1.2 (0.2-2.3)
	E-cigarette use	186	26.5 (23.2-29.8)	206	25.9 (22.9-28.9)	192	27.0 (23.5-30.5)	263	29.3 (25.4-33.3)
	Pouch use	19	2.7 (1.5-3.9)	13	1.6 (0.7-2.5)	21	2.9 (1.5-4.2)	19	2.1 (0.8-3.3)
Main device type used by e-cigarette users (N _{F2F} =259, N _{Tel} =417)	Disposable	96	40.3 (34.1-46.5)	124	31.2 (26.6-35.8)	98	38.2 (31.7-44.6)	166	33.6 (28.1-39.2)
	Rechargeable with pre-filled cartridges	38	16.0 (11.3-20.7)	43	10.8 (7.8-13.8)	38	14.8 (10.2-19.4)	60	12.1 (8.1-16.1)
	Rechargeable with tank to refill	73	30.7 (24.8-36.6)	165	41.5 (36.7-46.3)	84	32.8 (26.4-39.2)	189	38.2 (32.7-43.8)
	Mod system	31	13.0 (8.7-17.3)	54	13.6 (10.2-17.0)	37	14.2 (9.4-19.1)	69	14.0 (9.9-18.1)
	Don't know	0	0 (0-0)	12	3.0 (1.3-4.7)	0	0 (0-0)	10	2.1 (0.7-3.4)

Abbreviations: NRT, nicotine replacement therapy; HTP, heated tobacco product.

Table S3: Estimates for alcohol use from face-to-face (F2F) and telephone (Tel) data collected in 2022 and 2024 (N_{F2F}=4381 and N_{Tel}=4982).

Alcohol use		Unweighted				Weighted			
		F2F, n	F2F, % (95% CI)	Tel, n	Tel, % (95% CI)	F2F, n	F2F, % (95% CI)	Tel, n	Tel, % (95% CI)
AUDIT 1	Never	1408	32.1 (30.7-33.5)	1208	24.2 (23.0-25.4)	1363	31.1 (29.7-32.5)	1246	25.0 (23.5-26.5)
	Monthly or less	930	21.2 (20.0-22.4)	1030	20.7 (19.6-21.8)	954	21.8 (20.5-23.1)	1072	21.5 (20.1-22.9)
	2-4 per month	750	17.1 (16.0-18.2)	918	18.4 (17.3-19.5)	782	17.8 (16.6-19.1)	925	18.6 (17.3-19.9)
	2-3 per week	817	18.6 (17.4-19.8)	1141	22.9 (21.7-24.1)	823	18.8 (17.6-20.0)	1082	21.7 (20.4-23.1)
	4-5 per week	210	4.8 (4.2-5.4)	314	6.3 (5.6-7.0)	203	4.6 (4.0-5.3)	297	6.0 (5.2-6.7)
	6+ per week	229	5.2 (4.5-5.9)	282	5.7 (5.1-6.3)	222	5.1 (4.4-5.7)	274	5.5 (4.8-6.2)
	Don't know	4	0.1 (0.0-0.2)	43	0.9 (0.6-1.2)	4	0.1 (0.0-0.2)	46	0.9 (0.6-1.3)
	Refused	33	0.8 (0.5-1.1)	46	0.9 (0.6-1.2)	31	0.7 (0.5-1.0)	40	0.8 (0.5-1.1)
AUDIT 2 (N _{F2F} =2973, N _{Tel} =3774)	1-2	1425	48.3 (46.5-50.1)	1715	45.4 (43.8-47.0)	1439	48.0 (46.1-49.9)	1667	44.6 (42.7-46.5)
	3-4	824	27.9 (26.3-29.5)	957	25.4 (24.0-26.8)	849	28.3 (26.6-30.0)	939	25.1 (23.5-26.8)
	5-6	341	11.6 (10.4-12.8)	425	11.3 (10.3-12.3)	354	11.8 (10.6-13.1)	424	11.3 (10.1-12.5)
	7-9	175	5.9 (5.0-6.8)	268	7.1 (6.3-7.9)	172	5.7 (4.8-6.6)	273	7.3 (6.3-8.3)
	10-12	86	2.9 (2.3-3.5)	135	3.6 (3.4-2.0)	83	2.8 (2.2-3.4)	139	3.7 (2.9-4.5)
	13-15	37	1.3 (0.9-1.7)	34	0.9 (0.6-1.2)	38	1.3 (0.8-1.7)	48	1.3 (0.8-1.8)
	16+	37	1.3 (0.9-1.7)	72	1.9 (1.5-2.3)	37	1.2 (0.8-1.7)	67	1.8 (1.3-2.3)
	Don't know	11	0.4 (0.2-0.6)	106	2.8 (2.3-3.3)	11	0.4 (0.1-0.6)	129	3.5 (2.6-4.3)
Refused	14	0.5 (0.3-0.7)	62	1.6 (1.2-2.0)	14	0.5 (0.2-0.8)	51	1.4 (1.0-1.8)	
AUDIT 3 (N _{F2F} =2973, N _{Tel} =3774)	Never	1403	47.6 (45.8-49.4)	1557	41.3 (39.7-42.9)	1397	46.6 (44.7-48.5)	1504	40.2 (38.4-42.1)
	Less than monthly	802	27.2 (25.6-28.8)	1073	28.4 (27.0-29.8)	851	28.4 (26.7-30.1)	1099	29.4 (27.7-31.2)
	Monthly	388	13.2 (12.0-14.4)	556	14.7 (13.6-15.8)	393	13.1 (11.8-14.4)	558	14.9 (13.5-16.3)
	Weekly	300	10.2 (9.1-11.3)	414	11.0 (10.0-12.0)	299	10.0 (8.8-11.1)	412	11.0 (9.8-12.3)
	Daily or almost daily	35	1.2 (0.8-1.6)	71	1.9 (1.5-2.3)	35	1.2 (0.8-1.6)	73	1.9 (1.4-2.5)
	Don't know	7	0.2 (0.0-0.4)	51	1.4 (1.0-1.8)	8	0.3 (0.1-0.5)	49	1.3 (0.9-1.8)
Refused	15	0.5 (0.2-0.8)	52	1.4 (1.0-1.8)	15	0.5 (0.2-0.8)	41	1.1 (0.7-1.5)	
AUDIT-C ≥5		1090	25.2 (23.9-26.5)	1455	30.6 (29.3-31.9)	1095	25.3 (23.9-26.7)	1447	30.4 (28.9-32.0)

AUDIT 1: how often drinking; AUDIT 2: how many standard drinks on a typical drinking day; AUDIT 3: how often ≥6 standard drinks on one occasion; AUDIT-C ≥5: increasing and higher risk drinking.

3. Comparing 2024 data using telephone data from February 2024

This section includes face-to-face data collected between January and March 2024 and telephone data collected in February 2024.

Table S4: Missing values during face-to-face (F2F) and telephone (Tel) data collected in 2024 (F2F: Jan-Mar; Tel: Feb; N_{F2F}=2317, N_{Tel}=2375).

Variable	F2F missing values		Tel missing values	
	n	% (95% CI)	n	% (95% CI)
Age	0	0 (0-0)	0	0 (0-0)
Gender	8	0.3 (0.1-0.5)	24	1.0 (0.6-1.4)
Social grade	0	0 (0-0)	0	0 (0-0)
Smoking status	0	0 (0-0)	0	0 (0-0)
AUDIT 1	0	0 (0-0)	0	0 (0-0)
AUDIT 2 and 3 (N _{F2F} =1541, N _{Tel} =1766) ¹	1	0.1 (0.0-0.3)	0	0 (0-0)
AUDIT-C ≥5	18	0.8 (0.4-1.2)	83	3.5 (2.8-4.2)
Quit attempt in past year (N _{F2F} =317, N _{Tel} =364) ²	20	6.3 (3.6-9.0)	13	3.6 (1.7-5.5)
E-cigarette device used (N _{F2F} =139, N _{Tel} =241) ³	13	9.4 (4.5-14.3)	2	0.8 (0.0-1.9)

¹ only assessed for people who reported drinking alcohol for AUDIT 1 question.

² only assessed for people who reported smoking in the past year.

³ only assessed for people who reported using e-cigarettes.

Table S5: Sociodemographic estimates from face-to-face (F2F) and telephone (Tel) data collected in 2024 (F2F: Jan-Mar; Tel: Feb; N_{F2F}=2317 and N_{Tel}=2375).

Sociodemographic characteristics		Unweighted				Weighted			
		F2F, n	F2F, % (95% CI)	Tel, n	Tel, % (95% CI)	F2F, n	F2F, % (95% CI)	Tel, n	Tel, % (95% CI)
Age	16-24 years	257	11.1 (9.8-12.4)	234	9.9 (8.7-11.1)	298	12.9 (11.3-14.4)	306	12.9 (11.2-14.6)
	25-34 years	333	14.4 (13.0-15.8)	277	11.7 (10.4-13.0)	384	16.6 (14.9-18.3)	392	16.5 (14.6-18.4)
	35-44 years	375	16.2 (14.7-17.7)	326	13.7 (12.3-15.1)	367	15.8 (14.3-17.4)	375	15.8 (14.1-17.5)
	45-54 years	314	13.6 (12.2-15.0)	403	17.0 (15.5-18.5)	374	16.2 (14.5-17.8)	383	16.1 (14.5-17.7)
	55-64 years	361	15.6 (14.1-17.1)	409	17.2 (15.7-18.7)	364	15.7 (14.1-17.2)	373	15.7 (14.1-17.3)
	65+ years	677	29.2 (27.3-31.1)	725	30.5 (28.6-32.4)	530	22.9 (21.2-24.5)	546	23.0 (21.2-24.7)
	Refused	0	0 (0-0)	1	0.0 (0.0-0.1)	0	0 (0-0)	1	0.0 (0.0-0.1)
Gender	Men	1167	50.5 (48.5-52.5)	1233	52.4 (50.4-54.4)	1132	49.0 (46.9-51.2)	1153	49.0 (46.7-51.4)
	Women	1142	49.5 (47.5-51.5)	1118	47.6 (45.6-49.6)	1177	51.0 (48.8-53.1)	1198	51.0 (48.6-53.3)
	Non-binary	0	0 (0-0)	0	0 (0-0)	0	0 (0-0)	0	0 (0-0)
Social grade	AB	606	26.2 (24.4-28.0)	687	28.9 (27.1-30.7)	610	26.3 (24.4-28.2)	624	26.3 (24.3-28.2)
	C1	782	33.8 (31.9-35.7)	936	39.4 (37.4-41.4)	688	29.7 (27.8-31.6)	704	29.7 (27.7-31.6)
	C2	411	17.7 (16.1-19.3)	342	14.4 (13.0-15.8)	471	20.3 (18.5-22.1)	482	20.3 (18.3-22.3)
	D	302	13.0 (11.6-14.4)	178	7.5 (6.4-8.6)	336	14.5 (12.9-16.1)	344	14.5 (12.4-16.6)
	E	216	9.3 (8.1-10.5)	232	9.8 (8.6-11.0)	212	9.1 (7.9-10.3)	221	9.3 (8.0-10.5)

Table S6: Estimates for smoking and use of other nicotine products from face-to-face (F2F) and telephone (Tel) data collected in 2024 (F2F: Jan-Mar; Tel: Feb; N_{F2F}=2317 and N_{Tel}=2375).

Smoking and use of other nicotine products		Unweighted				Weighted			
		F2F, n	F2F, % (95% CI)	Tel, n	Tel, % 95% CI	F2F, n	F2F, % (95% CI)	Tel, n	Tel, % (95% CI)
Tobacco use in total population	Daily cigarette smoking	212	9.1 (7.9-10.3)	208	8.8 (7.7-9.9)	230	9.9 (8.6-11.2)	233	9.8 (8.4-11.3)
	Non-daily cigarette smoking	47	2.0 (1.4-2.6)	75	3.2 (2.5-3.9)	47	2.0 (1.4-2.7)	87	3.7 (2.8-4.6)
	Non-cigarette smoking	25	1.1 (0.7-1.5)	28	1.2 (0.8-1.6)	26	1.1 (0.7-1.6)	34	1.4 (0.8-2.0)
	Stopped last year	33	1.4 (0.9-1.9)	53	2.2 (1.6-2.8)	34	1.5 (0.9-2.0)	62	2.6 (1.8-3.4)
	Stopped 1+ year ago	445	19.2 (17.6-20.8)	582	24.5 (22.8-26.2)	423	18.2 (16.6-19.9)	563	23.7 (21.8-25.6)
	Never smoked	1551	66.9 (65.0-68.8)	1411	59.4 (57.4-61.4)	1553	67.0 (65.0-69.1)	1376	57.9 (55.6-60.2)
	Don't know	4	0.2 (0.0-0.4)	18	0.8 (0.5-1.1)	3	0.1 (0.0-0.3)	20	0.9 (0.4-1.3)
Other nicotine use in total population	NRT use	69	3.0 (2.3-3.7)	65	2.7 (2.0-3.4)	69	3.0 (2.3-3.7)	70	3.0 (2.2-3.7)
	HTP use	2	0.1 (0.0-0.2)	7	0.3 (0.1-0.5)	2	0.1 (0.0-0.2)	6	0.3 (0.1-0.5)
	E-cigarette use	139	6.0 (5.0-7.0)	241	10.1 (8.9-11.3)	164	7.1 (5.9-8.2)	299	12.6 (10.9-14.3)
	Pouch use	32	1.4 (0.9-1.9)	10	0.4 (0.1-0.7)	27	1.2 (0.8-1.6)	12	0.5 (0.1-0.9)
Quitting/other nicotine use in people smoking in past year (N _{F2F} =317, N _{Tel} =364)	Tried to stop in past year	113	38.0 (32.5-43.5)	118	33.6 (28.7-38.5)	121	38.0 (32.2-43.8)	132	32.7 (27.1-38.4)
	Quit in past year	33	10.4 (7.0-13.8)	53	14.6 (11.0-18.2)	34	10.0 (6.6-13.5)	62	15.0 (10.8-19.2)
	NRT use	43	13.6 (9.8-17.4)	49	13.5 (10.0-17.0)	43	12.7 (9.0-16.3)	55	13.1 (9.3-16.9)
	HTP use	1	0.3 (0.0-0.9)	5	1.4 (0.2-2.6)	1	0.4 (0.0-1.0)	4	1.0 (0.1-1.9)
	E-cigarette use	66	20.8 (16.3-25.3)	105	28.8 (24.1-33.5)	78	23.0 (18.0-28.0)	136	32.6 (26.9-38.3)
	Pouch use	8	2.5 (0.8-4.2)	9	2.5 (0.9-4.1)	9	2.6 (0.7-4.4)	11	2.6 (0.7-4.6)
Main device type used by e-cigarette users (N _{F2F} =139, N _{Tel} =241)	Disposable	47	37.3 (28.9-45.7)	92	38.5 (32.3-44.7)	55	36.8 (28.0-45.5)	125	42.0 (34.8-49.2)
	Rechargeable with pre-filled cartridges	17	13.5 (7.5-19.5)	19	7.9 (4.5-11.3)	17	11.5 (6.0-17.1)	27	9.1 (4.8-13.4)
	Rechargeable with tank to refill	38	30.2 (22.2-38.2)	96	40.2 (34.0-46.4)	48	32.0 (23.3-40.6)	107	36.0 (29.3-42.8)
	Mod system	24	19.0 (12.1-25.9)	26	10.9 (7.0-14.8)	30	19.8 (12.3-27.2)	33	11.2 (6.6-15.9)
	Don't know	0	0 (0-0)	6	2.5 (0.5-4.5)	0	0 (0-0)	5	1.7 (0.2-3.1)

Abbreviations: NRT, nicotine replacement therapy; HTP, heated tobacco product.

Table S7: Estimates for alcohol use from face-to-face (F2F) and telephone (Tel) data collected in 2024 (F2F: Jan-Mar; Tel: Feb; N_{F2F}=2317 and N_{Tel}=2375).

Alcohol use		Unweighted				Weighted			
		F2F, n	F2F, % (95% CI)	Tel, n	Tel, % (95% CI)	F2F, n	F2F, % (95% CI)	Tel, n	Tel, % (95% CI)
AUDIT 1	Never	775	33.4 (31.5-35.3)	609	25.6 (23.8-27.4)	733	31.6 (29.7-33.6)	628	26.4 (24.4-28.5)
	Monthly or less	522	22.5 (20.8-24.2)	483	20.3 (18.7-21.9)	542	23.4 (21.5-25.2)	521	21.9 (20.0-23.9)
	2-4 per month	408	17.6 (16.0-19.2)	433	18.2 (16.6-19.8)	436	18.8 (17.1-20.5)	435	18.3 (16.5-20.1)
	2-3 per week	390	16.8 (15.3-18.3)	530	22.3 (20.6-24.0)	397	17.1 (15.5-18.8)	507	21.3 (19.5-23.2)
	4-5 per week	103	4.4 (3.6-5.2)	154	6.5 (5.5-7.5)	98	4.2 (3.4-5.1)	130	5.5 (4.5-6.5)
	6+ per week	108	4.7 (3.8-5.6)	133	5.6 (4.7-6.5)	100	4.3 (3.5-5.1)	122	5.1 (4.2-6.1)
	Don't know	2	0.1 (0.0-0.2)	15	0.6 (0.3-0.9)	2	0.1 (0.0-0.2)	14	0.6 (0.2-0.9)
	Refused	9	0.4 (0.1-0.7)	18	0.8 (0.5-1.1)	9	0.4 (0.1-0.7)	18	0.8 (0.4-1.2)
AUDIT 2 (N _{F2F} =1541, N _{Tel} =1766)	1-2	739	48.0 (45.5-50.5)	810	45.9 (43.6-48.2)	728	46 (43.3-48.6)	765	43.8 (41.1-46.4)
	3-4	446	28.9 (26.6-31.2)	466	26.4 (24.3-28.5)	471	29.8 (27.3-32.2)	462	26.4 (24.1-28.8)
	5-6	172	11.2 (9.6-12.8)	202	11.4 (9.9-12.9)	189	11.9 (10.2-13.7)	208	11.9 (10.2-13.7)
	7-9	90	5.8 (4.6-7.0)	125	7.1 (5.9-8.3)	96	6.0 (4.8-7.3)	133	7.6 (6.2-9.1)
	10-12	44	2.9 (2.1-3.7)	53	3.0 (2.2-3.8)	46	2.9 (2.0-3.8)	55	3.2 (2.2-4.2)
	13-15	17	1.1 (0.6-1.6)	18	1.0 (0.5-1.5)	17	1.1 (0.5-1.6)	27	1.5 (0.8-2.3)
	16+	17	1.1 (0.6-1.6)	33	1.9 (1.3-2.5)	20	1.2 (0.6-1.9)	33	1.9 (1.1-2.6)
	Don't know	5	0.3 (0.0-0.6)	38	2.2 (1.5-2.9)	6	0.4 (0.0-0.7)	44	2.5 (1.6-3.5)
Refused	11	0.7 (0.3-1.1)	21	1.2 (0.7-1.7)	12	0.8 (0.3-1.3)	20	1.1 (0.6-1.7)	
AUDIT 3 (N _{F2F} =1541, N _{Tel} =1766)	Never	741	48.1 (45.6-50.6)	728	41.2 (38.9-43.5)	716	45.2 (42.6-47.8)	683	39.1 (36.5-41.7)
	Less than monthly	420	27.3 (25.1-29.5)	511	28.9 (26.8-31.0)	460	29.0 (26.6-31.5)	539	30.9 (28.3-33.4)
	Monthly	195	12.7 (11.0-14.4)	266	15.1 (13.4-16.8)	215	13.6 (11.7-15.4)	267	15.3 (13.3-17.2)
	Weekly	151	9.8 (8.3-11.3)	187	10.6 (9.2-12.0)	156	9.9 (8.3-11.5)	184	10.5 (8.9-12.2)
	Daily or almost daily	19	1.2 (0.6-1.8)	33	1.9 (1.3-2.5)	20	1.2 (0.7-1.8)	36	2.1 (1.2-2.9)
	Don't know	4	0.3 (0.0-0.6)	21	1.2 (0.7-1.7)	4	0.3 (0.0-0.5)	19	1.1 (0.5-1.7)
	Refused	11	0.7 (0.3-1.1)	20	1.1 (0.6-1.6)	12	0.8 (0.3-1.3)	19	1.1 (0.5-1.6)
AUDIT-C ≥5		540	23.5 (21.8-25.2)	680	29.7 (27.8-31.6)	567	24.7 (22.8-26.6)	675	29.5 (27.3-31.6)

AUDIT 1: how often drinking; AUDIT 2: how many standard drinks on a typical drinking day; AUDIT 3: how often ≥6 standard drinks on one occasion; AUDIT-C ≥5: increasing and higher risk drinking.

4. Comparing 2024 data using telephone data from January to March 2024

This section includes face-to-face data collected between January and March 2024 and telephone data collected between January and March 2024.

Reweighting applied to telephone data:

- for January in which 21% were collected, weights were multiplied by approximately 0.27 [21%/(65%+14%)];
- for February in which 65% were collected, weights were multiplied by approximately 1.86 [65%/(21%+14%)];
- for March in which 14% were collected, weights were multiplied by approximately 0.16 [14%/(21%+65%)].

Table S8: Sociodemographic estimates from face-to-face (F2F) and telephone (Tel) data collected in Jan-Mar 2024 (N_{F2F}=2317 and N_{Tel-3-months}=7157).

Sociodemographic characteristics		Unweighted				Weighted			
		F2F, n	F2F, % (95% CI)	Tel, n	Tel, % (95% CI)	F2F, n	F2F, % (95% CI)	Tel, n	Tel, % (95% CI)
Age	16-24 years	257	11.1 (9.8-12.4)	732	10.2 (9.5-10.9)	298	12.9 (11.3-14.4)	921	12.9 (11.9-13.9)
	25-34 years	333	14.4 (13.0-15.8)	862	12.0 (11.2-12.8)	384	16.6 (14.9-18.3)	1189	16.6 (15.5-17.7)
	35-44 years	375	16.2 (14.7-17.7)	893	12.5 (11.7-13.3)	367	15.8 (14.3-17.4)	1132	15.8 (14.8-16.9)
	45-54 years	314	13.6 (12.2-15.0)	1151	16.1 (15.2-17.0)	374	16.2 (14.5-17.8)	1149	16.1 (15.1-17.0)
	55-64 years	361	15.6 (14.1-17.1)	1313	18.3 (17.4-19.2)	364	15.7 (14.1-17.2)	1121	15.7 (14.7-16.6)
	65+ years	677	29.2 (27.3-31.1)	2200	30.7 (29.6-31.8)	530	22.9 (21.2-24.5)	1639	22.9 (21.9-23.9)
	Refused	0	0 (0-0)	6	0.1 (0.0-0.2)	0	0 (0-0)	6	0.1 (0.0-0.2)
Gender	Men	1167	50.5 (48.5-52.5)	3566	50.4 (49.2-51.6)	1132	49.0 (46.9-51.2)	3462	49.0 (47.6-50.3)
	Women	1142	49.5 (47.5-51.5)	3495	49.4 (48.2-50.6)	1177	51.0 (48.8-53.1)	3599	50.9 (49.5-52.3)
	Non-binary	0	0 (0-0)	8	0.1 (0.0-0.2)	0	0 (0-0)	8	0.1 (0.0-0.2)
Social grade	AB	606	26.2 (24.4-28.0)	2142	29.9 (28.8-31.0)	610	26.3 (24.4-28.2)	1878	26.2 (25.1-27.3)
	C1	782	33.8 (31.9-35.7)	2825	39.5 (38.4-40.6)	688	29.7 (27.8-31.6)	2128	29.7 (28.6-30.8)
	C2	411	17.7 (16.1-19.3)	969	13.5 (12.7-14.3)	471	20.3 (18.5-22.1)	1449	20.3 (19.0-21.5)
	D	302	13.0 (11.6-14.4)	512	7.2 (6.6-7.8)	336	14.5 (12.9-16.1)	1036	14.5 (13.2-15.7)
	E	216	9.3 (8.1-10.5)	709	9.9 (9.2-10.6)	212	9.1 (7.9-10.3)	665	9.3 (8.6-10.0)

Table S9: Estimates for smoking and use of other nicotine products from face-to-face (F2F) and telephone (Tel) data collected in Jan-Mar 2024 (N_{F2F}=2317 and N_{Tel-3-months}=7157).

Smoking and use of other nicotine products		Unweighted				Weighted			
		F2F, n	F2F, % (95% CI)	Tel, n	Tel, % 95% CI	F2F, n	F2F, % (95% CI)	Tel, n	Tel, % (95% CI)
Tobacco use in total population	Daily cigarette smoking	212	9.1 (7.9-10.3)	628	8.8 (8.1-9.5)	230	9.9 (8.6-11.2)	750	10.5 (9.6-11.4)
	Non-daily cigarette smoking	47	2.0 (1.4-2.6)	204	2.9 (2.5-3.3)	47	2.0 (1.4-2.7)	242	3.4 (2.9-3.9)
	Non-cigarette smoking	25	1.1 (0.7-1.5)	112	1.6 (1.3-1.9)	26	1.1 (0.7-1.6)	121	1.7 (1.3-2.1)
	Stopped last year	33	1.4 (0.9-1.9)	167	2.3 (2.0-2.6)	34	1.5 (0.9-2.0)	202	2.8 (2.3-3.3)
	Stopped 1+ year ago	445	19.2 (17.6-20.8)	1771	24.7 (23.7-25.7)	423	18.2 (16.6-19.9)	1667	23.3 (22.2-24.4)
	Never smoked	1551	66.9 (65.0-68.8)	4224	59.0 (57.9-60.1)	1553	67.0 (65.0-69.1)	4119	57.5 (56.2-58.9)
	Don't know	4	0.2 (0.0-0.4)	51	0.7 (0.5-0.9)	3	0.1 (0.0-0.3)	56	0.8 (0.5-1.0)
Other nicotine use in total population	NRT use	69	3.0 (2.3-3.7)	189	2.6 (2.2-3.0)	69	3.0 (2.3-3.7)	206	2.9 (2.4-3.4)
	HTP use	2	0.1 (0.0-0.2)	12	0.2 (0.1-0.3)	2	0.1 (0.0-0.2)	13	0.2 (0.1-0.3)
	E-cigarette use	139	6.0 (5.0-7.0)	676	9.4 (8.7-10.1)	164	7.1 (5.9-8.2)	847	11.8 (10.9-12.8)
	Pouch use	32	1.4 (0.9-1.9)	39	0.5 (0.3-0.7)	27	1.2 (0.8-1.6)	49	0.7 (0.4-0.9)
Quitting/other nicotine use in people smoking in past year (N _{F2F} =317, N _{Tel-3-months} =1111)	Tried to stop in past year	113	38.0 (32.5-43.5)	362	33.9 (31.1-36.7)	121	38.0 (32.2-43.8)	434	34.3 (30.9-37.7)
	Quit in past year	33	10.4 (7.0-13.8)	167	15.0 (12.9-17.1)	34	10.0 (6.6-13.5)	202	15.4 (12.9-17.8)
	NRT use	43	13.6 (9.8-17.4)	136	12.2 (10.3-14.1)	43	12.7 (9.0-16.3)	156	11.9 (9.6-14.1)
	HTP use	1	0.3 (0.0-0.9)	8	0.7 (0.2-1.2)	1	0.4 (0.0-1.0)	8	0.6 (0.2-1.1)
	E-cigarette use	66	20.8 (16.3-25.3)	312	28.1 (25.5-30.7)	78	23.0 (18.0-28.0)	410	31.2 (27.9-34.5)
	Pouch use	8	2.5 (0.8-4.2)	28	2.5 (1.6-3.4)	9	2.6 (0.7-4.4)	36	2.7 (1.6-3.9)
Main device type used by e-cigarette users (N _{F2F} =139, N _{Tel-3-months} =676)	Disposable	47	37.3 (28.9-45.7)	241	36.0 (32.4-39.6)	55	36.8 (28.0-45.5)	324	38.5 (34.2-42.9)
	Rechargeable with pre-filled cartridges	17	13.5 (7.5-19.5)	71	10.6 (8.3-12.9)	17	11.5 (6.0-17.1)	92	10.9 (8.1-13.7)
	Rechargeable with tank to refill	38	30.2 (22.2-38.2)	261	39.0 (35.3-42.7)	48	32.0 (23.3-40.6)	308	36.7 (32.5-40.8)
	Mod system	24	19.0 (12.1-25.9)	89	13.3 (10.7-15.9)	30	19.8 (12.3-27.2)	110	13.1 (10.1-16.1)
	Don't know	0	0 (0-0)	8	1.2 (0.4-2.0)	0	0 (0-0)	7	0.9 (0.2-1.5)

Abbreviations: NRT, nicotine replacement therapy; HTP, heated tobacco product.

Table S10: Estimates for alcohol use from face-to-face (F2F) and telephone (Tel) data collected in Jan-Mar 2024 (N_{F2F}=2317 and N_{Tel-3-months}=7157).

Alcohol use		Unweighted				Weighted			
		F2F, n	F2F, % (95% CI)	Tel, n	Tel, % (95% CI)	F2F, n	F2F, % (95% CI)	Tel, n	Tel, % (95% CI)
AUDIT 1	Never	775	33.4 (31.5-35.3)	1747	24.4 (23.4-25.4)	733	31.6 (29.7-33.6)	1829	25.6 (24.4-26.8)
	Monthly or less	522	22.5 (20.8-24.2)	1534	21.4 (20.4-22.4)	542	23.4 (21.5-25.2)	1646	23.0 (21.8-24.2)
	2-4 per month	408	17.6 (16.0-19.2)	1314	18.4 (17.5-19.3)	436	18.8 (17.1-20.5)	1331	18.6 (17.5-19.7)
	2-3 per week	390	16.8 (15.3-18.3)	1597	22.3 (21.3-23.3)	397	17.1 (15.5-18.8)	1514	21.2 (20.1-22.2)
	4-5 per week	103	4.4 (3.6-5.2)	467	6.5 (5.9-7.1)	98	4.2 (3.4-5.1)	398	5.6 (5.0-6.1)
	6+ per week	108	4.7 (3.8-5.6)	420	5.9 (5.4-6.4)	100	4.3 (3.5-5.1)	363	5.1 (4.5-5.6)
	Don't know	2	0.1 (0-0.2)	41	0.6 (0.4-0.8)	2	0.1 (0.0-0.2)	40	0.6 (0.4-0.8)
	Refused	9	0.4 (0.1-0.7)	37	0.5 (0.3-0.7)	9	0.4 (0.1-0.7)	36	0.5 (0.3-0.7)
AUDIT 2 (N _{F2F} =1541, N _{Tel-3-months} = 5410)	1-2	739	48 (45.5-50.5)	2520	46.6 (45.3-47.9)	728	46 (43.3-48.6)	2357	44.2 (42.7-45.8)
	3-4	446	28.9 (26.6-31.2)	1420	26.2 (25.0-27.4)	471	29.8 (27.3-32.2)	1420	26.7 (25.3-28)
	5-6	172	11.2 (9.6-12.8)	619	11.4 (10.6-12.2)	189	11.9 (10.2-13.7)	632	11.9 (10.9-12.9)
	7-9	90	5.8 (4.6-7)	356	6.6 (5.9-7.3)	96	6.0 (4.8-7.3)	389	7.3 (6.4-8.2)
	10-12	44	2.9 (2.1-3.7)	193	3.6 (3.1-4.1)	46	2.9 (2.0-3.8)	207	3.9 (3.3-4.5)
	13-15	17	1.1 (0.6-1.6)	48	0.9 (0.7-1.1)	17	1.1 (0.5-1.6)	63	1.2 (0.8-1.6)
	16+	17	1.1 (0.6-1.6)	99	1.8 (1.4-2.2)	20	1.2 (0.6-1.9)	107	2.0 (1.5-2.5)
	Don't know	5	0.3 (0-0.6)	105	1.9 (1.5-2.3)	6	0.4 (0.0-0.7)	108	2.0 (1.6-2.5)
Refused	11	0.7 (0.3-1.1)	50	0.9 (0.6-1.2)	12	0.8 (0.3-1.3)	45	0.8 (0.6-1.1)	
AUDIT 3 (N _{F2F} =1541, N _{Tel-3-months} = 5410)	Never	741	48.1 (45.6-50.6)	2260	41.8 (40.5-43.1)	716	45.2 (42.6-47.8)	2077	39.0 (37.5-40.5)
	Less than monthly	420	27.3 (25.1-29.5)	1636	30.2 (29.0-31.4)	460	29.0 (26.6-31.5)	1730	32.5 (31.0-34.0)
	Monthly	195	12.7 (11.0-14.4)	792	14.6 (13.7-15.5)	215	13.6 (11.7-15.4)	803	15.1 (13.9-16.2)
	Weekly	151	9.8 (8.3-11.3)	531	9.8 (9.0-10.6)	156	9.9 (8.3-11.5)	529	9.9 (9.0-10.9)
	Daily or almost daily	19	1.2 (0.6-1.8)	109	2.0 (1.6-2.4)	20	1.2 (0.7-1.8)	107	2.0 (1.6-2.4)
	Don't know	4	0.3 (0.0-0.6)	44	0.8 (0.6-1.0)	4	0.3 (0.0-0.5)	45	0.8 (0.5-1.1)
	Refused	11	0.7 (0.3-1.1)	38	0.7 (0.5-0.9)	12	0.8 (0.3-1.3)	36	0.7 (0.4-0.9)
AUDIT-C ≥5		540	23.5 (21.8-25.2)	2089	30.1 (29.0-31.2)	567	24.7 (22.8-26.6)	2072	29.8 (28.6-31.1)

AUDIT 1: how often drinking; Audit 2: how many standard drinks on a typical drinking day; AUDIT 3: how often ≥6 standard drinks on one occasion; AUDIT-C ≥5: increasing and higher risk drinking.

5. Exploring differences in e-cigarette use

This section further explores differences in e-cigarette use by comparing e-cigarette use in the total population stratified by age and (given e-cigarette use was assessed within several different survey questions) by counting the number and proportion of participants who reported e-cigarette use in response to the different survey questions that contributed to the derived variable “e-cigarette use in total population”.

Table S11: Estimates for e-cigarette use in the total population stratified by age from face-to-face (F2F) and telephone (Tel) data collected in 2022 and 2024 ($N_{F2F}=4381$ and $N_{Tel}=4982$).

Age group	Unweighted				Weighted			
	F2F, n	F2F, % (95% CI)	Tel, n	Tel, % (95% CI)	F2F, n	F2F, % (95% CI)	Tel, n	Tel, % (95% CI)
16-24	59	9.8 (7.1-12.5)	99	19.5 (17.3-21.7)	58	10.1 (7.5-12.7)	141	21.5 (17.2- 25.8)
25-34	58	8.9 (8.0-9.8)	98	15.1(14.0-16.2)	65	9.0 (6.7-11.3)	148	17.8 (14.1-21.5)
35-44	41	6.0 (3.3-8.7)	78	12.4 (10.1-14.7)	46	6.7 (4.6-8.7)	101	13.0 (9.9-16.0)
45-54	40	6.7 (5.9-7.5)	57	6.5 (5.5-7.5)	52	7.2 (5.0-9.4)	53	6.6 (4.7-8.4)
55-64	40	5.7 (3.0-8.4)	44	5.1 (2.8-7.4)	39	5.7 (3.9-7.5)	38	5.0 (3.3-6.7)
65+	21	1.8 (1.1-2.5)	40	2.7 (1.8-3.6)	19	1.9 (1.1-2.8)	38	3.3 (2.1-4.4)

Table S12: Unweighted number and proportion of positive responses to each variable that contributed to the derived variable “current e-cigarette use in total population”, based on face-to-face (F2F) and telephone (Tel) data collected in 2022 and 2024 ($N_{F2F}=4381$ and $N_{Tel}=4982$). Responses not mutually exclusive.

Survey variable with e-cigarette or Juul as response option	F2F			Tel		
	N	n	% (95% CI)	N	n	% (95% CI)
Using e-cigarette to stop smoking, cut down, or any other reason ¹	702	140	19.9 (16.9-22.9)	794	150	18.9 (16.2-21.6)
Using Juul to stop smoking, cut down, or any other reason ¹	702	12	1.7 (0.7-2.7)	794	5	0.6 (0.0-1.2)
Using e-cigarette to help cut down amount smoked ²	277	82	29.6 (24.2-35.0)	362	101	27.9 (23.3-32.5)
Using Juul to help cut down amount smoked ²	277	7	2.5 (0.7-4.3)	362	6	1.7 (0.4-3.0)
Regularly using e-cigarette when not allowed to smoke ²	636	118	18.6 (15.6-21.6)	700	107	15.3 (12.6-18.0)
Regularly using Juul when not allowed to smoke ²	636	9	1.4 (0.5-2.3)	700	5	0.7 (0.1-1.3)
Using any of the following – e-cigarette ³	3667	110	3.0 (2.4-3.6)	4131	209	5.1 (4.4-5.8)
Using any of the following – Juul ³	3667	7	0.2 (0.1-0.3)	4131	2	0.0 (0.0-0.1)

¹ Asked to people who smoked in the past year

² Asked to people who currently smoke

³ Asked to people who have not smoke in the past year