

Supplementary Table 1: Recruitment group and timing for individuals that participated two or three times.

Participant	Sample #1	Sample #2	Sample #3
1	SRC (2021)	MSK (2022)	
2	SRC (2021)	SRC (2023)	
3	MSK (2021)	SRC (2024)	
4	MSK (2021)	HC (2022)	
5	SRC (2021)	SRC (2024)	
6	SRC (2021)	HC (2022)	
7	HC (2022)	SRC (2022)	
8	HC (2022)	SRC (2023)	
9	HC (2022)	SRC (2022)	
10	HC (2022)	HC (2023)	SRC (2024)
11	SRC (2022)	SRC (2022)	
12	SRC (2022)	SRC (2024)	
13	SRC (2022)	HC (2023)	SRC (2023)
14	HC (2023)	SRC (2024)	SRC (2024)
15	HC (2023)	SRC (2024)	
16	SRC (2023)	SRC (2024)	
17	HC (2023)	SRC (2024)	
18	SRC (2023)	SRC (2024)	
19	HC (2023)	SRC (2024)	

(year) denotes the recruitment year.

Abbreviations: SRC, sport-related concussion; MSK, musculoskeletal; HC, healthy control

Supplementary Table 2: Study participants that self-reported sustaining a concussion in the six months prior to recruitment.

Participant	Group	Time since previous concussion (self-reported; days)
1	SRC	29
2	SRC	9
3	SRC	29
4	HC	112
5	SRC	50
6	SRC	84

Abbreviations: SRC, sport-related concussion; HC, healthy control.

Supplementary Table 3: Area under the curve statistics for serum GFAP and SCAT symptom severity for supplementary analyses.

Time Window		SRC Sample Size	GFAP Raw	GFAP Adjusted	SCAT Symptoms	SCAT Symptoms + GFAP (Raw)
Next or Subsequent Day	i)	156	0.74 (0.68-0.80)	0.74 (0.68-0.80)	0.93 (0.89-0.96)	0.95 (0.93-0.98)
	ii)	151	0.73 (0.67-0.79)	0.74 (0.67-0.80)	0.92 (0.89-0.96)	0.95 (0.93-0.98)
	iii)	139	0.73 (0.67-0.80)	0.74 (0.68-0.80)	0.93 (0.90-0.97)	0.96 (0.94-0.99)
Next Day	i)	131	0.78 (0.72-0.84)	0.78 (0.73-0.84)	0.93 (0.90-0.96)	0.96 (0.94-0.98)
	ii)	127	0.77 (0.71-0.83)	0.78 (0.72-0.84)	0.93 (0.89-0.96)	0.96 (0.93-0.98)
	iii)	115	0.78 (0.72-0.84)	0.79 (0.73-0.85)	0.94 (0.91-0.97)	0.96 (0.94-0.99)
16-24 hours	i)	63	0.83 (0.76-0.90)	0.83 (0.76-0.90)	0.92 (0.87-0.97)	0.98 (0.95-1.00)
	ii)	60	0.83 (0.76-0.90)	0.83 (0.76-0.90)	0.91 (0.86-0.97)	0.97 (0.95-1.00)
	iii)	50	0.83 (0.75-0.91)	0.83 (0.75-0.91)	0.94 (0.90-0.99)	0.99 (0.97-1.00)
24-32 hours	i)	68	0.72 (0.64-0.81)	0.76 (0.68-0.83)	0.93 (0.89-0.97)	0.95 (0.91-0.98)
	ii)	67	0.72 (0.64-0.80)	0.75 (0.68-0.83)	0.93 (0.89-0.97)	0.94 (0.91-0.98)
	iii)	65	0.74 (0.65-0.82)	0.76 (0.69-0.84)	0.93 (0.89-0.97)	0.94 (0.90-0.98)
36-52 hours	i)	25	0.48 (0.33-0.62)	0.50 (0.35-0.66)	0.92 (0.86-0.99)	0.93 (0.87-1.00)
	ii)	24	0.48 (0.33-0.64)	0.52 (0.36-0.68)	0.92 (0.85-0.99)	0.93 (0.87-1.00)
	iii)	24	0.48 (0.33-0.63)	0.50 (0.34-0.66)	0.92 (0.85-0.99)	0.93 (0.87-1.00)

Next day defined as 16-32h, subsequent day 36-52h. Data presented as AUC (95% CI). Adjusted indicates GFAP AUC adjusted for age and BMI. i) denotes AUC analysis for all data. ii) denotes analysis with exclusion of study participants that self-reported sustaining a concussion in the 6 months prior to recruitment. iii) denotes analysis with the exclusion of the second (and third) participation of an individual. Five control participants and 1 SRC participant had no SCAT symptom data and were not included in associated AUC calculations.

Abbreviations: SRC, sport-related concussion; GFAP, glial fibrillary acidic protein; SCAT, sport concussion assessment tool. AUC, area under curve; 95% CI, 95% confidence interval; BMI, body mass index.

Supplementary Table 4: Distribution of GFAP levels for supplementary analyses.

	Sample size	GFAP Median (IQR)	Sample size	GFAP Median (IQR)
	SRC		Control	
All data	156	100.07 (63.32 – 169.22)	98	66.00 (49.46 – 81.23)
Concussion in prior six months excluded	151	100.00 (62.83 – 166.39)	97	65.87 (49.12 – 81.30)
Repeated participation excluded	139	98.81 (62.83 – 160.99)	93	66.14 (48.03 – 81.05)
	MSK		HC	
MSK vs HC	14	70.48 (49.88 – 79.87)	84	64.84 (50.15 - 81.93)

Abbreviations: SRC, sport-related concussion; GFAP, glial fibrillary acidic protein; MSK, musculoskeletal injury; HC, healthy control; IQR, interquartile range.