

1 **Pulmonary ¹²⁹Xe Magnetic Resonance Gas-exchange Abnormalities in Moderate-Severe**
2 **Asthma**

3
4 Sam Tcherner^{1,2}, Ali Mozaffaripour^{1,3}, Alexander M Matheson^{1,2}, Alexander Biancaniello², Cory
5 Yamashita⁴ and Grace Parraga¹⁻⁴

6
7 ¹Robarts Research Institute; ²Department of Medical Biophysics; ³School of Biomedical
8 Engineering; ⁴Division of Respiriology, Department of Medicine; Western University, London,
9 Canada.

10
11 **Corresponding Author**

12 Grace Parraga PhD, FCAHS Robarts Research Institute, room 5135 1151 Richmond St N,
13 Western University, London Canada, N6A 5B7, gparraga@uwo.ca, +1-519-931-5265

14
15 **Word count:1189; Table & Figure count:3; Reference count:22**

16
17 **Summary of Conflict of Interest Statements:**

18 Sam Tcherner BSc: No conflicts of interest.

19 Ali Mozaffaripour BMSc: No conflicts of interest.

20 Alexander M Matheson PhD: No conflicts of interest.

21 Alexander Biancaniello MSc: No conflicts of interest.

22 Cory Yamashita MD FRCPC: No conflicts of interest.

23 Grace Parraga PhD FCAHS: Dr Parraga reports speaking honoraria (GSK; Polarean)

24 **ABSTRACT**

25 **BACKGROUND:** Asthma is recognized as an inflammatory disease of the airways, but
26 inflammation may also affect the parenchyma and pulmonary vasculature. Hyperpolarized ^{129}Xe
27 MRI and MR spectroscopy (MRS) provide a way to quantify the transfer of gas from the airways
28 through the alveolar membrane and its binding to hemoglobin in the red blood cells (RBC) of the
29 pulmonary microvasculature. The vast majority of ^{129}Xe MRS studies have investigated
30 interstitial lung disease and the ratio of ^{129}Xe binding to the RBC and ^{129}Xe present in the
31 alveolar membrane, (RBC:membrane) which is a surrogate of oxygen gas-transfer to the blood.
32 We wondered if ^{129}Xe RBC:membrane would differ in asthma patients as compared to healthy
33 volunteers because of recent work showing abnormally diminished pulmonary vascular small-
34 vessel structure in severe- asthma.

35 **RESEARCH QUESTION:** Do ^{129}Xe MRI gas-transfer measurements differ significantly in
36 patients with moderate-severe asthma?

37 **STUDY DESIGN AND METHODS:** In this retrospective study, healthy (NCT02484885) and
38 asthma (NCT04651777; NCT02351141) participants were evaluated who provided written
39 informed consent.

40 **RESULTS:** Thirty-one participants with asthma (mean age=55 years \pm 18; 22 females) and 32
41 healthy volunteers (mean age=31 years \pm 14; 12 females) with ^{129}Xe MRS were evaluated. FEV₁,
42 VDP and DL_{CO}/K_{CO} were significantly different in asthma compared to healthy participants.
43 Age-corrected $^1\text{RBC:membrane}$ was significantly different in moderate-severe asthma
44 (0.32 \pm 0.09) as compared to healthy participants (0.47 \pm 0.12, P =.01). RBC:membrane was
45 significantly related to pulse-oximetry hemoglobin estimates (ρ =.29; P =.04) and DL_{CO} (ρ =.71;

46 $P < .001$). Significant relationships between ^{129}Xe RBC:membrane and age were observed in
47 healthy ($\rho = -.55$; $P = .002$) and asthma participants ($\rho = -0.49$; $P = .006$), adjusted for sex. A
48 significant ANCOVA model also revealed the influence of age ($P = .002$), sex ($P < .001$),
49 hemoglobin ($P = .003$) and asthma status ($P = .02$) on RBC:membrane.

50 **INTERPRETATION:** ^{129}Xe RBC:membrane values were significantly different in moderate-
51 severe asthma compared to healthy volunteers and were explained by age, sex, hemoglobin, and
52 asthma status.

53 INTRODUCTION

54 Although asthma is recognized as an inflammatory airway disease, other lung compartments
55 such as the parenchyma and pulmonary vasculature may be modified directly or indirectly, by
56 inflammation.¹⁻³ For example, evidence from cross-sectional and longitudinal studies in severe
57 asthma suggested pulmonary vascular differences in the small vessel volume fraction.^{4,5} Reversal
58 of these abnormalities was reported in asthma patients following biologic therapy, as evidenced
59 by a redistribution of blood volume from larger to smaller vessels.⁵ It is possible that some of
60 these abnormalities stem from inflammation and pulmonary vascular remodeling.^{2,3}

61 Hyperpolarized ¹²⁹Xe MRI and MR spectroscopy (MRS) provide a way to quantify the transfer
62 of gas from the airways through the alveolar membrane and its binding to hemoglobin in the red
63 blood cells (RBC) of the pulmonary microvasculature. In these studies, the ratio of ¹²⁹Xe binding
64 to the RBC to ¹²⁹Xe which has participated in transmembrane diffusion into the alveolar
65 membrane, (RBC:membrane) is utilized as a surrogate of oxygen gas-transfer to the blood.⁶
66 Recent work revealed the reproducibility and age-dependence of RBC:membrane values.⁷⁻¹⁰ The
67 vast majority of such ¹²⁹Xe “multi-compartment” studies have investigated interstitial lung
68 disease,¹¹⁻¹⁵ mainly because RBC:membrane is a sensitive marker of alveolar membrane
69 thickening and fibrosis.¹⁶ In a small number of patients with asthma, a pilot study revealed both
70 abnormally increased and abnormally decreased RBC:membrane values, without
71 RBC:membrane post-bronchodilator response.¹⁷

72 Because of these previous contradictory findings, our objective was to acquire and evaluate ¹²⁹Xe
73 MRS gas-exchange measurements in patients with moderate-severe asthma and compare these
74 directly with healthy volunteers. We also aimed to measure the quantitative relationships of

75 RBC:membrane measurements with age, sex, asthma severity, diffusing-capacity of the lung for
76 carbon-monoxide and hemoglobin values.

77

78 **METHODS**

79 *Study participants and Design*

80 We retrospectively evaluated 31 participants with asthma and 32 healthy volunteers who
81 provided written informed consent (NCT02484885; NCT04651777; NCT02351141) to
82 pulmonary function tests and MRI. Inclusion criteria for participants with asthma consisted of
83 male and female non-smokers, 18-75 years of age and documented diagnosis of asthma and
84 treated with low-to-high dose ICS/LABA. Healthy volunteers were males and females 18-85
85 years of age, with ≤ 1 pack year smoking history and no previous diagnosis or history of chronic
86 respiratory disease.

87 *Pulmonary Function Tests*

88 Spirometry and DL_{CO} measurements were undertaken according to American Thoracic Society
89 guidelines using a MedGraphic Elite Series system (MedGraphics; St. Paul, MN). Participants
90 with asthma withheld short-acting β -agonists for 6 hours, and long-acting β -agonists for 12 hours
91 prior to study visits,¹⁸ and completed the Asthma Control Questionnaire (ACQ-6),¹⁹ Asthma
92 Quality-of-Life Questionnaire (AQLQ),²⁰ and St. George's Respiratory Questionnaire (SGRQ).²¹

93 *MRI Acquisition and Analysis*

94 ¹²⁹Xe gas was polarized to 30% to 55% (XeniSpin 9820; Polarean, Durham, NC, USA).
95 Anatomic (¹H) MRI, functional (¹²⁹Xe) MRI and MRS were acquired at 3.0 Tesla (Discovery
96 MR750; GE Healthcare, Milwaukee, WI, USA), as previously described.²² Spectroscopic
97 measurements were reported as normalized ratios (RBC:membrane, Membrane:gas, RBC:gas)

98 using the area-under-the-curve values from the corresponding RBC, membrane and gas
99 resonance peaks.

100 *Statistical Analysis*

101 SPSS Statistics version 29.0 (IBM) was used for analysis. Normality was assessed with Shapiro-
102 Wilk tests, and non-parametric tests were used for non-normal data. Differences between
103 participant groups were assessed with independent samples t-tests or Mann-Whitney U-tests.
104 Analysis of covariance (ANCOVA) was used to generate significant models explaining
105 RBC:membrane values. Results were considered statistically significant if the probability of a
106 Type I error was less than 5% ($P<.05$).

107

108 **RESULTS**

109 Table 1 shows demographic, pulmonary function, and imaging measurements for 31 participants
110 with asthma (mean age, 56 years \pm 17; 21 females) and 32 healthy participants (mean age, 31
111 years \pm 14; 12 females). The two groups were significantly different for age ($P<.001$), BMI
112 ($P=.009$), FEV₁ ($P=.001$), FEV₁/FVC ($P=.004$), DL_{CO} ($P=.01$), K_{CO} ($P=.003$) and VDP
113 ($P<.001$). Mean ¹²⁹Xe RBC:membrane was significantly different when age-adjusted in
114 moderate-severe asthma (0.32 ± 0.09) as compared to the healthy volunteers (0.47 ± 0.12 ; $P=.01$).

115 Figure 1 (top panel) shows box and whisker plots with males (blue) and females (red) identified
116 the mean RBC:membrane significant difference ($P=.01$) in patients with moderate-severe
117 asthma. The middle panel shows significant sex-dependent linear relationships for
118 RBC:membrane with age in healthy ($\rho=-.55$, $P=.002$, $Y=-0.0035x+0.5776$) and asthma ($\rho=-.49$,
119 $P=.006$, $Y=-0.0022x+0.4484$) participants. A significant ANCOVA model revealed the
120 significant influence of age ($P=.005$), sex ($P<.001$), hemoglobin ($P=.008$), and asthma status

121 ($P=.02$) on RBC:membrane values. As shown in Figure 2, significant relationships for
122 RBC:membrane with hemoglobin ($\rho=.29$, $P=.04$) and DL_{CO} ($\rho=.71$, $P<.001$) were also observed.

123

124 **DISCUSSION**

125 ^{129}Xe MRI offers novel insights into airway, terminal airway and gas-transfer function in patients
126 with asthma because of its modest Ostwald solubility in biologic membranes, including the
127 alveolar and RBC membranes. Once inhaled, ^{129}Xe gas rapidly flows through airways and to
128 terminal airways throughout the lung to provide reproducible measurements of airways
129 dysfunction. ^{129}Xe also participates in alveolar transmembrane diffusion and binds to
130 microvascular RBC, displacing molecular oxygen on hemoglobin-bound heme. Because the
131 ^{129}Xe gas, alveolar membrane and RBC signals resonate at different frequencies, their individual
132 resonance peaks may be quantified directly using the area-under-the-curve (AUC) and
133 normalized to one another. The ratio of the RBC signal peak AUC to the alveolar membrane
134 signal peak AUC results in RBC:membrane, which has been proposed as a voxel-specific
135 surrogate of gas-transfer.

136 We recently discovered that bulk blood volume in the large- (BV_{10}) and small-vessels (BV_5)⁴
137 differed in severe asthma, compared to healthy volunteers, which was consistent with the notion
138 of pulmonary vascular pruning in severe asthma,⁴ hypoxic vasoconstriction and/or vascular wall
139 structural remodeling.² It is impossible to undertake serial histology to determine the temporal
140 dynamics of these processes, but ^{129}Xe MRI makes this possible non-invasively, *in vivo*, with
141 high spatial-temporal resolution, without radiation, and with obvious advantages over histology.
142 To better understand the potential influence of pulmonary vascular abnormalities on gas-
143 exchange in patients with moderate-severe asthma, here we evaluated a small group of patients
144 and healthy volunteers using ^{129}Xe MR RBC:membrane and pulmonary function tests. We

145 observed: 1) mean ^{129}Xe RBC:membrane was significantly different in asthma compared to the
146 healthy volunteers ($P=.01$), 2) RBC:membrane values were both sex- and age-dependent in
147 healthy ($P=.002$) and asthma participants ($P=.006$) participants, and, 3) in a significant model,
148 age ($P=.005$), sex ($P<.001$), hemoglobin ($P=.008$), and asthma status ($P=.02$) influenced
149 RBC:membrane values.

150 The difference between asthma and healthy volunteer RBC:membrane values, even after age-
151 correction, was unexpected. The fact that Membrane:gas did not differ but there was a trend
152 towards diminished RBC:gas values in asthma participants, suggests RBC pulmonary vascular
153 differences may be responsible for the abnormal RBC:membrane values in asthma. This
154 hypothesis is supported by the finding of abnormally diminished small vessel blood distribution
155 (coined pruning) observed in the Severe Asthma Research Program.^{4,5} Lending support to the
156 finding of lower RBC:membrane in asthma participants, a significant ANCOVA model also
157 revealed the contributions of sex, age, hemoglobin and asthma to RBC:membrane.

158 Similar to previous work, we observed an age-related decrease in the RBC:Membrane ratio.⁸⁻¹⁰
159 In addition, the finding of moderate relationships for all participants between RBC:membrane
160 and both DL_{CO} and hemoglobin is consistent with our understanding of gas-exchange and the
161 role of hemoglobin and heme binding by ^{129}Xe atoms.

162 In conclusion, we report significantly different ^{129}Xe RBC:membrane values in moderate-severe
163 asthma participants as compared to healthy volunteers, which may be related to pulmonary
164 vascular remodeling in these patients.

165 **REFERENCES**

- 166 1. Weitoft M, Andersson C, Andersson-Sjoland A, et al. Controlled and uncontrolled
167 asthma display distinct alveolar tissue matrix compositions. *Respir Res.* 2014;15(1):67.
- 168 2. Mostaco-Guidolin LB, Yang CX, Hackett TL. Pulmonary Vascular Remodeling Is an
169 Early Feature of Fatal and Nonfatal Asthma. *Am J Respir Cell Mol Biol.* 2021;65(1):114-
170 118.
- 171 3. Synn AJ, Margerie-Mellon C, Jeong SY, et al. Vascular remodeling of the small
172 pulmonary arteries and measures of vascular pruning on computed tomography. *Pulm*
173 *Circ.* 2021;11(4):20458940211061284.
- 174 4. Ash SY, Rahaghi FN, Come CE, et al. Pruning of the Pulmonary Vasculature in Asthma.
175 The Severe Asthma Research Program (SARP) Cohort. *Am J Respir Crit Care Med.*
176 2018;198(1):39-50.
- 177 5. McIntosh MJ, Matheson AM, Kooner HK, et al. Pulmonary Vascular Differences in
178 Eosinophilic Asthma after 2.5 Years of Anti-IL-5/Alpha Treatment. *Am J Respir Crit*
179 *Care Med.* 2023;208(9):998-1001.
- 180 6. Mugler JP, 3rd, Altes TA, Ruset IC, et al. Simultaneous magnetic resonance imaging of
181 ventilation distribution and gas uptake in the human lung using hyperpolarized xenon-
182 129. *Proc Natl Acad Sci U S A.* 2010;107(50):21707-21712.
- 183 7. Hahn AD, Kammerman J, Evans M, et al. Repeatability of regional pulmonary functional
184 metrics of Hyperpolarized (129) Xe dissolved-phase MRI. *J Magn Reson Imaging.*
185 2019;50(4):1182-1190.
- 186 8. Plummer JW, Willmering MM, Cleveland ZI, Towe C, Woods JC, Walkup LL.
187 Childhood to adulthood: Accounting for age dependence in healthy-reference
188 distributions in (129) Xe gas-exchange MRI. *Magn Reson Med.* 2023;89(3):1117-1133.
- 189 9. Mummy D, Zhang S, Bechtel A, et al. Functional gas exchange measures on (129)Xe
190 MRI and spectroscopy are associated with age, sex, and BMI in healthy subjects. *Front*
191 *Med (Lausanne).* 2024;11:1342499.
- 192 10. Collier GJ, Smith LJ, Saunders LC, et al. Age, sex, and lung volume dependence of
193 dissolved xenon-129 MRI gas exchange metrics. *Magn Reson Med.* 2024;92(4):1471-
194 1483.
- 195 11. Wang JM, Robertson SH, Wang Z, et al. Using hyperpolarized (129)Xe MRI to quantify
196 regional gas transfer in idiopathic pulmonary fibrosis. *Thorax.* 2018;73(1):21-28.
- 197 12. Collier GJ, Eaden JA, Hughes PJC, et al. Dissolved (129) Xe lung MRI with four-echo
198 3D radial spectroscopic imaging: Quantification of regional gas transfer in idiopathic
199 pulmonary fibrosis. *Magn Reson Med.* 2021;85(5):2622-2633.
- 200 13. Kammerman J, Hahn AD, Cadman RV, Malkus A, Mummy D, Fain SB. Transverse
201 relaxation rates of pulmonary dissolved-phase Hyperpolarized (129) Xe as a biomarker of
202 lung injury in idiopathic pulmonary fibrosis. *Magn Reson Med.* 2020;84(4):1857-1867.
- 203 14. Rankine LJ, Wang Z, Wang JM, et al. (129)Xenon Gas Exchange Magnetic Resonance
204 Imaging as a Potential Prognostic Marker for Progression of Idiopathic Pulmonary
205 Fibrosis. *Ann Am Thorac Soc.* 2020;17(1):121-125.
- 206 15. Stewart NJ, Leung G, Norquay G, et al. Experimental validation of the hyperpolarized
207 (129) Xe chemical shift saturation recovery technique in healthy volunteers and subjects
208 with interstitial lung disease. *Magn Reson Med.* 2015;74(1):196-207.

- 209 16. Kaushik SS, Freeman MS, Yoon SW, et al. Measuring diffusion limitation with a
210 perfusion-limited gas--hyperpolarized ^{129}Xe gas-transfer spectroscopy in patients with
211 idiopathic pulmonary fibrosis. *J Appl Physiol (1985)*. 2014;117(6):577-585.
- 212 17. Qing K, Teague WG, Mugler JP, Ruppert K, Mata JF, Shim YM. Reversal of airflow
213 limitation post-albuterol is not accompanied by improved xenon-129 gas exchange. *Am J*
214 *Respir Crit Care Med*. 2016:A6630-A6630.
- 215 18. Miller MR, Hankinson J, Brusasco V, et al. Standardisation of spirometry. *Eur Respir J*.
216 2005;26(2):319-338.
- 217 19. Juniper EF, O'Byrne PM, Guyatt GH, Ferrie PJ, King DR. Development and validation of
218 a questionnaire to measure asthma control. *Eur Respir J*. 1999;14(4):902-907.
- 219 20. Juniper EF, Buist AS, Cox FM, Ferrie PJ, King DR. Validation of a standardized version
220 of the Asthma Quality of Life Questionnaire. *Chest*. 1999;115(5):1265-1270.
- 221 21. Jones PW, Quirk FH, Baveystock CM, Littlejohns P. A self-complete measure of health
222 status for chronic airflow limitation. The St. George's Respiratory Questionnaire. *Am Rev*
223 *Respir Dis*. 1992;145(6):1321-1327.
- 224 22. Matheson AM, McIntosh MJ, Kooner HK, et al. Persistent (^{129}Xe) MRI Pulmonary and
225 CT Vascular Abnormalities in Symptomatic Individuals with Post-acute COVID-19
226 Syndrome. *Radiology*. 2022;305(2):466-476.

227

228 **TABLES:**

229 **Table 1. Demographic and Clinical Characteristics**

Parameter Mean (SD)	Healthy (n=32)	Asthma (n=31)	P
Age years	31 (14)	56 (17)	<.001
BMI kg/m ²	25 (4)	29 (7)	.009
Female n (%)	12 (38)	21 (68)	
GINA-4 n (%)	–	25 (81)	
GINA-5 n (%)	–	6 (19)	
Pack-years			
Mean	0 (0)	7 (14)	<.001
Median [IQR]	0 [0–0]	0 [0–7]	
Duration of asthma years			
Mean	0 (0)	28 (20)	<.001
Median [IQR]	0 [0–0]	26 [10–39]	
SpHb g/dL	15.3 (1.4)	14.8 (1.6)	.2
SpO ₂ %	98 (1)	97 (2)	.01
FEV ₁ % _{pred}	92 (13)	79 (17)	.001
FVC % _{pred}	97 (14)	91 (12)	.09
FEV ₁ /FVC % _{pred}	95 (8)	85 (13)	.004
DL _{CO} mL/min/mmHg	34 (9)	23 (9)	.003
DL _{CO} % _{pred}	121 (19)	96 (28)	.01
VA (L)	5.8 (1.4)	5.1 (1.6)	.2
VA % _{pred}	103 (14)	96 (18)	.3
K _{CO} mL/min/mmHg/L	5.8 (0.9)	4.6 (1.3)	.004
K _{CO} % _{pred}	119 (18)	99 (17)	.003
ACQ-6	–	1.5 (1.0)	
AQLQ	–	5.3 (1.0)	
SGRQ	–	37 (21)	
VDP %	1 (1)	8 (8)	<.001
¹²⁹ Xe RBC:Membrane	0.47 (0.12)	0.32 (0.09)	.01*
¹²⁹ Xe Membrane:Gas	0.99 (0.26)	0.89 (0.36)	.5*
¹²⁹ Xe RBC:Gas	0.47 (0.19)	0.30 (0.16)	.08*

230 n=11 for Asthma DL_{CO}, VA, K_{CO}; n=12 for Healthy VDP; n=23 for Asthma SpHb; n=26 for
 231 Healthy SpHb; n=27 for Asthma SGRQ; n=28 for Healthy DL_{CO}, VA, K_{CO}.

232 P=Significance values for independent samples t-test; P*=Significance values for ANCOVA
 233 adjusted for age. BMI=body mass index; GINA=Global Initiative for Asthma; IQR=interquartile
 234 range; SpHb=continuous total hemoglobin; SpO₂=peripheral capillary oxygen saturation;
 235 FEV₁=forced expiratory volume in 1 second; %_{pred}=percent of predicted value; FVC=forced
 236 vital capacity; DL_{CO}=diffusing capacity of the lungs for carbon monoxide; VA=alveolar volume;
 237 K_{CO}=transfer coefficient of the lung for carbon monoxide; ACQ-6=Asthma Control
 238 Questionnaire; AQLQ=Asthma Quality-of-Life Questionnaire.

239

240 **FIGURES:**



241 **Figure 1. ¹²⁹Xe MRI RBC:membrane in Asthma and relationships with age & sex**

242 Box and whisker plots show there was significantly different RBC:Membrane ($P=.01$), but not
243 Membrane:Gas ($P=.5$) or RBC:Gas ($P=.08$) in healthy and asthma participants. Box represents
244 mean, whiskers represent min-max range. Significant relationship with age for RBC:membrane
245 for healthy ($\rho=-.55$, $P=.002$, $Y=-0.0035x+0.5776$) and asthma participants for RBC:Membrane
246 ($\rho=-.49$, $P=.006$, $Y=-0.0022x+0.4484$). P =Significance values for ANCOVA adjusted for sex.
247 ANCOVA model shows significant effects on RBC:membrane of age ($P=.005$), sex ($P<.001$),
248 hemoglobin ($P=.008$), and asthma status ($P=.02$). The model's explanatory power is high
249 ($R^2=.964$; $F=297.747$; $P<.001$). *BD=bronchodilator; RBC=red blood cell.*

250
251 **Figure 2. RBC:membrane Relationships.**

252 Significant relationships for RBC:membrane with pulse-oximetry measured hemoglobin ($\rho=.29$,
253 $P=.04$, $y=4.00x+13.48$) and DL_{CO} ($\rho=.71$, $P<.001$, $y=56.06x+6.50$). P =Significance values.
254 *RBC=red blood cell; DL_{CO} =diffusing capacity of the lungs for carbon monoxide.*

255
256



ANCOVA Model ($R^2 = .964$; $F = 297.747$; $P < .001$)

Variable (N=64)	F-value	Sig. P	Partial Eta Squared
Age y	8.756	.005	0.163
Sex	21.317	<.001	0.321
Hemoglobin g/dL	7.590	.008	0.144
Asthma Status	5.738	.02	0.113

