

## Mapping the genetic landscape of disorders on the autoimmune-autoinflammatory continuum: potential implications for classification and treatment

Vera Fominykh<sup>1\*</sup>, Alexey A. Shadrin<sup>1,2\*\*</sup>, Piotr Jaholkowski<sup>1\*\*</sup>, Julian Fuhrer<sup>1</sup>, Nadine Parker<sup>1</sup>, Erik D. Wiström<sup>3</sup>, Oleksandr Frei<sup>1,3</sup>, Olav B. Smeland<sup>1,3</sup>, Helga Sanner<sup>4,5</sup>, Srdjan Djurovic<sup>1,2,6</sup>, Ole A. Andreassen<sup>1,2,3</sup>

\*Corresponding author vera.fominykh@medisin.uio.no

\*\* contributed equally to this manuscript

### Affiliations:

1 Centre for Precision Psychiatry, Institute of Clinical Medicine, University of Oslo, Oslo, Norway

2 K.G. Jebsen Centre for Neurodevelopmental Disorders, University of Oslo and Oslo University Hospital, Oslo, Norway

3 Section for Precision Psychiatry, Oslo University Hospital, Oslo, Norway

4 Department of Rheumatology, Oslo University Hospital, Oslo, Norway

5 Oslo New University College, Oslo, Norway

6 Department of Medical Genetics, Oslo University Hospital, Oslo, Norway

### Abstract:

#### Abstract:

**Objectives:** Based on clinical, biomarker and genetic data, autoimmune and autoinflammatory disorders (AIDs) can be classified as a disease continuum from pure autoinflammatory to pure autoimmune with mixed diseases in between. However, the genetic architecture of AIDs has not been systematically described. Here we investigate the polygenic continuum of AIDs using genome-wide association studies (GWAS) and statistical genetics methods.

**Methods:** We mapped the genetic landscape of 15 AIDs using GWAS summary statistics and methods including genomic structural equation modelling (genomic SEM), linkage disequilibrium score regression, Local Analysis of [co]Variant Association, and Gaussian causal mixture modelling (MiXeR). We performed enrichment analyses of tissues and biological gene-sets using MAGMA.

**Results:** Genomic SEM suggested a continuum structure with four underlying latent factors from autoimmune diseases at one end to autoinflammatory on the opposite end. Across AIDs, we observed a balanced mixture of negative and positive local correlations within the major histocompatibility complex, while outside this region they were predominantly positive. MiXeR analysis showed large genetic overlap in accordance with the continuum landscape. MAGMA analysis implicated genes associated with monogenic immune diseases in autoimmune (factor 1) and autoinflammatory diseases (factor 4).

**NOTE:** This preprint reports new research that has not been certified by peer review and should not be used to guide clinical practice.

**Conclusions** Our findings support a polygenic continuum across AIDs, with four genetic clusters. The "polygenic autoimmune" and "polygenic autoinflammatory" clusters reside on margins of this continuum. The identified genetic patterns across different AIDs can potentially guide drug selection, as patients within the same clusters may benefit from the same therapies.

**Keywords:** immune, autoimmune, autoinflammatory diseases, genome-wide association studies, polygenicity, classification

## Introduction

During the last twenty years, autoimmune and autoinflammatory disorders (AIDs) have entered the frontline of the clinical field and pharmaceutical development in neurology and rheumatology [Mukai et al., 2023, Mueller et al., 2023]. Various new diseases have been described, including autoimmune encephalitis [Dalmau et al., 2018], myelin oligodendrocyte glycoprotein-associated diseases, monogenic autoinflammatory diseases (interferonopathies, adenosine deaminase 2 deficiency) and others [Canna et al., 2015, Di Donato et al., 2021, de Jesus et al., 2023]. The global rise in AIDs, as highlighted by epidemiological studies [Conrad et al., 2023, Miller et al, 2023], has sparked a need for urgent innovation in the development of curation approaches and treatment strategies.

In addition, the release of new therapeutic options for neuromyelitis optica [Held et al., 2021], myasthenia gravis [DeHart-McCoyle et al., 2022] and monogenic immune diseases [Perez et al., 2022] bring new attention of clinicians to immune disease conception, classification and genetics. The current understanding of AIDs relies on previous immunological concepts of autoinflammation and autoimmunity, which define autoinflammation as a dysregulated activation of innate immune cells, driven by an imbalance in the axis of pro-inflammatory cytokines, which leads to damage of host tissues without a break in immune tolerance [Savic et al., 2020, Szekanecz et al., 2021]. Conversely, autoimmunity is characterised by the loss of immune tolerance, the recognition of self-antigens and the activation of T cells and B cells, followed by the production of specific autoantibodies and the damage of multiple organs owing to a dysregulated adaptive immune response [McGonagle et al., 2006, Matzinger et al., 2022, Shirafkan et al, 2024]. Regarding the fact that AIDs is spread between different medical specialities (neurology, rheumatology, endocrinology, gastroenterology etc) the new methods and data driven approach can help with improving classification which is needed.

The conceptualization of AIDs as a continuum began with proof of concept publications which suggested the classification of immunological diseases based on clinical and laboratory data [McGonagle et al. 2006 updated by Ben-Chetrit et al, 2018 and Savic et al, 2020], and described AIDs as a continuum with 5 main classes (monogenic autoinflammatory, polygenic autoinflammatory, mixed pattern, polygenic autoimmune, monogenic autoimmune diseases) highlighting the importance of genetic factors for the classification [See Fig 1.]. This classification has become commonplace in rheumatology, where SLE represents the prototype of systemic autoimmunity with production of multiple autoantibodies. There is still no consensus on the precise classification of some diseases such as rheumatoid arthritis (RA),


juvenile idiopathic arthritis (JIA) and ankylosing spondylitis (AS) [Mauro et al., 2021], since they exhibit overlapping features of both autoimmunity and autoinflammation [Szekanecz et al., 2021]. In neurology some diseases are easier to place on the AIDs continuum (for example, myasthenia or anti-LGI1-positive autoimmune encephalitis are considered to be classical autoimmune diseases [Fichtner et al., 2020, Bink et al., 2017] but for the majority it remains unclear (for example, central nervous system vasculitis, acute encephalomyelitis, Höftberger et al., 2017, de Moraes et al., 2023]. Moreover, current efforts are based on clinical features and qualitative markers, leading to competing classifications due to different hierarchical structures of clinical terms and biomarkers [Grateu et al., 2013, Pathak et al., 2017].

AIDs have a high degree of comorbidity within families [Harroud et al., 2023, Barkhane et al., 2022] and twins [Gerussi et al., 2022] suggesting shared genetic risk factors. Additionally, the co-existence of them in one individual suggests shared genetic underpinnings across different immune pathologies [Remalante-Rayco et al., 2023, Fominykh et al., 2017]. Genetic-based disease classification has been used in a study from Japan Biobank [Sakaue et al., 2021] and a cross-disorder genetic analysis of immune disease [Demela et al., 2023, Lincoln et al., 2024, Topaloudi et al., 2023]. Nevertheless, most of the available cross-disorder genetic studies have used only a few disorders, without clearly defined selection, and focusing on a shared component but not on the characterising of separate clusters. Only a limited number of studies have systematically evaluated the AIDs continuum. A great effort was made to use the AutoCore network for the integration of a set of 186 inborn errors of immunity [Bousfiha et al., 2020] with predominant autoimmunity or autoinflammation into a comprehensive map of human immune dysregulation [Guthrie et al., 2023]. In France, the TransImmune project was started in 2015 to revisit the nosology of AIDs by combining clinical and biomarkers information [Lorenzon et al., 2018, Tchitchek et al., 2024]. State-of-art methods of statistical genetics provide an additional opportunity to address this heterogeneity between AIDs through classifying diseases into more homogeneous subgroups based on the underlying genes and pathways that drive disease.

The main aims of our study are a) to evaluate the hypothesis of AIDs continuum based on GWAS data and b) to identify genetically driven clusters of closely related disorders and their shared genetic factors using publicly available large datasets and state-of-the art methodology. Bridging genetically driven classification with existing clinical knowledge may provide a better understanding of disease-specific molecular mechanisms, guiding further investigations of AIDs and facilitating identification of potential targets for drug repurposing.

The results can help to raise awareness of the distinct pathophysiology of AIDs, and to identify potential research directions in the field of immune-mediated disorders.

**Fig. 1. The immunological disease landscape.** AIDs described as a continuum with five main classes (monogenic autoinflammatory, polygenic autoinflammatory, mixed pattern, polygenic autoimmune, monogenic autoimmune disorders) highlighting the importance of genetic factors for the classification. Methods of statistical genetics can work with polygenic disorders and can help to characterise this continuum from genetics perspectives using available GWAS data.



## 2 Materials and methods

### 2.1. Samples

We curated a collection of well-powered publicly available GWAS summary statistics which belong to neurological, rheumatological, gastroenterological, and endocrine system AIDs resulting in data on 15 immune-linked diseases (Table 1). The cut-off for GWAS inclusion was August 2023. First, we have 26 immune-linked disorders in the list (Suppl.1, Table 1) with existing GWASs and applied criteria with N effective sample size greater than 5000 [Bulik-Sullivan et al., 2015, Tylee et al., 2018] and excluded datasets with less than 200,000 SNPs overlap with the linkage disequilibrium (LD) score regression reference panel [Zheng et al., 2017]. N effective was computed with the formula  $4/(1/n \text{ cases} + 1/n \text{ controls})$  or for internal meta-analysis (described below) as the sum of N effective for contributing cohorts. This resulted in seventeen diseases. However, we had to exclude ankylosing

spondylitis and narcolepsy because only ImmunoChip data was available which didn't provide genome-wide coverage. For SLE, JIA, PS and CeD we performed in-house meta-analyses using METAL from different publicly available datasets (FinnGen R9 release and available published GWAS, the source of data mentioned in Table 1 and Suppl. 1, Table 2) in order to increase power of current available GWASes. As well we can highlight the need for more powerful GWASes in immune-mediated neurological diseases because we can keep only two among neurological diseases. All GWAS data were limited to participants of European ancestry. After data harmonisation and pre-processing of the GWAS summary statistics with PythonConvert ([https://github.com/precimed/python\\_convert/](https://github.com/precimed/python_convert/)), we conducted cross-trait analyses using a variety of analytical tools as described below.

**Table 1. Overview of the GWAS used in the study.**

	<b>Diseases</b>	<b>Abbreviation</b>	<b>N cases</b>	<b>N controls</b>	<b>N effective</b>	<b>Source of GWAS (PMID/ GWAS catalog)</b>
<b>1</b>	Autoimmune thyroiditis	AITD	30234	725172	114296	32581359
<b>2</b>	Celiac disease	CeD	6897	334824	22142	20190752, 34278373
<b>3</b>	Crohn's disease	CD	12194	34915	36151	28067908
<b>4</b>	Juvenile idiopathic arthritis	JIA	4799	294231	15670	FinnGen, 33106285
<b>5</b>	Multiple sclerosis	MS	14802	26703	38093	31604244
<b>6</b>	Myasthenia gravis	MG	1873	36370	7125	35074870
<b>7</b>	Primary biliary cholangitis	PBC	8021	16489	21584	34033851
<b>8</b>	Primary sclerosing cholangitis	PSC	2 871	12019	9270	27992413
<b>9</b>	Primary Sjogren's syndrome	SjS	3232	17481	10911	35896530
<b>10</b>	Psoriasis (including PsA)	PS	17255	693100	64320	FinnGen, 34278373, 24482804
<b>11</b>	Rheumatoid arthritis	RA	22350	74823	68838	GCST 90132223
<b>12</b>	Systemic lupus erythematosus	SLE	5595	361571	15096	FinnGen, 26502338, 29848360
<b>13</b>	Systemic sclerosis	SS	9095	17584	23978	GCST 31672989
<b>14</b>	Type 1 diabetes	T1D	18 942	501 638	73011	34012112
<b>15</b>	Ulcerative colitis	UC	12366	34915	36527	28067908

## **2.2 Analytical tools**

### **2.2.1 Genomic structural equation modelling (SEM)**

Genomic SEM [Grotzinger et al., 2019] models the multivariate genetic architecture across traits and may reveal latent factors underlying genetic correlations and clusters of correlated traits and determine how latent factors correlate with each other. We conducted exploratory and confirmatory factor analyses (EFA and CFA, respectively) of the 15 immune phenotypes. First, the multivariable extension of LD score regression (LDSC) employed in genomic SEM was used to derive a genetic covariance matrix (S) and sampling covariance matrix (V). Population prevalence was taken from the literature (Suppl.1, Table 3). Next, EFA

with promax rotation was conducted on the standardised S matrix using the R genomic SEM package (R version: R 4.3.2, <https://github.com/GenomicSEM/>, Suppl 1., Table 4). Results from the EFA were used to guide CFA for a one-, two-, three-, four- and five-factor model. CFA was performed using Genomic SEM, and correlated factors with standardised loadings  $> 0.25$  were retained for CFA. Model fit (Table 2 and Suppl. 1, Table 5) for each factor model was assessed using recommended fit indices: standardised root mean square residual (SRMR), model  $\chi^2$  statistic, Akaike Information Criterion (AIC), and Comparative Fit Index (CFI). Model fit was considered acceptable for CFI values  $\geq 0.90$  and SRMR  $< 0.1$  [for interpretation of model fit Grotzinger et al., 2019].

### **2.2.2 Linkage disequilibrium score regression (LDSC) and Gaussian causal mixture modelling (MiXeR)**

For each phenotype, we estimated the SNP-heritability using LDSC and used LDSC for establishing the genome-wide genetic correlation ( $r_g$ ) [Bulik-Sullivan et al., 2015]. We used MiXeR (univariate and bivariate, Frei et al., 2019, <https://github.com/precimed/mixer>) to estimate the number of variants influencing the trait and the number of variants shared between pairs of traits (trait-specific and shared polygenicity) as well as mean effect size across trait influencing variants (discoverability). For MiXeR analyses we excluded the MHC region as described by Frei et al. 2019 (MHC region, 26-34 Mb), due to the intricate LD structure. For assessment of model robustness, delta AIC  $> 0$  and visual evaluation of log-likelihood plots were used. Results are presented in heatmap (Fig. 3) showing the proportion of trait-specific and shared trait-influencing SNPs followed by the standard deviation across 20 independent runs, and log-likelihood plots and tables with parameters estimated by the MiXeR model (Suppl.1., Table 7 and 8). For the analysis of disease triplets, we used trivariate MiXeR [Shadrin et al., 2024, <https://codeberg.org/intercm/mix3r>] constituting a modification of bivariate MiXeR with the possibility to access overlap between 3 phenotypes simultaneously.

### **2.2.3. Local Analysis of [co]Variant Association (LAVA)**

For local  $r_g$  analyses, we used Local Analysis of [co]Variant Association (LAVA) version 1.3.8, following the protocol with the LD reference panel based on 1000 Genomes phase 3 genotype data for European samples, and the partition of the genome into 2495 regions with an average size of 1 Mb as described elsewhere [Werme J et al, 2022]. We applied Bonferroni correction to account for multiple comparisons within each pairwise analysis. The statistical tests conducted were all two-sided. Then we identified ten regions with the greatest number of significant local genetic correlations across immune traits and created network plots

displaying these associations. Only regions revealing significant estimated SNP heritability ( $p < 0.05/2495$ ) in both diseases were used to estimate local genetic correlations between the traits. Also, we described significantly correlated loci in the MHC region (26-34 Mb) across all phenotypes.

#### **2.2.4. Functional Annotation**

We followed the FUMA protocol for the identification of genomic loci and genes, as well as performing MAGMA analysis [Watanabe et al., 2017, de Leeuw et al, 2015]. We applied MAGMA to summary statistics to test for enrichment of GWAS signals in 54 tissues. The 54 gene sets were defined by gene-expression levels from 54 GTEx tissues [GTEx Consortium, 2015]. We also analysed genes revealed by MAGMA in each of the 15 diseases and common for all phenotypes, as well as unique genes for diseases assigned to marginal latent factors obtained in genomic SEM analysis. Additionally, unique gene sets were detected for each factor (we applied Bonferroni correction for each trait and then again for 15 comparisons). We used Cytoscape version 3.9.1 [Shannon P et al 2003] for pathway analysis.


### **3. Results:**

#### **3.1. Genomic SEM**

To investigate whether there is a common genetic factor underlying the 15 AIDs, we first estimated genetic correlations using the multivariate LDSC implementation in genomic SEM. Factor is a united genetic entity underlying the set of disorders and combines it to the cluster. Next, we uncovered latent factors which represent shared variance components across diseases, we modelled the genetic variance-covariance matrices across traits using genomic SEM (Fig. 2.). The model with four latent factors provides the best fit among all tested models presented in Table 2. Unfortunately, we were unable to include T1D and MS into the four-factor model as the EFA revealed that both traits had multiple, below threshold loadings across the factors (see 4 factor EFA in Suppl. 1., Table 4), leading to a decreased CFA model fit.

Factor 1 consists of rheumatic diseases with autoimmune origin (SjS, SLE, SS). Factor 2 consists of MG and CeD which are confirmed to be autoimmune according to the current autoimmune concept, but MG also shared part of genetics with Factor 3, which belongs to mixer pathology. Factor 3 consists of AITD, JIA, PS which are considered as autoimmune, autoinflammatory or mixed according to different classifications. Factor 4 consists of diseases with gastrointestinal tract with autoinflammatory origin (CD, UC and PSC). Several diseases cross-loaded on two factors, such as RA (equally belongs to F1 and F2), PBC (F1 and F3) and MG (F2 and F3). Loadings of T1D and MS were below the selected cutoff for the minimal loading (0.25) and therefore don't belong to any of these factors.





**Fig. 2. Four groups of immune-mediated diseases identified in genomic SEM analysis.**


Genomic SEM model with four latent factors representing four diseases clusters. Green represents factor 1 (F1, “autoimmune”), orange factor 2 (F2, “autoimmune-mixed”), blue factor 3 (F3, “mixed pattern”), purple factor 4 (F4, “autoinflammatory”), and diseases cross-load on two factors are shown in its own colour. The arrows connecting the latent variables with diseases are shown “factor loadings” obtained from confirmatory factor analysis. The rounded and dashed arrows on the indicators (traits) are residual variances in the genetic indicators not explained by the common factor. The dashed arrows connecting factors are covariances and give an idea about the factors’ mutual associations.

**Table 2.** genomic SEM model performance criteria: AIC — Akaike Information Criterion, CFI — Comparative Fit Index, SRMR — standardised root mean square residual,  $\chi^2$  — model chi-square, reflecting index of exact fit to observed data df and  $p_{\chi^2}$  — degrees of freedom and p-value for the model  $\chi^2$ .


<b>N factors</b>	<b><math>\chi^2</math></b>	<b>df</b>	<b><math>P\chi^2</math></b>	<b>AIC</b>	<b>CFI</b>	<b>SRMR</b>
1	1321.83	90	4.56E-218	1381.83	0.658	0.124
2	1073.7	88	3.08E-169	1137.69	0.727	0.111
3	779.35	87	9.09E-112	845.35	0.808	0.094
<b>4</b>	<b>215.59</b>	<b>56</b>	<b>1.42E-20</b>	<b>285.59</b>	<b>0.919</b>	<b>0.077</b>
5	238.37	54	5.25E-25	312.37	0.907	0.083

### 3.2. LDSC and MiXeR

We ran LDSC analysis to quantify SNP-heritability, observed scale (Fig. 3, and Suppl. 2, Fig. 1), and univariate and bivariate MiXeR for all phenotypes to characterise polygenicity and overlap between diseases. Univariate MiXeR models exhibited good model fit (Suppl 1., Table 7). As shown below, CD, SLE and PBC have a higher SNP-heritability compared to others (more than 0.45). CeD and SjS have the lowest polygenicity, and RA and AITD have the highest polygenicity (Fig. 3). Bivariate analysis with several diseases, including CeD, SjS and PSC as well as pairs of diseases including SLE-SS and RA-SS didn't fulfil model robustness criteria and therefore are not presented. Bivariate results, for those diseases with acceptable model fit, are presented in Suppl. 2, Fig. 2, and Suppl. 1, Table 8. We ran trivariate MiXeR for F1 and F4 clusters to show that diseases from the same factor have more overlap compared to another factor, which revealed SLE/SS has less overlap with UC compared with RA from the same factor (Fig. 4).



**Fig.3. Genetic architecture characteristics.** Left part: single-nucleotide polymorphism-based heritability, observed scale, for 15 immune-mediated diseases according to linkage disequilibrium score regression. Right part: the polygenicity value multiplied by 1000 according to Gaussian causal mixture modelling. The colour of dots corresponds to colour in genomic SEM figure (green factor 1, orange factor 2, blue factor 3, purple factor 4 and own colour for mixed diseases).




**Fig.4. Trivariate Gaussian causal mixture modelling.** The illustrations of overlap inside “autoimmune” factor 1 (Systemic Sclerosis, Systemic Lupus Erythematosus and Rheumatoid Arthritis), and between “autoimmune” factor 1 (Systemic Sclerosis, Systemic Lupus Erythematosus) and “autoinflammatory” factor 4 (Ulcerative Colitis). For each triad of phenotypes, for every area of the diagram, its percentage is shown with respect to the combined total area of three phenotypes (rounded to the closest integer). The colour of circles corresponds to colour in genomic SEM figure (green F1, purple F4 and beige for the Rheumatoid Arthritis).

### 3.3. LAVA across immune-related diseases

Local genetic correlation analyses complemented global genetic correlation between the immune-related phenotypes. This step allowed us to show in which immune disorders genetic correlations are restricted to specific genomic regions, and to identify the shared genetic factors located within these genomic regions.

We identified significant regional correlations between immune diseases and genomic loci, including loci that contain genes known to be implicated in immune processes. SLE, JIA and MG were excluded due to the methods failing to converge. During assessment of the pattern of intercorrelations (Suppl. 2, Fig 3) we revealed that all diseases are highly correlated. We confirmed that the local correlation and heritability were prominent in the MHC region, where strong genetic risk was shown. Among 20 regions with top significant local genetic correlation (see Suppl. 3, Figures and Tables) ten were in the MHC region (26-34 Mb) with the top 1 being in HLA DRB1 gene, top 2 in HLA DQB1 and the top 4-5 in MHC region with different genes. The pattern of local genetic correlation in the MHC region (Fig. 5) involved a mixture of concordant and discordant effects across the diseases. For example, in loci on chr 6 [32539568 - 32586784] UC was negatively correlated with PBC, PSC, PS, AITD and RA but positively correlated with MS. In comparison, fewer diseases displayed significant correlations in loci outside the MHC, but the correlations were generally strongly positive between the diseases [for example, chr 2: 191051955 - 193033982].



**Fig. 5. LAVA characteristics for MHC locus.** Correlation matrix of positive and negative significant correlation, number of loci within MHC (upper triangular part), with loci outside MHC (bottom triangular part). Numbers in cells show the number of regions with significant heritability in both traits (Bonferroni corrected), where the correlation was assessed. Colour indicates the direction of effect (Aquamarine - positive; green - negative) for mean across all correlation values ( $r_g$ ) within the regions with significant  $h^2$  for each phenotype.


### 3.4. Functional Annotation

#### 3.4.1. MAGMA shared gene results:

MAGMA analysis for all 15 summary statistics revealed whole blood and spleen to be the most common tissues for all diseases except PSC, SjS and MG, and spleen for CeD. Also, Small Intestine Terminal Ileum, Cells\_EBV-transformed\_lymphocytes, and Lung participated in 11, 8 and 7 diseases respectively (Suppl. 2, Fig.5). We did not detect specific grouping of tissues by factor.

We assessed the most frequently associated genes across traits, which for 11 diseases were (Suppl. 2, Fig 4.): *GABBR1* (gamma-aminobutyric acid type B receptor subunit 1), *ZKSCAN3* (Zinc Finger With KRAB And SCAN Domains 3), *PGBD1* (PiggyBac Transposable Element Derived 1) and for 10 diseases: *ZSCAN31* (Zinc Finger And SCAN Domain

Containing 31), *SCAND3* (SCAN Domain Containing 3) and *ZSCAN23* (Zinc Finger And SCAN Domain Containing 23). We also revealed genes only included in factor 1 or factor 4 (the most margin groups of autoimmune and autoinflammatory phenotypes, Suppl. 1, Table 9). In factor 1, 22 genes were unique and in factor 4, 159 genes were unique with some of them involved in monogenic immune/autoinflammatory diseases (see Fig. 6. below, Suppl. 2., Fig 6. and Discussion section)



**Fig. 6.** Genes uniquely involved in “autoimmune” factor 1. We marked genes linked to monogenic immune diseases with light blue circles. Lines between genes show the involvement in molecular networks (the stringApp for Cytoscape)

Subsequently, we assessed gene sets and revealed the uniqueness of each factor. There were no gene sets specific for factor 2 because factor 2 diseases share common genes with other factors. For factor 1, 9 unique gene sets were revealed (N02-mediated IL12 pathways, PID\_TCR\_pathway, as well leukocytes and lymphocytes signalling), for factor 3 – 16 unique

gene sets, for factor 4 – 41, and for out-models MS and T1D – 19 gene sets (Suppl. 1., Table 10 and discussion section).

### **Discussion:**

In this study of the genetic architecture of diseases on the AIDs continuum, we observe a striking convergence of findings across different statistical genetics methods, which indicate clustering along the AIDs continuum. Here we present novel evidence to support the recent call [AutoCore network paper, Bousfiha et al., 2020, Guthrie et al., 2023 and TransImmuno project [Lorenzon et al., 2018, Tchitchek et al., 2024] for a paradigm of AIDs nosology to incorporate genetics of polygenic diseases.

We found that by using the state-of-the-art statistical approaches, it is possible to reveal four clusters across the AIDs continuum based on GWAS data, with diseases belonging to factor 1 (autoimmune) and diseases belonging to factor 4 (autoinflammatory). The extreme positions for factor 1 and factor 4 diseases were further supported by trivariate MiXeR analysis. Significant overlap was observed within factor 1 diseases (SS, SLE, and RA), while there was a smaller overlap between factor 1 diseases (SLE, SS) and the factor 4 diseases (UC).

The classification of certain diseases such as CD, UC, and PSC is consistent with a previous study by Demela et al., 2023, and remains in the same cluster. Demela's study also utilises genomic SEM to demonstrate pathway convergence. While they suggest this factor corresponds to gastric disorders, we show that other gastroenterological phenotypes (PBC and CeD), cluster with different diseases (F1+3 and F3, respectively). Therefore, this grouping may not solely be due to involvement of the gastroenterological tract but may suggest grouping based on autoinflammatory pathogenesis. Topaloudi A. et al. [Topaloudi A. et al., 2023] also revealed four factors structure in line with a recent preprint [Breunig et al., 2024]. According to Topaloudi paper MS was in the same group as PSC and this connection in our paper was shown in LDSC and LAVA, as well as MG was in the same group as RA, and we were able to show that they share part of factor 2. Based on our findings and previous literature [Demela et al., 2023, Topaloudi et al. 2023, Williams et al., 2023] it appears that the factor structure in the genomic SEM model depends on 1) the initial set of disorders and 2) GWAS power. By including more diseases in our analyses compared to the previous studies, we were able to differentiate more factors within the autoimmune spectrum. This insight can enhance the classification of AIDs.

In the Transimmuno project [Tchitchek et al., 2024] five factors were revealed, but with inclusion of diseases with monogenic inheritance (Familial Mediterranean Fever) or

diseases without powerful GWAS (idiopathic inflammatory myopathies). Additionally, our factor 3 consists of diseases which were not studied in their project. Tchitchek et al. revealed the belonging of CD and UC to the same factor which supports our findings on biomarkers level. According to Tchitchek's classification, RA and T1D were in the same group, and we shown the significant overlap in MiXeR (rg 0.33, shared fraction 38%) and had strong local positive correlations in some loci as indicated by LAVA disregarding the fact that T1D is out of model in our study.

Also, in our study RA and SLE belong in factor one and RA in both factor 1 and 2, indicating that RA has a different genetic background compared to SLE. SLE is regarded as a classical autoimmune disease with autoantibody production, whereas RA has both autoimmune (classical seropositive RA) and mixed/autoinflammatory features (seronegative subtypes, Szekanecz et al., 2021). A notable observation is the grouping of MG together with CeD, which can help to reveal background for co-incidence of both diseases in case-reports without confirmed evidence at the population level [Thawani et al., 2018]. If we focus on MS, which were out of our genomic SEM model, but it was shown that MS has a high correlation and overlap with factor 4 diseases and PBC by other methods. That was previously shown [Kim et al., 2022] and support the fact that MS has common genetic background not only with autoimmune but also with autoinflammatory clusters.

LAVA analysis showed a mixed pattern (positive and negative) of correlations inside the MHC region in AIDs, but in loci outside MHC correlations were mostly positive. The same pattern for mixed correlation in the main MHC hotspot [chr6: 32539568 - 32586784] was shown for AIDs in the original LAVA paper [Werme et al., 2022]. Most MHC loci with significant local correlation belong to the MHC II subgroup. On the contrary PS which belongs to MHC-1-opathy according to EULAR classification [Kuiper et al., 2023, Matzaraki et al., 2017] in most of the cases has a negative correlation with other diseases in top-MHC loci. The MHC findings could be limited by non-specialized genotyping chips in GWAS and resolution of the MHC region, as well as limitations of the LAVA method due to LD structure and GWAS power.

The most frequent genes across diseases were involved in cell signalling and interaction and can be common for all types of immune disorders, consistent with Lincoln et al. (2024). Then we assessed the unique genes in marginal "autoimmune" and "autoinflammatory factors". In "autoimmune" factor, 22 genes were unique, among them neutrophil cytosolic factor 2, which is associated with autosomal recessive chronic granulomatous disease 2 with neutrophils lacking superoxide-generating act [Nunoi et al., 1998]. Mutation in Interferon Regulatory



Factor 7 was described as a cause of immunodeficiency-39 in a child [Ciancanelli, M. J. et al, 2015] and as a gene which is linked to intrinsic and innate immunity according to the classification of inborn errors of immunity [Tangye et al., 2022]. Mutation in IKZF3 was described in Immunodeficiency-84, which is an autosomal dominant primary immunologic disease associated with low levels of B cells and impaired early B-cell development [Yamashita, M. et al., 2021]. For “autoinflammatory” factor 4, 159 genes were unique, but less percentage participated in the immunological process [Suppl. 2., Fig. 6]. For example, the NOD2 gene plays a role in Blau syndrome, a rare autosomal dominant autoinflammatory syndrome classified as an autoinflammatory phenotype according to Tangye et al., 2022. These findings are in line with AutoCore's idea that monogenic AIDs may represent genetically determined, more severe forms of more common polygenic AIDs. That also enables the transfer of knowledge between rare and complex diseases [Guthrie et al., 2023].

In the assessment of gene sets, we identified the involvement of Interleukin 12 (IL12) and IL23 gene sets in factor 4 diseases. A monoclonal antibody which targets IL12/IL23 (ustekinumab) is approved for the treatment of UC and CD as well as PS [Aggeletopoulou et al., 2018, Koutruba et al., 2010]. Also, IL3, IL4, IL18, IL27, T-helper and IFN gene sets involved in “autoinflammatory” factor 4 provide valuable insights that can inform the potential application of drugs that targets these interleukins and related pathways.

For factor 1, NOD2-mediated IL12 pathways in NK cells were involved as well as the IL12 pathway, and that was shown before to be associated with SLE genetic risk [Larosa M et al., 2019]. Also, TCR signalling in naïve CD4+ T cells and leukocyte-associated pathways participated in “autoimmune” factor 1. These pathways can also be important for SjS and SS where fewer studies about the involvement of IL pathways exist due to the lack of powerful data [Shi et al., 2024].

In factor 3 diseases a lot of pathways were involved linked to T-cells, B-cells and JAK pathways as well as IL17. Involvement of IL17 in JIA pathogenesis was recently shown in different studies [Paroli et al., 2022] as well secukinumab, IL17 blocker, was approved for treating of 2 of 7 JIA subtypes (juvenile psoriatic arthritis and enthesitis-related arthritis) as well as for PS treatment. Also, for MS and T1D which were out of any factors in genomic SEM, different IL2, IL6 and T-cell pathways were important.

The accumulated evidence from this study and previous findings can help inform immunotherapy for AIDs. Already designed drugs which suppress major pro-inflammatory signalling pathways as IL-17, JAK inhibitors have tremendous success compared to traditional systemic therapies. Unfortunately, there are still unaddressed medical requirements in terms of

both long-term safety and overall effectiveness, as a considerable number of patients do not attain disease remission. Enhanced understanding of the genetically informed mechanisms and diversity within AIDs would create opportunities to address these challenges, resulting in a more personalised and efficient treatment approach.

A limitation of our study is that it only focussed on GWAS performed on populations of European ancestry. This is because genomic SEM and LD score regression require the samples to be drawn from the same ancestry, as well lack of powerful GWAS for non-European ancestry for most of the diseases. The second limitation was that we could use only powerful GWASes for polygenic disease for statistical methods available and had to ignore small GWASes as well as data freeze performed before analysis (August 2023). Also, cohort sample size and SNP-sets are drivers of the limited resolution of genetic mapping and the ability to detect robust disease associations [Uffelmann et al. 2021]. An additional limitation was to focus only on genetics but not on the deep immunophenotyping via cells/immune molecules due to the lack of powerful data for these phenotypes. Our findings about genetic clusters and overlap does not necessarily disqualify the current clinical system but bring attention to distribution of genetic factors. As well data on infections, vaccine regimen and the treatment with anti-inflammatory drugs lack in used GWAS and that is a common limitation of studies performed on summary statistics data.

To summarise, the revealing of four-factors-clustering can help to define the disease groups for which it is possible to use the same therapies. Furthermore, our results can be instrumental in assembling cluster-guided multi-parametric analyses that include genetics and omics data and enable deep phenotyping of patients leading to personalised drug selection. The current study sheds light on the autoimmune-autoinflammatory continuum from a genetic perspective and can inform future studies in this field.

## **Acknowledgements**

All GWAS investigated in the present study were approved by the local ethics committees, and informed consent was obtained from all participants. The authors thank the researchers of the IMSGC <https://imsgc.net/>, and consortia for access to data, and for all participants who provided DNA samples. We want to acknowledge the participants and investigators of the FinnGen study: <https://www.finnngen.fi/> [Kurki MI et al., 2023].

We gratefully acknowledge support from the the Research Council of Norway (#223273, 248778, 300309, 324252, 324499, 326813, 344121), European Union's Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie Actions Grant

801133 (Scientia fellowship); the South-East Norway Regional Health Authority (2022-073 and 2023-087), KG Jebsen Stiftelsen. This project has received funding from the European Union's Horizon 2020 research and innovation programme under grant agreement No 847776 and 964874, and EEA grants (EEA-RO-NO-2018-0535, EEA-RO-NO-2018-0573).

This work was partly performed on the TSD (Tjeneste for Sensitive Data) facilities, owned by the University of Oslo and operated and developed by the TSD service group at the University of Oslo, IT Department (USIT), with resources provided by UNINETT Sigma2 - the National Infrastructure for High Performance Computing and Data Storage in Norway.

Computations were also performed on resources provided by the National Infrastructure for High-Performance Computing and Data Storage in Norway.

### **Declaration of Interests**

O.A.A. has received speaker fees from Lundbeck, Janssen, Otsuka, and Sunovion and is a consultant to Cortechs.ai. and Precision Health. Dr. Frei is a consultant to Precision Health. Remaining authors have no conflicts of interest to declare.

### **Data availability and Computational tools**

GWAS data are publicly available or available on request.

Statistical analyses were performed with Python and R using existing tools available on

FUMA <https://fuma.ctglab.nl/>

GitHub <https://github.com/precimed/>

Genomic SEM <https://github.com/GenomicSEM/>

METAL <https://github.com/statgen/METAL>

MiXeR bivariate <https://github.com/precimed/mixer>

MiXeR trivariate <https://codeberg.org/intercm/mix3r>

LAVA <https://github.com/josefin-werme/LAVA>

LDSC <https://github.com/bulik/ldsc>

MAGMA <https://cncr.nl/research/magma/>

Python Convert [https://github.com/precimed/python\\_convert](https://github.com/precimed/python_convert)

## **Abbreviations:**

AD — autoimmune diseases

AIC — Akaike Information Criterion

AIDs — autoimmune and autoinflammatory disorders

AIF — autoinflammatory disorders

AITD — autoimmune thyroiditis

EFA — exploratory and confirmatory factor analysis

CeD — celiac disease

CD — Crohn's disease

CFA — confirmatory factor analysis

CFI — Comparative Fit Index

CNS — central nervous system

Genomic SEM — genomic structural equation modelling

IL — interleukin

JAK — Janus kinase

JIA — juvenile idiopathic arthritis

LAVA — Local Analysis of [co]Variant Association

LD — linkage disequilibrium

LDSC — linkage disequilibrium score regression

MHC — major histocompatibility complex

MS — multiple sclerosis

MiXeR — Gaussian causal mixture modelling

MG — myasthenia gravis

PBC — primary biliary cholangitis

PS — psoriasis

PSC — primary sclerosing cholangitis

RA — rheumatoid arthritis

SjS — primary Sjogren's syndrome

SLE — systemic lupus erythematosus

SRMR — standardised root mean square residual

SS — systemic sclerosis

T1D — type 1 diabetes

UC — ulcerative colitis

## References:

1. Mukai T, Ida H, Ueki Y and Nishikomori R (2023) Editorial: A new frontier in translational research on autoinflammatory diseases - various aspects of innate immunity on human diseases. *Front. Immunol.* 14:1147202. doi: 10.3389/fimmu.2023.1147202
2. Mueller F, Taubmann J, Voelkl S, Laura Bucci, Christina Bergmann, Michael Aigner, Artur Wilhelm, Tobias Rothe, Ioanna Minopoulou, Johannes Knitza, Soraya Kharboutli, Silvia Spoerl, Ingrid Vasova, Dagmar Werner, Hannah Reimann, Sascha Kretschmann, Dimitrios Mougiakakos, Gerhard Krönke, Georg Schett, Andreas Mackensen, CD19-Targeted CAR-T Cells in Refractory Systemic Autoimmune Diseases: A Monocentric Experience from the First Fifteen Patients, *Blood*, Volume 142, Supplement 1, 2023, Page 220, <https://doi.org/10.1182/blood-2023-180547>.
3. Dalmau J, Graus F. Antibody-Mediated Encephalitis. *N Engl J Med.* 2018 Mar 1;378(9):840-851. doi: 10.1056/NEJMra1708712.
4. Canna SW, Goldbach-Mansky R. New monogenic autoinflammatory diseases--a clinical overview. *Semin Immunopathol.* 2015 Jul;37(4):387-94. doi: 10.1007/s00281-015-0493-5.
5. Di Donato G, d'Angelo DM, Breda L, Chiarelli F. Monogenic Autoinflammatory Diseases: State of the Art and Future Perspectives. *International Journal of Molecular Sciences.* 2021; 22(12):6360. <https://doi.org/10.3390/ijms22126360>
6. de Jesus AA, Chen G, Yang D, et al., Constitutively active Lyn kinase causes a cutaneous small vessel vasculitis and liver fibrosis syndrome. *Nat Commun.* 2023 Mar 17;14(1):1502. doi: 10.1038/s41467-023-36941-y.
7. Conrad N, Misra S, Verbakel JY, Verbeke G, Molenberghs G, Taylor PN, Mason J, Sattar N, McMurray JJV, McInnes IB, Khunti K, Cambridge G. Incidence, prevalence, and co-occurrence of autoimmune disorders over time and by age, sex, and socioeconomic status: a population-based cohort study of 22 million individuals in the UK. *Lancet.* 2023 Jun 3;401(10391):1878-1890. doi: 10.1016/S0140-6736(23)00457-9.
8. Miller FW. The increasing prevalence of autoimmunity and autoimmune diseases: an urgent call to action for improved understanding, diagnosis, treatment, and prevention. *Curr Opin Immunol.* 2023 Feb;80:102266. doi: 10.1016/j.coi.2022.102266.

9. Held F, Klein AK, Berthele A. Drug Treatment of Neuromyelitis Optica Spectrum Disorders: Out with the Old, in with the New? *Immunotargets Ther.* 2021 Mar 19;10:87-101. doi: 10.2147/ITT.S287652.
10. DeHart-McCoyle M, Patel S, Du X. New and emerging treatments for myasthenia gravis. *BMJ Med.* 2023 Aug 7;2(1):e000241. doi: 10.1136/bmjmed-2022-000241.
11. Perez E. Future of Therapy for Inborn Errors of Immunity. *Clin Rev Allergy Immunol.* 2022 Aug;63(1):75-89. doi: 10.1007/s12016-021-08916-8.
12. Savic S, Caseley EA, McDermott MF. Moving towards a systems-based classification of innate immune-mediated diseases. *Nat Rev Rheumatol.* 2020 Apr;16(4):222-237. doi: 10.1038/s41584-020-0377-5.
13. Szekanecz Z, McInnes IB, Schett G, Szamosi S, Benkő S, Szűcs G. Autoinflammation and autoimmunity across rheumatic and musculoskeletal diseases. *Nat Rev Rheumatol.* 2021 Oct;17(10):585-595. doi: 10.1038/s41584-021-00652-9.
14. McGonagle D, McDermott MF. A proposed classification of the immunological diseases. *PLoS Med.* 2006 Aug;3(8):e297. doi: 10.1371/journal.pmed.0030297.
15. Matzinger P. Autoimmunity: Are we asking the right question? *Front Immunol.* 2022 Nov 3;13:864633. doi: 10.3389/fimmu.2022.864633.
16. Shirafkan F, Hensel L, Rattay K. Immune tolerance and the prevention of autoimmune diseases essentially depend on thymic tissue homeostasis. *Front Immunol.* 2024 Mar 20;15:1339714. doi: 10.3389/fimmu.2024.1339714.
17. Ben-Chetrit E, Gattorno M, Gul A, Kastner DL, Lachmann HJ, Touitou I, Ruperto N; Paediatric Rheumatology International Trials Organisation (PRINTO) and the AIDs Delphi study participants. Consensus proposal for taxonomy and definition of the autoinflammatory diseases (AIDs): a Delphi study. *Ann Rheum Dis.* 2018 Nov;77(11):1558-1565. doi: 10.1136/annrheumdis-2017-212515.
18. Mauro D, Thomas R, Guggino G, Lories R, Brown MA, Ciccia F. Ankylosing spondylitis: an autoimmune or autoinflammatory disease? *Nat Rev Rheumatol.* 2021 Jul;17(7):387-404. doi: 10.1038/s41584-021-00625-y.
19. Fichtner ML, Jiang R, Bourke A, Nowak RJ, O'Connor KC. Autoimmune Pathology in Myasthenia Gravis Disease Subtypes Is Governed by Divergent Mechanisms of Immunopathology. *Front Immunol.* 2020 May 27;11:776. doi: 10.3389/fimmu.2020.00776.

20. Binks SNM, Klein CJ, Waters P, Pittock SJ, Irani SR. LGI1, CASPR2 and related antibodies: a molecular evolution of the phenotypes. *J Neurol Neurosurg Psychiatry*. 2018 May;89(5):526-534. doi: 10.1136/jnnp-2017-315720.
21. de Moraes MPM, do Nascimento RRNR, Abrantes FF, Pedroso JL, Perazzio SF, Barsottini OGP. What General Neurologists Should Know about Autoinflammatory Syndromes? *Brain Sci*. 2023 Sep 21;13(9):1351. doi: 10.3390/brainsci13091351.
22. Höftberger R, Lassmann H. Inflammatory demyelinating diseases of the central nervous system. *Handb Clin Neurol*. 2017;145:263-283. doi: 10.1016/B978-0-12-802395-2.00019-5.
23. Grateau G, Hentgen V, Stojanovic KS, Jéru I, Amselem S, Steichen O. How should we approach classification of autoinflammatory diseases? *Nat Rev Rheumatol*. 2013 Oct;9(10):624-9. doi: 10.1038/nrrheum.2013.101.
24. Pathak S, McDermott MF, Savic S. Autoinflammatory diseases: update on classification diagnosis and management. *J Clin Pathol*. 2017 Jan;70(1):1-8. doi: 10.1136/jclinpath-2016-203810.
25. Harroud A, Hafler DA. Common genetic factors among autoimmune diseases. *Science*. 2023 May 5;380(6644):485-490. doi: 10.1126/science.adg2992.
26. Barkhane Z, Elmadi J, Satish Kumar L, Pugalenti LS, Ahmad M, Reddy S. Multiple Sclerosis and Autoimmunity: A Veiled Relationship. *Cureus*. 2022 Apr 19;14(4):e24294. doi: 10.7759/cureus.24294.
27. Gerussi A, Soskic B, Asselta R, Invernizzi P, Gershwin ME. GWAS and autoimmunity: What have we learned and what next. *J Autoimmun*. 2022 Dec;133:102922. doi: 10.1016/j.jaut.2022.102922.
28. Remalante-Rayco P, Espiritu AI, Daghistani Y, Chim T, Atenafu E, Keshavarzi S, Jha M, Gladman DD, Oh J, Haroon N, Inman RD. Incidence and predictors of demyelinating disease in spondyloarthritis: data from a longitudinal cohort study. *Rheumatology (Oxford)*. 2023 Oct 4;kead527. doi: 10.1093/rheumatology/kead527.
29. Fominykh V, Shevtsova T, Arzumanian N, Brylev L. Coexistence of multiple sclerosis and ankylosing spondylitis: Report of four cases from Russia and review of the literature. *J Clin Neurosci*. 2017 Oct;44:230-233. doi: 10.1016/j.jocn.2017.06.031.
30. Sakaue S, Kanai M, Tanigawa Y, Karjalainen J, Kurki M, Koshiha S, Narita A, Konuma T, Yamamoto K, Akiyama M, Ishigaki K, Suzuki A, Suzuki K, Obara W, Yamaji K, Takahashi K, Asai S, Takahashi Y, Suzuki T, Shinozaki N, Yamaguchi H,



- Minami S, Murayama S, Yoshimori K, Nagayama S, Obata D, Higashiyama M, Masumoto A, Koretsune Y; FinnGen; Ito K, Terao C, Yamauchi T, Komuro I, Kadowaki T, Tamiya G, Yamamoto M, Nakamura Y, Kubo M, Murakami Y, Yamamoto K, Kamatani Y, Palotie A, Rivas MA, Daly MJ, Matsuda K, Okada Y. A cross-population atlas of genetic associations for 220 human phenotypes. *Nat Genet*. 2021 Oct;53(10):1415-1424. doi: 10.1038/s41588-021-00931-x.
31. Demela P, Pirastu N, Soskic B. Cross-disorder genetic analysis of immune diseases reveals distinct gene associations that converge on common pathways. *Nat Commun*. 2023 May 12;14(1):2743. doi: 10.1038/s41467-023-38389-6.
32. Lincoln, M.R., Connally, N., Axisa, PP. *et al*. Genetic mapping across autoimmune diseases reveals shared associations and mechanisms. *Nat Genet* 56, 838–845 (2024). <https://doi.org/10.1038/s41588-024-01732-8>
33. Topaloudi A, Jain P, Martinez MB, Bryant JK, Reynolds G, Zagoriti Z, Lagoumintzis G, Zamba-Papanicolaou E, Tzartos J, Poulas K, Kleopa KA, Tzartos S, Georgitsi M, Drineas P, Paschou P. PheWAS and cross-disorder analysis reveal genetic architecture, pleiotropic loci and phenotypic correlations across 11 autoimmune disorders. *Front Immunol*. 2023 Sep 21;14:1147573. doi: 10.3389/fimmu.2023.1147573.
34. Bousfiha A, Jeddane L, Picard C, Al-Herz W, Ailal F, Chatila T, Cunningham-Rundles C, Etzioni A, Franco JL, Holland SM, Klein C, Morio T, Ochs HD, Oksenhendler E, Puck J, Torgerson TR, Casanova JL, Sullivan KE, Tangye SG. Human Inborn Errors of Immunity: 2019 Update of the IUIS Phenotypical Classification. *J Clin Immunol*. 2020 Jan;40(1):66-81. doi: 10.1007/s10875-020-00758-x.
35. Guthrie J, Köstel Bal S, Lombardo SD, Müller F, Sin C, Hütter CVR, Menche J, Boztug K. AutoCore: A network-based definition of the core module of human autoimmunity and autoinflammation. *Sci Adv*. 2023 Sep;9(35):eadg6375. doi: 10.1126/sciadv.adg6375.
36. Lorenzon R, Mariotti-Ferrandiz E, Aheng C, Ribet C, Toumi F, Pitoiset F, Chaara W, Derian N, Johanet C, Drakos I, Harris S, Amselem S, Berenbaum F, Benveniste O, Bodaghi B, Cacoub P, Grateau G, Amouyal C, Hartemann A, Saadoun D, Sellam J, Seksik P, Sokol H, Salem JE, Vicaut E, Six A, Rosenzweig M, Bernard C, Klatzmann D. Clinical and multi-omics cross-phenotyping of patients with autoimmune and autoinflammatory diseases: the observational TRANSIMMUNOM protocol. *BMJ Open*. 2018 Aug 30;8(8):e021037. doi: 10.1136/bmjopen-2017-021037.

37. Tchitchek N, Binivignat M, Roux A, et al Deep immunophenotyping reveals that autoimmune and autoinflammatory disorders are spread along two immunological axes capturing disease inflammation levels and types *Annals of the Rheumatic Diseases* Published Online First: 05 January 2024. doi: 10.1136/ard-2023-225179
38. Bulik-Sullivan BK, Loh PR, Finucane HK, Ripke S, Yang J; Schizophrenia Working Group of the Psychiatric Genomics Consortium; Patterson N, Daly MJ, Price AL, Neale BM. LD Score regression distinguishes confounding from polygenicity in genome-wide association studies. *Nat Genet.* 2015 Mar;47(3):291-5. doi: 10.1038/ng.3211.
39. Tylee DS, Sun J, Hess JL, Tahir MA, Sharma E, Malik R, Worrall BB, Levine AJ, Martinson JJ, Nejentsev S, Speed D, Fischer A, Mick E, Walker BR, Crawford A, Grant SFA, Polychronakos C, Bradfield JP, Sleiman PMA, Hakonarson H, Ellinghaus E, Elder JT, Tsoi LC, Trembath RC, Barker JN, Franke A, Dehghan A; 23 and Me Research Team; Inflammation Working Group of the CHARGE Consortium; METASTROKE Consortium of the International Stroke Genetics Consortium; Netherlands Twin Registry; neuroCHARGE Working Group; Obsessive Compulsive and Tourette Syndrome Working Group of the Psychiatric Genomics Consortium; Faraone SV, Glatt SJ. Genetic correlations among psychiatric and immune-related phenotypes based on genome-wide association data. *Am J Med Genet B Neuropsychiatr Genet.* 2018 Oct;177(7):641-657. doi: 10.1002/ajmg.b.32652.
40. Zheng J, Erzurumluoglu AM, Elsworth BL, Kemp JP, Howe L, Haycock PC, Hemani G, Tansey K, Laurin C; Early Genetics and Lifecourse Epidemiology (EAGLE) Eczema Consortium; Pourcain BS, Warrington NM, Finucane HK, Price AL, Bulik-Sullivan BK, Anttila V, Paternoster L, Gaunt TR, Evans DM, Neale BM. LD Hub: a centralized database and web interface to perform LD score regression that maximizes the potential of summary level GWAS data for SNP heritability and genetic correlation analysis. *Bioinformatics.* 2017 Jan 15;33(2):272-279. doi: 10.1093/bioinformatics/btw613.
41. Saevarsdottir S, Olafsdottir TA, Ivarsdottir EV, Halldorsson GH, Gunnarsdottir K, Sigurdsson A, Johannesson A, Sigurdsson JK, Juliusdottir T, Lund SH, Arnthorsson AO, Styrnisdottir EL, Gudmundsson J, Grondal GM, Steinsson K, Alfredsson L, Askling J, Benediktsson R, Bjarnason R, Geirsson AJ, Gudbjornsson B, Gudjonsson H, Hjaltason H, Hreidarsson AB, Klareskog L, Kockum I, Kristjansdottir H, Love TJ, Ludviksson BR, Olsson T, Onundarson PT, Orvar KB, Padyukov L, Sigurgeirsson B, Tragante V, Bjarnadottir K, Rafnar T, Masson G, Sulem P, Gudbjartsson DF, Melsted

- P, Thorleifsson G, Norddahl GL, Thorsteinsdottir U, Jonsdottir I, Stefansson K. FLT3 stop mutation increases FLT3 ligand level and risk of autoimmune thyroid disease. *Nature*. 2020 Aug;584(7822):619-623. doi: 10.1038/s41586-020-2436-0.
42. Dubois PC, Trynka G, Franke L, Hunt KA, Romanos J, Curtotti A, Zhernakova A, Heap GA, Adány R, Aromaa A, Bardella MT, van den Berg LH, Bockett NA, de la Concha EG, Dema B, Fehrmann RS, Fernández-Arquero M, Fiatal S, Grandone E, Green PM, Groen HJ, Gwilliam R, Houwen RH, Hunt SE, Kaukinen K, Kelleher D, Korponay-Szabo I, Kurppa K, MacMathuna P, Mäki M, Mazzilli MC, McCann OT, Mearin ML, Mein CA, Mirza MM, Mistry V, Mora B, Morley KI, Mulder CJ, Murray JA, Núñez C, Oosterom E, Ophoff RA, Polanco I, Peltonen L, Platteel M, Rybak A, Salomaa V, Schweizer JJ, Sperandeo MP, Tack GJ, Turner G, Veldink JH, Verbeek WH, Weersma RK, Wolters VM, Urcelay E, Cukrowska B, Greco L, Neuhausen SL, McManus R, Barisani D, Deloukas P, Barrett JC, Saavalainen P, Wijmenga C, van Heel DA. Multiple common variants for celiac disease influencing immune gene expression. *Nat Genet*. 2010 Apr;42(4):295-302. doi: 10.1038/ng.543.
43. Glanville KP, Coleman JRI, O'Reilly PF, Galloway J, Lewis CM. Investigating Pleiotropy Between Depression and Autoimmune Diseases Using the UK Biobank. *Biol Psychiatry Glob Open Sci*. 2021 Jun;1(1):48-58. doi: 10.1016/j.bpsgos.2021.03.002.
44. de Lange KM, Moutsianas L, Lee JC, Lamb CA, Luo Y, Kennedy NA, Jostins L, Rice DL, Gutierrez-Achury J, Ji SG, Heap G, Nimmo ER, Edwards C, Henderson P, Mowat C, Sanderson J, Satsangi J, Simmons A, Wilson DC, Tremelling M, Hart A, Mathew CG, Newman WG, Parkes M, Lees CW, Uhlig H, Hawkey C, Prescott NJ, Ahmad T, Mansfield JC, Anderson CA, Barrett JC. Genome-wide association study implicates immune activation of multiple integrin genes in inflammatory bowel disease. *Nat Genet*. 2017 Feb;49(2):256-261. doi: 10.1038/ng.3760.
45. Kurki, M.I., Karjalainen, J., Palta, P. *et al.* FinnGen provides genetic insights from a well-phenotyped isolated population. *Nature* **613**, 508–518 (2023). <https://doi.org/10.1038/s41586-022-05473-8>
46. López-Isac E, Smith SL, Marion MC, Wood A, Sudman M, Yarwood A, Shi C, Gaddi VP, Martin P, Prahalad S, Eyre S, Orozco G, Morris AP, Langefeld CD, Thompson SD, Thomson W, Bowes J. Combined genetic analysis of juvenile idiopathic arthritis clinical subtypes identifies novel risk loci, target genes and key regulatory mechanisms. *Ann Rheum Dis*. 2021 Mar;80(3):321-328. doi: 10.1136/annrheumdis-2020-218481.

47. International Multiple Sclerosis Genetics Consortium. Multiple sclerosis genomic map implicates peripheral immune cells and microglia in susceptibility. *Science*. 2019 Sep 27;365(6460):eaav7188. doi:10.1126/science.aav7188.
48. Chia R, Saez-Atienzar S, Murphy N, Chiò A, Blauwendraat C; International Myasthenia Gravis Genomics Consortium; Roda RH, Tienari PJ, Kaminski HJ, Ricciardi R, Guida M, De Rosa A, Petrucci L, Evoli A, Provenzano C, Drachman DB, Traynor BJ. Identification of genetic risk loci and prioritization of genes and pathways for myasthenia gravis: a genome-wide association study. *Proc Natl Acad Sci U S A*. 2022 Feb 1;119(5):e2108672119. doi:10.1073/pnas.2108672119.
49. Cordell HJ, Fryett JJ, Ueno K, Darlay R, Aiba Y, Hitomi Y, Kawashima M, Nishida N, Khor SS, Gervais O, Kawai Y, Nagasaki M, Tokunaga K, Tang R, Shi Y, Li Z, Juran BD, Atkinson EJ, Gerussi A, Carbone M, Asselta R, Cheung A, de Andrade M, Baras A, Horowitz J, Ferreira MAR, Sun D, Jones DE, Flack S, Spicer A, Mulcahy VL, Byan J, Han Y, Sandford RN, Lazaridis KN, Amos CI, Hirschfield GM, Seldin MF, Invernizzi P, Siminovitch KA, Ma X, Nakamura M, Mells GF; PBC Consortia; Canadian PBC Consortium; Chinese PBC Consortium; Italian PBC Study Group; Japan-PBC-GWAS Consortium; US PBC Consortium; UK-PBC Consortium. An international genome-wide meta-analysis of primary biliary cholangitis: Novel risk loci and candidate drugs. *J Hepatol*. 2021 Sep;75(3):572-581. doi: 10.1016/j.jhep.2021.04.055. Epub 2021 May 23. Erratum in: *J Hepatol*. 2022 Feb;76(2):489. doi: 10.1016/j.jhep.2021.11.015. Erratum in: *J Hepatol*. 2023 Apr;78(4):883. doi:10.1016/j.jhep.2022.12.001.
50. Ji SG, Juran BD, Mucha S, Folseraas T, Jostins L, Melum E, Kumasaka N, Atkinson EJ, Schlicht EM, Liu JZ, Shah T, Gutierrez-Achury J, Boberg KM, Bergquist A, Vermeire S, Eksteen B, Durie PR, Farkkila M, Müller T, Schramm C, Sterneck M, Weismüller TJ, Gotthardt DN, Ellinghaus D, Braun F, Teufel A, Laudes M, Lieb W, Jacobs G, Beuers U, Weersma RK, Wijmenga C, Marschall HU, Milkiewicz P, Pares A, Kontula K, Chazouillères O, Invernizzi P, Goode E, Spiess K, Moore C, Sambrook J, Ouwehand WH, Roberts DJ, Danesh J, Floreani A, Gulamhusein AF, Eaton JE, Schreiber S, Coltescu C, Bowlus CL, Luketic VA, Odin JA, Chopra KB, Kowdley KV, Chalasani N, Manns MP, Srivastava B, Mells G, Sandford RN, Alexander G, Gaffney DJ, Chapman RW, Hirschfield GM, de Andrade M; UK-PSC Consortium; International IBD Genetics Consortium; International PSC Study Group; Rushbrook SM, Franke A, Karlsen TH, Lazaridis KN, Anderson CA. Genome-wide association study of primary

sclerosing cholangitis identifies new risk loci and quantifies the genetic relationship with inflammatory bowel disease. *Nat Genet.* 2017 Feb;49(2):269-273. doi: 10.1038/ng.3745.

51. Khatri B, Tessneer KL, Rasmussen A, Aghakhanian F, Reksten TR, Adler A, Alevizos I, Anaya JM, Aqrawi LA, Baecklund E, Brun JG, Bucher SM, Eloranta ML, Engelke F, Forsblad-d'Elia H, Glenn SB, Hammenfors D, Imgenberg-Kreuz J, Jensen JL, Johnsen SJA, Jonsson MV, Kvarnström M, Kelly JA, Li H, Mandl T, Martín J, Nocturne G, Norheim KB, Palm Ø, Skarstein K, Stolarczyk AM, Taylor KE, Teruel M, Theander E, Venuturupalli S, Wallace DJ, Grundahl KM, Hefner KS, Radfar L, Lewis DM, Stone DU, Kaufman CE, Brennan MT, Guthridge JM, James JA, Scofield RH, Gaffney PM, Criswell LA, Jonsson R, Eriksson P, Bowman SJ, Omdal R, Rönnblom L, Warner B, Rischmueller M, Witte T, Farris AD, Mariette X, Alarcon-Riquelme ME; PRECISESADS Clinical Consortium; Shiboski CH; Sjögren's International Collaborative Clinical Alliance (SICCA); Wahren-Herlenius M, Ng WF; UK Primary Sjögren's Syndrome Registry; Sivils KL, Adrianto I, Nordmark G, Lessard CJ. Genome-wide association study identifies Sjögren's risk loci with functional implications in immune and glandular cells. *Nat Commun.* 2022 Jul 27;13(1):4287. doi: 10.1038/s41467-022-30773-y.
52. Ellinghaus D, Ellinghaus E, Nair RP, Stuart PE, Esko T, Metspalu A, Debrus S, Raelson JV, Tejasvi T, Belouchi M, West SL, Barker JN, Kõks S, Kingo K, Balschun T, Palmieri O, Annese V, Gieger C, Wichmann HE, Kabesch M, Trembath RC, Mathew CG, Abecasis GR, Weidinger S, Nikolaus S, Schreiber S, Elder JT, Weichenthal M, Nothnagel M, Franke A. Combined analysis of genome-wide association studies for Crohn disease and psoriasis identifies seven shared susceptibility loci. *Am J Hum Genet.* 2012 Apr 6;90(4):636-47. doi: 10.1016/j.ajhg.2012.02.020.
53. Ishigaki K, Sakaue S, Terao C, Luo Y, Sonehara K, Yamaguchi K, Amariuta T, Too CL, Laufer VA, Scott IC, Viatte S, Takahashi M, Ohmura K, Murasawa A, Hashimoto M, Ito H, Hammoudeh M, Emadi SA, Masri BK, Halabi H, Badsha H, Uthman IW, Wu X, Lin L, Li T, Plant D, Barton A, Orozco G, Verstappen SMM, Bowes J, MacGregor AJ, Honda S, Koido M, Tomizuka K, Kamatani Y, Tanaka H, Tanaka E, Suzuki A, Maeda Y, Yamamoto K, Miyawaki S, Xie G, Zhang J, Amos CI, Keystone E, Wolbink G, van der Horst-Bruinsma I, Cui J, Liao KP, Carroll RJ, Lee HS, Bang SY, Siminovitch KA, de Vries N, Alfredsson L, Rantapää-Dahlqvist S, Karlson EW, Bae SC, Kimberly RP, Edberg JC, Mariette X, Huizinga T, Dieudé P, Schneider M, Kerick

- M, Denny JC; BioBank Japan Project; Matsuda K, Matsuo K, Mimori T, Matsuda F, Fujio K, Tanaka Y, Kumanogoh A, Traylor M, Lewis CM, Eyre S, Xu H, Saxena R, Arayssi T, Kochi Y, Ikari K, Harigai M, Gregersen PK, Yamamoto K, Louis Bridges S Jr, Padyukov L, Martin J, Klareskog L, Okada Y, Raychaudhuri S. Multi-ancestry genome-wide association analyses identify novel genetic mechanisms in rheumatoid arthritis. *Nat Genet.* 2022 Nov;54(11):1640-1651. doi: 10.1038/s41588-022-01213-w.
54. Bentham J, Morris DL, Graham DSC, Pinder CL, Tombleson P, Behrens TW, Martín J, Fairfax BP, Knight JC, Chen L, Replogle J, Syvänen AC, Rönnblom L, Graham RR, Wither JE, Rioux JD, Alarcón-Riquelme ME, Vyse TJ. Genetic association analyses implicate aberrant regulation of innate and adaptive immunity genes in the pathogenesis of systemic lupus erythematosus. *Nat Genet.* 2015 Dec;47(12):1457-1464. doi: 10.1038/ng.3434.
55. Julià A, López-Longo FJ, Pérez Venegas JJ, Bonàs-Guarch S, Olivé À, Andreu JL, Aguirre-Zamorano MÁ, Vela P, Nolla JM, de la Fuente JLM, Zea A, Pego-Reigosa JM, Freire M, Díez E, Rodríguez-Almaraz E, Carreira P, Blanco R, Taboada VM, López-Lasanta M, Corbeto ML, Mercader JM, Torrents D, Absher D, Marsal S, Fernández-Nebro A. Genome-wide association study meta-analysis identifies five new loci for systemic lupus erythematosus. *Arthritis Res Ther.* 2018 May 30;20(1):100. doi: 10.1186/s13075-018-1604-1.
56. López-Isac E, Acosta-Herrera M, Kerick M, Assassi S, Satpathy AT, Granja J, Mumbach MR, Beretta L, Simeón CP, Carreira P, Ortego-Centeno N, Castellvi I, Bossini-Castillo L, Carmona FD, Orozco G, Hunzelmann N, Distler JHW, Franke A, Lunardi C, Moroncini G, Gabrielli A, de Vries-Bouwstra J, Wijmenga C, Koeleman BPC, Nordin A, Padyukov L, Hoffmann-Vold AM, Lie B; European Scleroderma Group†; Proudman S, Stevens W, Nikpour M; Australian Scleroderma Interest Group (ASIG); Vyse T, Herrick AL, Worthington J, Denton CP, Allanore Y, Brown MA, Radstake TRDJ, Fonseca C, Chang HY, Mayes MD, Martin J. GWAS for systemic sclerosis identifies multiple risk loci and highlights fibrotic and vasculopathy pathways. *Nat Commun.* 2019 Oct 31;10(1):4955. doi: 10.1038/s41467-019-12760-y.
57. Chiou J, Geusz RJ, Okino ML, Han JY, Miller M, Melton R, Beebe E, Benaglio P, Huang S, Korgaonkar K, Heller S, Kleger A, Preissl S, Gorkin DU, Sander M, Gaulton KJ. Interpreting type 1 diabetes risk with genetics and single-cell epigenomics. *Nature.* 2021 Jun;594(7863):398-402. doi: 10.1038/s41586-021-03552-w.

58. Grotzinger AD, Rhemtulla M, de Vlaming R, Ritchie SJ, Mallard TT, Hill WD, Ip HF, Marioni RE, McIntosh AM, Deary IJ, Koellinger PD, Harden KP, Nivard MG, Tucker-Drob EM. Genomic structural equation modelling provides insights into the multivariate genetic architecture of complex traits. *Nat Hum Behav.* 2019 May;3(5):513-525. doi: 10.1038/s41562-019-0566-x.
59. Frei O, Holland D, Smeland OB, Shadrin AA, Fan CC, Maeland S, O'Connell KS, Wang Y, Djurovic S, Thompson WK, Andreassen OA, Dale AM. Bivariate causal mixture model quantifies polygenic overlap between complex traits beyond genetic correlation. *Nat Commun.* 2019 Jun 3;10(1):2417. doi: 10.1038/s41467-019-10310-0.
60. Shadrin AA, Hindley G, Hagen E, Parker N, Tesfaye M, Jaholkowski P, Rahman Z, Kutrolli G, Fominykh V, Djurovic S, Smeland OB, O'Connell KS, van der Meer D, Frei O, Andreassen OA, Dale AM. Dissecting the genetic overlap between three complex phenotypes with trivariate MiXeR. *medRxiv [Preprint].* 2024 Feb 27:2024.02.23.24303236. doi: 10.1101/2024.02.23.24303236.
62. Werme J, van der Sluis S, Posthuma D, de Leeuw CA. An integrated framework for local genetic correlation analysis. *Nat Genet.* 2022 Mar;54(3):274-282. doi: 10.1038/s41588-022-01017-y.
63. Watanabe K, Taskesen E, van Bochoven A, Posthuma D. Functional mapping and annotation of genetic associations with FUMA. *Nat Commun.* 2017 Nov 28;8(1):1826. doi: 10.1038/s41467-017-01261-5.
64. de Leeuw CA, Mooij JM, Heskes T, Posthuma D. MAGMA: generalized gene-set analysis of GWAS data. *PLoS Comput Biol.* 2015 Apr 17;11(4):e1004219. doi: 10.1371/journal.pcbi.1004219.
65. 1000 Genomes Project Consortium; Auton A, Brooks LD, Durbin RM, Garrison EP, Kang HM, Korbel JO, Marchini JL, McCarthy S, McVean GA, Abecasis GR. A global reference for human genetic variation. *Nature.* 2015 Oct 1;526(7571):68-74. doi: 10.1038/nature15393.
66. Shannon P, Markiel A, Ozier O, Baliga NS, Wang JT, Ramage D, Amin N, Schwikowski B, Ideker T. Cytoscape: a software environment for integrated models of biomolecular interaction networks. *Genome Res.* 2003 Nov;13(11):2498-504. doi: 10.1101/gr.1239303.
67. Breunig S, Lee YH, Karlson EW, Krishnan A, Lawrence JM, Schaffer LS, Grotzinger AD. Examining the Genetic Links between Clusters of Immune-mediated Diseases and

- Psychiatric Disorders. medRxiv [Preprint]. 2024 Jul 19:2024.07.18.24310651. doi: 10.1101/2024.07.18.24310651.
68. Williams, C.M., Poore, H., Tanksley, P.T. *et al.* Guidelines for Evaluating the Comparability of Down-Sampled GWAS Summary Statistics. *Behav Genet* **53**, 404–415 (2023). <https://doi.org/10.1007/s10519-023-10152-z>
69. Thawani SP, Brannagan TH, Lebwohl B, Green PHR, Ludvigsson JF. Celiac disease and risk of myasthenia gravis - nationwide population-based study. *BMC Neurol*. 2018 Mar 12;18(1):28. doi: 10.1186/s12883-018-1035-2.
70. Kim W, Patsopoulos NA. Genetics and functional genomics of multiple sclerosis. *Semin Immunopathol*. 2022 Jan;44(1):63-79. doi: 10.1007/s00281-021-00907-3. Epub 2022 Jan 12. PMID: 35022889.
71. Kuiper JJ, Prinz JC, Stratikos E, Kuśnierczyk P, Arakawa A, Springer S, Mintoff D, Padjen I, Shumnalieva R, Vural S, Kötter I, van de Sande MG, Boyvat A, de Boer JH, Bertsias G, de Vries N, Krieckaert CL, Leal I, Vidovič Valentinčič N, Tugal-Tutkun I, El Khaldi Ahanach H, Costantino F, Glatigny S, Mrazovac Zimak D, Lötscher F, Kerstens FG, Bakula M, Viera Sousa E, Böhm P, Bosman K, Kenna TJ, Powis SJ, Breban M, Gul A, Bowes J, Lories RJ, Nowatzky J, Wolbink GJ, McGonagle DG, Turkstra F; EULAR studygroup MHC-I-opathies. EULAR study group on 'MHC-I-opathy': identifying disease-overarching mechanisms across disciplines and borders. *Ann Rheum Dis*. 2023 Jul;82(7):887-896. doi: 10.1136/ard-2022-222852.
72. Matzaraki V, Kumar V, Wijmenga C, Zhernakova A. The MHC locus and genetic susceptibility to autoimmune and infectious diseases. *Genome Biol*. 2017 Apr 27;18(1):76. doi: 10.1186/s13059-017-1207-1.
73. Nuno H, Iwata M, Tatsuzawa S, Onoe Y, Shimizu S, Kanegasaki S, Matsuda I. AG dinucleotide insertion in a patient with chronic granulomatous disease lacking cytosolic 67-kD protein. *Blood*. 1995 Jul 1;86(1):329-33. PMID: 7795241.
74. Ciancanelli MJ, Huang SX, Luthra P, Garner H, Itan Y, Volpi S, Lafaille FG, Trouillet C, Schmolke M, Albrecht RA, Israelsson E, Lim HK, Casadio M, Hermesh T, Lorenzo L, Leung LW, Pedergnana V, Boisson B, Okada S, Picard C, Ringuier B, Troussier F, Chaussabel D, Abel L, Pellier I, Notarangelo LD, García-Sastre A, Basler CF, Geissmann F, Zhang SY, Snoeck HW, Casanova JL. Infectious disease. Life-threatening influenza and impaired interferon amplification in human IRF7 deficiency. *Science*. 2015 Apr 24;348(6233):448-53. doi: 10.1126/science.aaa1578.



75. Tangye SG, Al-Herz W, Bousfiha A, Cunningham-Rundles C, Franco JL, Holland SM, Klein C, Morio T, Oksenhendler E, Picard C, Puel A, Puck J, Seppänen MRJ, Somech R, Su HC, Sullivan KE, Torgerson TR, Meyts I. Human Inborn Errors of Immunity: 2022 Update on the Classification from the International Union of Immunological Societies Expert Committee. *J Clin Immunol*. 2022 Oct;42(7):1473-1507. doi: 10.1007/s10875-022-01289-3.
76. Yamashita M, Kuehn HS, Okuyama K, Okada S, Inoue Y, Mitsui N, Imai K, Takagi M, Kanegane H, Takeuchi M, Shimojo N, Tsumura M, Padhi AK, Zhang KYJ, Boisson B, Casanova JL, Ohara O, Rosenzweig SD, Taniuchi I, Morio T. A variant in human AIOLOS impairs adaptive immunity by interfering with IKAROS. *Nat Immunol*. 2021 Jul;22(7):893-903. doi: 10.1038/s41590-021-00951-z.
77. Aggeletopoulou I, Assimakopoulos SF, Konstantakis C, Triantos C. Interleukin 12/interleukin 23 pathway: Biological basis and therapeutic effect in patients with Crohn's disease. *World J Gastroenterol*. 2018 Sep 28;24(36):4093-4103. doi: 10.3748/wjg.v24.i36.4093.
78. Koutruba N, Emer J, Lebwohl M. Review of ustekinumab, an interleukin-12 and interleukin-23 inhibitor used for the treatment of plaque psoriasis. *Ther Clin Risk Manag*. 2010 Apr 15;6:123-41. doi: 10.2147/tcrm.s5599.
79. Larosa M, Zen M, Gatto M, Jesus D, Zanatta E, Iaccarino L, Inês L, Doria A. IL-12 and IL-23/Th17 axis in systemic lupus erythematosus. *Exp Biol Med (Maywood)*. 2019 Jan;244(1):42-51. doi: 10.1177/1535370218824547.
80. Shi W, Xu Y, Zhang A, Jia X, Liu S, Hu Z. Inflammatory cytokines and their potential role in Sjogren's syndrome risk: insights from a mendelian randomization study. *Adv Rheumatol*. 2024 Feb 16;64(1):14. doi: 10.1186/s42358-024-00354-2.
81. Paroli M, Spadea L, Caccavale R, Spadea L, Paroli MP, Nante N. The Role of Interleukin-17 in Juvenile Idiopathic Arthritis: From Pathogenesis to Treatment. *Medicina (Kaunas)*. 2022 Oct 28;58(11):1552. doi: 10.3390/medicina58111552.
82. Uffelmann, E., Huang, Q.Q., Munung, N.S. et al. Genome-wide association studies. *Nat Rev Methods Primers* 1, 59 (2021). <https://doi.org/10.1038/s43586-021-00056-9>