A MAE of 24-2 for Each Test Point																						
						3.	16	3.	17	3.2	21	3.4	7									
	2.99			3.07		3.00		3.02		3.15		3.25		3.44							ľ	- 4.0
			.99	2.	2.92		2.98		3.02		26	3.49		3.41		3.55						
	3.26	3.	.06 2.		93 2.		91	3.00		3.04		3.31				3.61						0.0
	3.64		3.45		18	3.03		2.86		2.68		3.19				3.71						- 3.8
		3.3		3.05		3.00		2.81		2.95		3.21		3.33		3.43						
				3.2		24 3.1		0 3.1		3.21		3.19		3.36								- 3.6
					3		52 3.4		48	8 3.4		3.5	1									0.0
														ale								
B				MA	٩Ε c	of 1	0-2	? fo	r E	ach	Te	est I	20	int					-			- <sup>-</sup> 3.5 Value Scale
							3.5	54	3.	61												lue
			3.2	22	3.2	24	3.3	38	3.	63	3.	79	3	.87								Va
	3.1	6	3.13		3.26		3.35		3.	66	6 3.		.85 3		3	3.88					ŀ	- 3.2
	3.24		3.37		3.54		3.68		3.	77 3.		.67 3		.58 3		.68						
3.83	3 3.8	7	3.89		4.08		3.99		3.	41 3.		.27 3		3.20 3		3.33 3.1		70				
3.88	8 3.7	9	3.73		3.73		3.48		3.15		2.96		2.95		3.00		3.	66			ŀ	- 3.0
	3.4	3.40		3.40		3.42		31	3.19		3.11		3.10		3.19							
	3.2	6	3.2	25	3.2	20	3.2	23	3.	29	3.	28	3	.27	3	.52						
			3.12		3.18		3.21		3.33		3.47		3.52								ŀ	- 2.8
							3.2	21	3.	37												

1

Figure S7. Heatmaps of the MAEs for each test point. (A) 24-2 test pattern, (B) 10-2 test pattern. The color scale represents the magnitude of the MAE at each test point, revealing the spatial distribution of estimation accuracy across the VF for both test patterns. We horizontally flipped the left eye data and integrated them with the right eye data. For the 24-2 pattern, MAEs are generally consistent across most VFs, with slight increases in peripheral areas beyond the OCT scan range and regions surrounding the physiological blind spot. The 10-2 pattern shows a relatively stable distribution of MAEs, with a modest increase in the central nasal area.

9 MAE = mean absolute error; VF = visual field; OCT = optical coherence tomography.