

# Development and validation of a novel clinical risk score to predict hypoxemia in children with pneumonia using the WHO PREPARE dataset

## Supplementary material

### Table of Contents

<b>Table S1:</b> Multivariable regression model for predicting SpO <sub>2</sub> <92% in the development dataset (n=10,884).....	3
<b>Table S2.</b> Components of the PREPARE SpO <sub>2</sub> <92% risk score in the development dataset (n=10,884) .....	4
<b>Table S3.</b> Performance of the PREPARE SpO <sub>2</sub> <92% risk score at each cut-off in the validation dataset (n=3,625).....	5
<b>Table S4:</b> Characteristics of studies included in the development and validation of the PREPARE hypoxemia risk score without respiratory distress .....	6
<b>Table S5.</b> Baseline characteristics of children with pneumonia (fast breathing and/or chest indrawing), pulse oximetry measurements and complete data on all candidate predictors included in the PREPARE risk score development and validation when excluding respiratory distress (n=15,592)	9
<b>Table S6:</b> Multivariable regression model for predicting hypoxemia in the development dataset without respiratory distress as a candidate predictor (n=11,731).....	10
<b>Table S7.</b> Components of the PREPARE hypoxemia risk score in the development dataset without respiratory distress as a candidate predictor (n=11,731).....	11
<b>Table S8.</b> Performance of the PREPARE hypoxemia risk score without respiratory distress at each cut-off in the validation dataset (n=3,861) .....	12
<b>Table S9:</b> Characteristics of studies included in the development and validation of the PREPARE hypoxemia risk score when including wheezing as a candidate predictor .....	13
<b>Table S10.</b> Baseline characteristics of children with pneumonia (fast breathing and/or chest indrawing), pulse oximetry measurements and complete data on all candidate predictors included in the PREPARE risk score development and validation when including wheezing as a candidate predictor (n=13,792) .....	15
<b>Table S11.</b> Multivariable regression model for predicting hypoxemia in the development dataset when including wheezing as a candidate predictor (n=13,792) .....	16
<b>Table S12.</b> Components of the PREPARE hypoxemia risk score in the development dataset when including wheezing as a candidate predictor (n=10,397) .....	17
<b>Table S13.</b> Comparison of risk score weights from the main analysis and the sensitivity analyses ....	18
<b>Figure S1.</b> Decision curve analysis of the PREPARE hypoxemia model .....	19
<b>Figure S2.</b> Receiver operating characteristic curve for the PREPARE SpO <sub>2</sub> <92% clinical prediction model for children 2–59 months of age with pneumonia .....	20
<b>Figure S3.</b> Calibration plot of the PREPARE SpO <sub>2</sub> <92% clinical prediction model in the validation dataset.....	21

<b>Figure S4</b> Receiver operating characteristic curve for the PREPARE hypoxemia clinical prediction model excluding respiratory distress as a candidate predictor for children 2–59 months of age with pneumonia .....	22
<b>Figure S5:</b> Calibration plot of the PREPARE hypoxemia clinical prediction model in the validation dataset when excluding respiratory distress as a candidate predictor.....	23
<b>Figure S6.</b> Receiver operating characteristic curve for the PREPARE hypoxemia clinical prediction model when including wheezing as a candidate predictor for children 2–59 months of age with pneumonia .....	24
<b>Figure S7:</b> Calibration plot of the PREPARE hypoxemia clinical prediction model in the validation dataset when including wheezing as a candidate predictor .....	25

**Table S1:** Multivariable regression model for predicting SpO2<92% in the development dataset (n=10,884)

Characteristic	SpO2 92-100% (n=9,117)	SpO2 <92% (n=1,767)	Odds ratio (95% CI)	Adjusted OR (95% CI)
<b>Age</b>				
2-5 months	2,787 (30.6%)	613 (34.7%)	1.31 (1.16, 1.48)	1.25 (1.10, 1.42)
6-11 months	2,344 (25.7%)	486 (27.5%)	1.24 (1.09, 1.41)	1.31 (1.15, 1.50)
12-59 months	3,986 (43.7%)	688 (37.8%)	(reference)	(reference)
<b>Sex</b>				
Male	5,099 (55.9%)	1,026 (58.1%)	(reference)	(reference)
Female	4,018 (44.1%)	741 (41.9%)	0.92 (0.83, 1.02)	0.92 (0.83, 1.02)
<b>Pneumonia classification</b>				
Fast breathing only	3,673 (40.3%)	379 (21.5%)	(reference)	(reference)
Lower chest indrawing	5,444 (59.7%)	1,388 (78.5%)	2.47 (2.19, 2.79)	1.50 (1.29, 1.73)
<b>Any sign of respiratory distress<sup>b</sup></b>				
No	5,299 (58.1%)	631 (35.7%)	(reference)	(reference)
Yes	3,818 (41.9%)	1,136 (64.3%)	2.50 (2.25, 2.78)	2.02 (1.78, 2.29)
<b>Respiratory rate</b>				
<age-adjusted tachypnea threshold <sup>a</sup>	110 (1.2%)	14 (0.8%)	(reference)	(reference)
0-9 breaths/min above cut-off	4,117 (45.2%)	483 (27.3%)	0.92 (0.52, 1.62)	1.07 (0.61, 1.90)
10-19 breaths/min above cut-off	3,298 (36.2%)	695 (39.3%)	1.66 (0.94, 2.90)	1.67 (0.95, 2.94)
>/=20 breaths/min above cut-off	1,592 (17.5%)	575 (32.5%)	2.83 (1.61, 4.99)	2.66 (1.51, 4.69)
<b>Axillary temperature</b>				
Normal (>/=35.5°C to 37.5°C)	5,839 (64.1%)	1,069 (60.5%)	(reference)	(reference)
>37.5°C	3,173 (34.8%)	681 (38.5%)	1.17 (1.06, 1.30)	1.02 (0.91, 1.14)
<35.5°C	105 (1.1%)	17 (1.0%)	0.88 (0.53, 1.48)	0.89 (0.52, 1.52)
<b>Weight for age z-score</b>				
>=-2	7,427 (81.5%)	1,350 (76.4%)	(reference)	(reference)
<-2 to -3	1,113 (12.2%)	251 (14.2%)	1.24 (1.07, 1.44)	1.16 (1.00, 1.36)
<-3	577 (6.3%)	166 (9.4%)	1.58 (1.32, 1.90)	1.59 (1.31, 1.93)

<sup>a</sup>IMCI chartbook threshold for tachypnea: age 2-11 months >=50 breaths per minute; age 12-59 months >=40 breaths per minute; <sup>b</sup> head nodding or nasal flaring or grunting

**Table S2.** Components of the PREPARE SpO<sub>2</sub> <92% risk score in the development dataset (n=10,884)

Factor	Adjusted log coefficient	PREPARE hypoxemia risk score <sup>a</sup>
<b>Age</b>		
2-5 months	0.22	--
6-11 months	0.27	+1
12-59 months	---	---
<b>Pneumonia classification</b>		
Fast breathing only	--	---
Lower chest indrawing	0.40	+1
<b>Respiratory rate</b>		
<age-adjusted tachypnea threshold <sup>b</sup>	--	---
0-9 breaths/min above cut-off	0.07	---
10-19 breaths/min above cut-off	0.51	---
>/=20 breaths/min above cut-off	0.98	+2
<b>Any sign of respiratory distress<sup>c</sup></b>		
No	---	--
Yes	0.70	+1
<b>Weight for age z-score</b>		
>=-2	--	---
<-2 to -3	0.15	--
<-3	0.46	+1

<sup>a</sup>Regression coefficients of each retained parameter with a statistically significant adjusted odds ratio (confidence interval (CI) did not overlap with 1.0) was rounded to the nearest 0.5 and then doubled to derive the points contributing to the PREPARE hypoxemia risk score; <sup>b</sup>IMCI chartbook threshold for tachypnea: age 2-11 months >=50 breaths per minute; age 12-59 months >=40 breaths per minute; <sup>c</sup> head nodding or nasal flaring or grunting

**Table S3.** Performance of the PREPARE SpO2 <92% risk score at each cut-off in the validation dataset (n=3,625)

Score	SpO2 <92%, n (%)*	Sensitivity (95% CI)*	Specificity (95% CI)*	Positive Likelihood Ratio (95% CI)*	Negative Likelihood Ratio (95% CI)*	Miss rate*† % (n/N)	False discovery rate*‡ % (n/N)	Proportion referred* ^ % (n/N)
0	37 (4.7%)	100.0	0.0	1.00	---	0% (0/275)	85.8% (3111/3625)	100% (3625/3625)
1	99 (12.5%)	92.8 (90.2, 94.9)	24.4 (22.9, 25.9)	1.23 (1.19, 1.27)	0.29 (0.21, 0.40)	7.2% (37/514)	83.1% (2353/2830)	78.1% (2830/3625)
2	157 (13.6%)	73.5 (69.5, 77.3)	46.6 (44.8, 48.3)	1.38 (1.29, 1.46)	0.57 (0.49, 0.66)	26.5% (136/514)	81.5% (1661/2039)	56.2% (2039/3625)
3	105 (20.5%)	43.0 (38.7, 47.4)	78.8 (77.3, 80.2)	2.02 (1.79, 2.28)	0.72 (0.67, 0.78)	57.0% (293/514)	74.9% (661/882)	24.3% (882/3625)
4	87 (28.4%)	22.6 (19.0, 26.4)	91.8 (90.8, 92.8)	2.76 (2.27, 3.37)	0.84 (0.80, 0.88)	77.4% (398/514)	68.7% (255/371)	10.2% (371/3254)
5	26 (43.3%)	5.6 (3.8, 8.0)	98.8 (98.4, 99.2)	5.01 (3.09, 8.13)	0.95 (0.93, 0.98)	94.4% (485/514)	55.4% (36/65)	1.8% (65/3625)
6	3 (75.0%)	0.6 (0.1, 1.7)	100.0 (99.8, 100.0)	18.2 (1.89, 174.0)	0.99 (0.99, 1.00)	99.4% (511/514)	25% (1/4)	0.1% (4/3625)

\*Calculated at  $\geq$  each respective cut-off

†Miss rate (also known as false negative rate = False Negative/(True Positive + False Negative)

‡False discovery rate = False Positive/(False Positive+True Positive)

^Referral used as a proxy for identification of hypoxemia, assuming that all children identified as at risk of hypoxemia would be referred

**Table S4:** Characteristics of studies included in the development and validation of the PREPARE hypoxemia risk score without respiratory distress

Study	Study design	Location	Age	Original inclusion criteria from the source studies	Original exclusion criteria from the source studies	Sample size included in the present analysis over the sample size from the original study, n/N (%)	Hypoxemia n/N (%)
Addo-Yobo 2004 <sup>39</sup>	Randomized controlled trial	Colombia, Ghana, India, Mexico, Pakistan, South Africa, Vietnam, Zambia	3-59 months	- Children with chest-indrawing pneumonia or WHO-defined severe pneumonia <sup>a</sup>	- Children with any danger sign during current illness (convulsions, abnormally sleepy or difficult to awake, stridor in a calm child). - Children with severe comorbidities and chronic conditions: severe malnutrition, measles, HIV infection, congenital cardiac or respiratory anomalies, etc - Children with other diseases requiring antibiotic therapy on presentation - Children with SpO <sub>2</sub> <80% - Children with known prior anaphylactic reaction to penicillin or amoxicillin - Children with known antibiotic therapy for 48 hours or more prior to admission - Children living outside the catchment area of the hospital - children unable to receive oral medications	1,251/1,702 (73.5%)	13.6% (170/1,251)
Basnet, 2012 <sup>40</sup>	Randomized controlled trial	Kathmandu, Nepal	2-35 months	- Cough for <14 days - Difficulty breathing ≤72 hours with presence of lower chest indrawing on examination	- Children with recurrent wheezing, heart disease, other severe illness, severe malnutrition,	192/641 (30.0%)	55.7% (107/192)

					dehydration, hemoglobin <70 g/L, chronic cough, effusion on chest x-ray, or history of documented tuberculosis.		
Bénet, 2017 <sup>41</sup>	Prospective case series	- Phnom Penh, Cambodia - Beijing, China - Port au Prince, Haiti - Lucknow, India - Pune-Vadu, India - Antananarivo, Madagascar - Bamako, Mali - Ulaanbaatar, Mongolia - San Lorenzo, Paraguay.	2-59 months	- Children with WHO-defined pneumonia with first symptoms lasting <14 days - Children with radiographic pneumonia (WHO criteria)	- Children with wheezing at auscultation	106/888 (11.9%)	5.7% (6/106)
Clara WS. Unpublished data 2012 <sup>42</sup>	Retrospective cohort	Chariqui Province, Panama	2-59 months	- Children with WHO-defined pneumonia of any severity <sup>b</sup>	None	9/85 (10.6%)	44.4% (4/9)
Cutts, 2005 <sup>38</sup>	Randomized controlled trial	Eastern Gambia	40-364 days	Children with WHO-defined pneumonia of any severity <sup>b</sup>	- Intent to move out within 4 months - Previous or uncertain receipt of diphtheria-pertussis-tetanus/ <i>Haemophilus influenzae</i> type b (DPT/Hib) or DPT vaccine - Inclusion in a previous vaccine trial - Serious chronic illness	323/1,716 (18.8%)	4.3% (14/323)
Ferolla, 2013 <sup>43</sup>	Prospective cohort	Buenos Aires, Argentina	< 2 years	- Children <2 years admitted with a diagnosis of severe acute respiratory illness, defined as the sudden onset of cough, wheezing, retractions and/or crackles with or without fever	None	59/5,043 (1.2%)	39.0% (23/59)
Gessner, 2005 <sup>44</sup>	Prospective cohort	Lombok, Indonesia	2-35 months	- Children with WHO-defined pneumonia of any severity <sup>b</sup>	- Received three doses of diphtheria-tetanus-pertussis (DTP) vaccine	4,464/6,221 (71.8%)	34.5% (1,542/4,464)
Klugman, 2003 <sup>45</sup>	Randomized controlled trial	Johannesburg, South Africa	2-59 months	ICD-10 codes for pneumonia	- Children with progressive underlying neurological disorder, history of seizures, or infantile spasms	1,517/10,114 (15.0%)	16.9% (256/1,517)

					- Low likelihood of receiving three doses of vaccine		
Lazzerini, 2016 <sup>46</sup>	Prospective cohort	Malawi	2-59 months	- Children with WHO-defined pneumonia of any severity <sup>b</sup>	None	2,640/16,475 (16.0%)	4.8% (127/2,640)
Mathew, 2015 <sup>47</sup>	Prospective cohort	Chandigarh, India	2-59 months	- Children with WHO-defined pneumonia of any severity <sup>b</sup>	- Duration of illness >7 days - Antibiotics for >24 hours - Previous hospitalization within the preceding 30 days - Children with wheeze who received a single dose of bronchodilator and whose symptoms disappeared	510/2400 (21.3%)	10.2% (52/510)
McCollum, 2017 <sup>48</sup>	Prospective cohort	Mchinji and Lilongwe Districts, Malawi	2-59 months	- Children with WHO-defined pneumonia of any severity <sup>b</sup>	None	3,601/6,764 (53.2%)	3.6% (129/3,601)
O'Grady, 2012 <sup>49</sup>	Randomized controlled trial	Central Australia	2-59 months	- Children with WHO-defined pneumonia of any severity <sup>b</sup>	Children with wheezing and chronic conditions	26/147 (17.7%)	0.0% (0/26)
Ugpo, 2009 <sup>50</sup>	Randomized controlled trial	Bohol, Philippines	2-59 months	- Children with WHO-defined pneumonia of any severity <sup>b</sup>	- Children with known seizure disorder presenting with febrile convulsions - Children with a history of benign febrile convulsion who present with fever and seizure with normal neurologic examination - Children with dengue shock syndrome	586/1,201 (48.8%)	14.0% (82/586)
Wadhwa, 2013 <sup>51</sup>	Randomized controlled trial	New Delhi, India	2-24 months	- Children with WHO-defined pneumonia of any severity <sup>b</sup> - Crepitations on auscultation	- Need for mechanical ventilation or inotropic medications - Any other serious underlying medical condition	308/550 (56.0%)	1.0% (3/308)

<sup>a</sup>WHO-defined severe pneumonia: presence of danger signs, stridor, central cyanosis, severe respiratory distress (nasal flaring, grunting, head-nodding), in children with cough and/or difficulty breathing

<sup>b</sup>WHO-defined pneumonia of any severity: presence of age-specific fast breathing, lower chest indrawing, or danger signs, in children with a cough and/or difficulty breathing.



**Table S5.** Baseline characteristics of children with pneumonia (fast breathing and/or chest indrawing), pulse oximetry measurements and complete data on all candidate predictors included in the PREPARE risk score development and validation when excluding respiratory distress (n=15,592)

Characteristic	Total (n=15,592)	Development dataset (n=11,731)	Validation dataset (n=3,861)
<b>Age<sup>a</sup></b>			
2-5 months	5,147 (33.0%)	4,041 (34.5%)	1,106 (28.6%)
6-11 months	4,257 (27.3%)	3,313 (28.2%)	944 (25.5%)
12-59 months	6,188 (39.7%)	4,377 (37.3%)	1,811 (46.9%)
<b>Sex</b>			
Male	8,022 (51.5%)	6,015 (51.3%)	2,007 (52.0%)
Female	7,570 (48.5%)	5,716 (48.7%)	1,854 (48.0%)
<b>Pneumonia classification</b>			
Fast breathing only	6,061 (38.9%)	4,489 (38.3%)	1,572 (40.7%)
Lower chest indrawing	9,531 (61.1%)	7,242 (61.7%)	2,289 (59.3%)
<b>Respiratory rate</b>			
<age-adjusted tachypnea threshold <sup>b</sup>	204 (1.3%)	90 (0.8%)	114 (3.0%)
0-9 breaths/min above cut-off	6,657 (42.7%)	4,995 (42.6%)	1,662 (43.1%)
10-19 breaths/min above cut-off	5,758 (36.9%)	4,351 (37.1%)	1,407 (36.4%)
>/=20 breaths/min above cut-off	2,973 (19.1%)	2,295 (19.6%)	678 (17.5%)
<b>Body temperature</b>			
Normal (>/=35.5°C to 37.5°C)	10,252 (65.7%)	7,678 (65.4%)	2,574 (66.7%)
>=37.5°C	5,250 (33.7%)	3,986 (34.0%)	1,264 (32.7%)
<35.5°C	90 (0.6%)	67 (0.6%)	23 (0.6%)
<b>Weight for age z-score</b>			
>=-2	11,690 (75.0%)	8,716 (74.3%)	2,974 (77.0%)
<-2 to -3	2,293 (14.7%)	1,770 (15.1%)	523 (13.6%)
<-3	1,609 (10.3%)	1,245 (10.6%)	364 (9.4%)
<b>Oxygen saturation (SpO<sub>2</sub>)</b>			
>=90%	13,077 (83.9%)	9,767 (83.3%)	3,310 (85.7%)
<90%	2,515 (16.1%)	1,964 (16.7%)	551 (14.3%)

<sup>a</sup>Some studies did not include patients in the whole 2-59 month age range; <sup>b</sup>IMCI chartbook threshold for tachypnea: age 2-11 months >=50 breaths per minute; age 12-59 months >=40 breaths per minute

**Table S6:** Multivariable regression model for predicting hypoxemia in the development dataset without respiratory distress as a candidate predictor (n=11,731)

Characteristic	No hypoxemia (n=9,767)	Hypoxemia (n=1,964)	Odds ratio (95% CI)	Adjusted OR (95% CI)
<b>Age</b>				
2-5 months	3,178 (32.5%)	863 (43.9%)	2.13 (1.89, 2.04)	1.92 (1.69, 2.19)
6-11 months	2,706 (27.7%)	607 (30.9%)	1.76 (1.55, 2.00)	1.65 (1.44, 1.90)
12-59 months	3,883 (39.8%)	4947 (25.2%)	(reference)	(reference)
<b>Sex</b>				
Male	5,083 (52.0%)	932 (47.4%)	(reference)	(reference)
Female	4,684 (48.0%)	1,032 (52.6%)	1.20 (1.09, 1.32)	1.25 (1.12, 1.39)
<b>Pneumonia classification</b>				
Fast breathing only	4,257 (43.6%)	232 (11.8%)	(reference)	(reference)
Lower chest indrawing	5,510 (56.4%)	1,732 (88.2%)	5.77 (5.00, 6.65)	4.44 (3.83, 5.15)
<b>Respiratory rate</b>				
<age-adjusted tachypnea threshold*	79 (0.8%)	11 (0.6%)	(reference)	(reference)
0-9 breaths/min above cut-off	4,475 (45.8%)	520 (26.5%)	0.83 (0.4, 1.58)	0.90 (0.46, 1.77)
10-19 breaths/min above cut-off	3,500 (35.8%)	851 (43.3%)	1.75 (0.93, 3.30)	1.44 (0.74, 2.81)
>=20 breaths/min above cut-off	1,713 (17.5%)	582 (29.6%)	2.44 (1.29, 4.62)	2.10 (1.08, 4.10)
<b>Body temperature</b>				
Normal (>=35.5°C to 37.5°C)	6,476 (66.3%)	1,202 (61.2%)	(reference)	(reference)
>37.5°C	3,230 (33.1%)	756 (38.5%)	1.26 (1.14, 1.39),	1.17 (1.05, 1.31)
<35.5°C	61 (0.6%)	6 (0.3%)	0.53 (0.23, 1.23)	0.75 (0.33, 1.73)
<b>Weight for age z-score</b>				
>=2	7,494 (76.7%)	1,222 (62.2%)	(reference)	(reference)
<-2 to -3	1,367 (14.0%)	403 (20.5%)	1.81 (1.59, 2.05)	1.67 (1.46, 1.91)
<-3	906 (9.3%)	339 (17.3%)	2.29 (2.00, 2.64)	1.93 (1.66, 2.25)

\*IMCI chartbook threshold for tachypnea: age 2-11 months >=50 breaths per minute; age 12-59 months >=40 breaths per minute

**Table S7.** Components of the PREPARE hypoxemia risk score in the development dataset without respiratory distress as a candidate predictor (n=11,731)

<b>Factor</b>	<b>Adjusted log coefficient</b>	<b>PREPARE hypoxemia risk score*</b>
<b>Age</b>		
2-5 months	0.65	+1
6-11 months	0.50	+1
12-59 months	--	---
<b>Pneumonia classification</b>		
Fast breathing only	--	---
Lower chest indrawing	1.49	+3
<b>Respiratory rate</b>		
<threshold	--	---
0-9 breaths above cut-off	-0.10	---
10-19 breaths above cut-off	0.37	---
>=20 breaths above cut-off	0.74	+1
<b>Weight for age z-score</b>		
>=-2	--	---
<-2 to -3	0.51	+1
<-3	0.66	+1

**Table S8.** Performance of the PREPARE hypoxemia risk score without respiratory distress at each cut-off in the validation dataset (n=3,861)

Score	Hypoxemia, n (%)	Sensitivity (95% CI) <sup>‡^* ^</sup>	Specificity (95% CI) <sup>‡* ^</sup>	Positive Likelihood Ratio (95% CI) <sup>‡</sup>	Negative Likelihood Ratio (95% CI) <sup>‡</sup>	Miss rate <sup>†</sup> % (n/N)	False discovery rate <sup>‡</sup> % (n/N)	Proportion referred <sup>^</sup> % (n/N)
0	3 (0.5%)	100.0	0.0	1.00	---	0% (0/551)	85.7% (3310/3861)	100% (3861/3861)
1	24 (3.1%)	99.5 (98.4, 99.9)	19.6 (18.3, 21.0)	1.24 (1.22, 1.26)	0.28 (0.01, 0.09)	0.5% (3/551)	82.9% (2660/3208)	83.1% (3208/3861)
2	6 (4.5%)	95.1 (93.0, 96.7)	42.5 (40.8, 44.2)	1.65 (1.60, 1.71)	0.11 (0.08, 0.17)	4.9% (27/551)	78.4% (1904/2428)	62.9% (2428/3861)
3	46 (11.0%)	94.0 (91.7, 95.8)	46.3 (44.6, 48.0)	1.75 (1.69, 1.82)	0.13 (0.09, 0.18)	6.0% (33/551)	77.4% (1777/2295)	59.4% (2295/3861)
4	278 (22.2%)	85.7 (82.5, 88.5)	57.6 (55.9, 59.3)	2.02 (1.92, 2.13)	0.25 (0.20, 0.31)	14.3% (79/551)	74.8% (1403/1875)	48.6% (1875/3861)
5	170 (30.1%)	35.2 (31.2, 39.4)	87.1 (85.9, 88.3)	2.74 (2.37, 3.16)	0.74 (0.70, 0.79)	64.8% (357/551)	68.7% (426/620)	16.1% (620/3861)
6	24 (42.9%)	4.4 (2.8, 6.4)	99.0 (98.6, 99.3)	4.51 (2.67, 7.59)	0.97 (0.95, 0.98)	95.6% (527/551)	57.1% (32/56)	1.5% (56/3861)

\*Calculated at  $\geq$  each respective cut-off

†Miss rate (also known as false negative rate = False Negative/(True Positive + False Negative)

‡False discovery rate = False Positive/(False Positive+True Positive)

^Referral used as a proxy for identification of hypoxemia, assuming that all children identified as at risk of hypoxemia would be referred

**Table S9:** Characteristics of studies included in the development and validation of the PREPARE hypoxemia risk score when including wheezing as a candidate predictor

Study	Study design	Location	Age	Original inclusion criteria from the source studies	Original exclusion criteria from the source studies	Sample size included in the present analysis over the sample size from the original study, n/N (%)	Hypoxemia % (n/N)
Basnet, 2012 <sup>29</sup>	Randomized controlled trial	Kathmandu, Nepal	2-35 months	- Cough for <14 days - Difficulty breathing ≤72 hours with presence of lower chest indrawing on examination	- Children with recurrent wheezing, heart disease, other severe illness, severe malnutrition, dehydration, hemoglobin <70 g/L, chronic cough, effusion on chest x-ray, or history of documented tuberculosis.	378/641 (59.0%)	66.4% (251/378)
Cutts, 2005 <sup>32</sup>	Randomized controlled trial	Eastern Gambia	40-364 days	Children with WHO-defined pneumonia of any severity*	- Intent to move out within 4 months - Previous or uncertain receipt of diphtheria-pertussis-tetanus/ <i>Haemophilus influenzae</i> type b (DPT/Hib) or DPT vaccine - Inclusion in a previous vaccine trial - Serious chronic illness	433/1,716 (25.2%)	5.1% (22/433)
Lazzeri, 2016 <sup>36</sup>	Prospective cohort	Malawi	2-59 months	- Children with WHO-defined pneumonia of any severity*	None	7254/16,475 (44.0%)	9.5% (687/7254)
Mathew, 2015 <sup>37</sup>	Prospective cohort	Chandigarh, India	2-59 months	- Children with WHO-defined pneumonia of any severity*	- Duration of illness >7 days - Antibiotics for >24 hours - Previous hospitalization within the preceding 30 days	1132/2400 (47.2%)	11.8% (134/1132)

					- Children with wheeze who received a single dose of bronchodilator and whose symptoms disappeared		
McCollum, 2017 <sup>38</sup>	Prospective cohort	Mchinji and Lilongwe Districts, Malawi	2-59 months	- Children with WHO-defined pneumonia of any severity*	None	4253/6,764 (62.9%)	6.2% (265/4253)
O'Grady, 2012 <sup>39</sup>	Randomized controlled trial	Central Australia	2-59 months	- Children with WHO-defined pneumonia of any severity*	None	26/147 (17.7%)	0.0% (0/26)
Wadhwa, 2013 <sup>41</sup>	Randomized controlled trial	New Delhi, India	2-24 months	- Children with WHO-defined pneumonia of any severity* - Crepitations on auscultation	- Need for mechanical ventilation or inotropic medications - Any other serious underlying medical condition	316/550 (56.0%)	0.9% (3/316)

\*WHO-defined pneumonia of any severity: presence of age-specific fast breathing, lower chest indrawing, or danger signs or stridor in a calm child, in children with a cough or difficulty breathing.

**Table S10.** Baseline characteristics of children with pneumonia (fast breathing and/or chest indrawing), pulse oximetry measurements and complete data on all candidate predictors included in the PREPARE risk score development and validation when including wheezing as a candidate predictor (n=13,792)

Characteristic	Total (n=13,792)	Development dataset (n=10,397)	Validation dataset (n=3,395)
<b>Age<sup>a</sup></b>			
2-5 months	4,050 (29.4%)	3,220 (31.0%)	830 (24.4%)
6-11 months	3,475 (25.2%)	2,705 (26.0%)	770 (22.7%)
12-59 months	6,267 (45.4%)	4,472 (43.0%)	1,795 (52.9%)
<b>Sex</b>			
Male	7,767 (56.3%)	5,843 (56.2%)	1,924 (56.7%)
Female	6,025 (43.7%)	4,554 (43.8%)	1,471 (43.3%)
<b>Pneumonia classification</b>			
Fast breathing only	5,416 (39.3%)	3,977 (38.2%)	1,439 (42.4%)
Lower chest indrawing	8,376 (60.7%)	6,420 (61.8%)	1,956 (57.6%)
<b>Respiratory rate</b>			
<age-adjusted tachypnea threshold <sup>b</sup>	157 (1.1%)	120 (1.1%)	37 (1.1%)
0-9 breaths/min above cut-off	5,881 (42.6%)	4,428 (42.6%)	1,453 (42.8%)
10-19 breaths/min above cut-off	5,047 (36.6%)	3,790 (36.4%)	1,257 (37.0%)
≥20 breaths/min above cut-off	2,707 (19.6%)	2,059 (19.8%)	648 (19.1%)
<b>Wheeze</b>			
No	9,805 (71.1%)	7,496 (72.1%)	2,309 (68.0%)
Yes	3,987 (28.9%)	2,901 (27.9%)	1,086 (32.0%)
<b>Any sign of respiratory distress<sup>c</sup></b>			
No	7,791 (56.5%)	5,823 (56.0%)	1,968 (58.0%)
Yes	6,001 (43.5%)	4,574 (44.0%)	1,427 (42.0%)
<b>Axillary temperature</b>			
Normal (≥35.5°C to 37.5°C)	8,940 (64.8%)	6,619 (63.7%)	2,321 (68.4%)
>37.5°C	4,708 (34.2%)	3,663 (35.2%)	1,045 (30.8%)
<35.5°C	144 (1.0%)	115 (1.1%)	29 (0.9%)
<b>Weight for age z-score</b>			
≥-2	11,155 (80.9%)	8,364 (80.5%)	2,791 (82.2%)
<-2 to -3	1,709 (12.4%)	1,316 (12.7%)	393 (11.6%)
<-3	928 (6.7%)	717 (6.9%)	211 (6.2%)
<b>Oxygen saturation (SpO<sub>2</sub>)</b>			
≥90%	12,430 (90.1%)	9,296 (89.4%)	3,134 (92.3%)
<90%	1,362 (9.9%)	1,101 (10.6%)	261 (7.7%)

<sup>a</sup>Some studies did not include patients in the whole 2-59 month age range; <sup>b</sup>IMCI chartbook threshold for tachypnea: age 2-11 months ≥50 breaths per minute; age 12-59 months ≥40 breaths per minute; <sup>c</sup> head nodding or nasal flaring or grunting

**Table S11.** Multivariable regression model for predicting hypoxemia in the development dataset when including wheezing as a candidate predictor (n=13,792)

Characteristic	No hypoxemia (n=9,296)	Hypoxemia (n=1,101)	Odds ratio (95% CI)	Adjusted OR (95% CI)
<b>Age</b>				
2-5 months	2,833 (30.5%)	387 (35.2%)	1.38 (1.19, 1.60)	1.34 (1.14, 1.56)
6-11 months	2,393 (25.7%)	312 (28.3%)		
12-59 months	4,070 (43.8%)	402 (36.5%)	1.32 (1.13, 1.54) (reference)	1.44 (1.22, 1.70) (reference)
<b>Sex</b>				
Male	5,196 (55.9%)	647 (58.8%)	(reference)	(reference)
Female	4,100 (44.1%)	454 (41.2%)	0.89 (0.78, 1.01)	0.90 (0.79, 1.03)
<b>Pneumonia classification</b>				
Fast breathing only	3,764 (40.5%)	213 (19.3%)	(reference)	(reference)
Lower chest indrawing	5,532 (59.5%)	888 (80.7%)	2.84 (2.43, 3.31)	1.60 (1.34, 1.93)
<b>Respiratory rate</b>				
<age-adjusted tachypnea threshold <sup>a</sup>	110 (1.2%)	10 (0.9%)	(reference)	(reference)
0-9 breaths/min above cut-off	4,155 (44.7%)	273 (24.8%)	0.72 (0.37, 1.40)	0.87 (0.45, 1.70)
10-19 breaths/min above cut-off	3,371 (36.3%)	419 (38.1%)	1.37 (0.71, 2.63)	1.39 (0.72, 2.69)
>=20 breaths/min above cut-off	1,660 (17.9%)	399 (36.2%)	2.64 (1.37, 5.10)	2.47 (1.27, 4.79)
<b>Wheeze</b>				
No	6,809 (73.2%)	687 (62.4%)	(reference)	(reference)
Yes	2,487 (26.8%)	414 (37.6%)	1.65 (1.45, 1.88)	1.28 (1.11, 1.47)
<b>Any sign of respiratory distress<sup>b</sup></b>				
No	5,453 (58.7%)	370 (33.6%)	(reference)	(reference)
Yes	3,843 (41.3%)	731 (66.4%)	2.80 (2.46, 3.20)	2.25 (1.84, 2.52)
<b>Body temperature</b>				
Normal (>=35.5°C to 37.5°C)	5,964 (64.2%)	655 (59.5%)	(reference)	(reference)
>37.5°C	3,227 (34.7%)	436 (39.6%)	1.23 (1.08, 1.40)	1.06 (0.93, 1.22)
<35.5°C	105 (1.1%)	10 (0.9%)	0.87 (0.45, 1.67)	0.94 (0.48, 1.85)
<b>Weight for age z-score</b>				
>=-2	7,562 (81.3%)	802 (72.8%)	(reference)	(reference)
<-2 to -3	1,141 (12.3%)	175 (15.9%)	1.45 (1.21, 1.72)	1.35 (1.12, 1.62)
<-3	593 (6.4%)	124 (11.3%)	1.97 (1.60, 2.42)	1.98 (1.59, 2.48)

<sup>a</sup>IMCI chartbook threshold for tachypnea: age 2-11 months >=50 breaths per minute; age 12-59 months >=40 breaths per minute; <sup>b</sup> head nodding or nasal flaring or grunting



**Table S12.** Components of the PREPARE hypoxemia risk score in the development dataset when including wheezing as a candidate predictor (n=10,397)

Factor	Adjusted log coefficient	PREPARE hypoxemia risk score including wheezing*
<b>Age</b>		
2-5 months	0.29	+1
6-11 months	0.37	+1
12-59 months	---	--
<b>Pneumonia classification</b>		
Fast breathing only	--	--
Lower chest indrawing	0.47	+1
<b>Wheeze</b>		
No	---	--
Yes	0.24	+0
<b>Respiratory rate</b>		
< age-adjusted tachypnea threshold <sup>a</sup>	--	--
0-9 breaths/min above cut-off	-0.14	--
10-19 breaths/min above cut-off	0.33	--
>/=20 breaths/min above cut-off	0.90	+2
<b>Any sign of respiratory distress<sup>b</sup></b>		
No	--	--
Yes	0.77	+2
<b>Weight for age z-score</b>		
>=-2	--	+0
-2 to -3	0.30	+1
<-3	0.68	+1

\*Regression coefficients of each retained parameter with a statistically significant adjusted odds ratio (confidence interval (CI) did not overlap with 1.0) was rounded to the nearest 0.5 and then doubled to derive the points contributing to the PREPARE hypoxemia risk score. Maximum score is 7.

<sup>a</sup>IMCI chartbook threshold for tachypnea: age 2-11 months >=50 breaths per minute; age 12-59 months >=40 breaths per minute; <sup>b</sup> head nodding or nasal flaring or grunting


**Table S13.** Comparison of risk score weights from the main analysis and the sensitivity analyses

	Main score	< 92%	Without respiratory distress	With wheezing
Age 2-5 months	1	-	1	1
Age 6-11 months	1	1	1	1
Lower chest indrawing	1	1	3	1
Sign of respiratory distress	2	1	-	2
Respiratory rate $\geq 20$ breaths/min <sup>a</sup>	2	2	1	2
WAZ <-2 to 3	1	-	1	1
WAZ <-3	1	1	1	1
<b>TOTAL</b>	<b>7</b>	<b>6</b>	<b>6</b>	<b>7</b>

WAZ: Weight-for-age z-score

<sup>a</sup> a above age-adjusted tachypnea threshold. IMCI chartbook threshold for tachypnea: age 2-11 months  $\geq 50$  breaths per minute; age 12-59 months  $\geq 40$  breaths per minute


**Figure S1.** Decision curve analysis of the PREPARE hypoxemia model




**Legend:** The net benefit of the different score thresholds (score\_cat4 = refer children with a score of  $\geq 4$ ; score\_cat5 = refer children with a score of  $\geq 5$ ; score\_cat6 = refer children with a score of  $\geq 6$ ; and Pr(spocat1) = Children with a predicted probability of hypoxemia greater than or equal to the threshold probability [referral threshold] are referred) are compared to a “Treat all” approach where all children are referred, and a “Treat None” approach where no children are referred. A threshold probability of 20% indicates a management strategy whereby any child with a  $\geq 20\%$  probability of being hypoxemic is referred (i.e., a scenario where the value of one correct referral is equivalent to four incorrect referrals or a number-needed-to-refer [NNR] of 5).

**Figure S2.** Receiver operating characteristic curve for the PREPARE SpO<sub>2</sub><92% clinical prediction model for children 2–59 months of age with pneumonia


**A. Development dataset (n=10,884), AUROC 0.67, 95% CI 0.67 to 0.69**



**B. Validation dataset (n=3,625), AUROC 0.67, 95% CI 0.65 to 0.70**




**Figure S3.** Calibration plot of the PREPARE SpO<sub>2</sub> <92% clinical prediction model in the validation dataset




**Legend:** Reference (dashed line) indicates perfect calibration; Lowess (blue line) indicates calibration slope; Red rug plots indicate the distribution of predicted risk of participants with hypoxemia (above the line), and those without hypoxemia (below the line). E:O = expected to observed events ratio; CITL = calibration-in-the-large (intercept); AUC = area under the receiver operating characteristic curve.

**Figure S4** Receiver operating characteristic curve for the PREPARE hypoxemia clinical prediction model excluding respiratory distress as a candidate predictor for children 2–59 months of age with pneumonia


**A. Development dataset (n=11,731), AUROC 0.73, 95% CI 0.72 to 0.75**



**B. Validation dataset (n=3,861), AUROC 0.78, 95% CI 0.76 to 0.80**




**Figure S5:** Calibration plot of the PREPARE hypoxemia clinical prediction model in the validation dataset when excluding respiratory distress as a candidate predictor




**Legend:** Reference (dashed line) indicates perfect calibration; Lowess (blue line) indicates calibration slope; Red rug plots indicate the distribution of predicted risk of participants with hypoxemia (above the line), and those without hypoxemia (below the line). E:O = expected to observed events ratio; CITL = calibration-in-the-large (intercept); AUC = area under the receiver operating characteristic curve.

**Figure S6.** Receiver operating characteristic curve for the PREPARE hypoxemia clinical prediction model when including wheezing as a candidate predictor for children 2–59 months of age with pneumonia

**A. Development dataset (n=10,397), AUROC 0.71, 95% CI 0.69 to 0.72**




**B. Validation dataset (n=3,395), AUROC 0.72, 95% CI 0.69 to 0.75**





**Figure S7:** Calibration plot of the PREPARE hypoxemia clinical prediction model in the validation dataset when including wheezing as a candidate predictor



**Legend:** Reference (dashed line) indicates perfect calibration; Lowess (blue line) indicates calibration slope; Red rug plots indicate the distribution of predicted risk of participants with hypoxemia (above the line), and those without hypoxemia (below the line). E:O = expected to observed events ratio; CITL = calibration-in-the-large; AUC = area under the receiver operating characteristic curve.