

Appendices

Appendix 1: Projected population and live-births in 2014 in the catchment areas of the six acute jaundice surveillance hospitals, Bangladesh

Surveillance hospitals	No. of unions	Census population ¹	Population projection ²			Projected population ≥ 14 years ⁴	Projected live-birth ⁵
			2012	2013	2014		
Bogra							
Rural	12	427,635	433,494	439,432	445,453		
Urban ³	2	34,414	-		-		
Total	14				479,867	324,870	10,653
Barisal							
Rural	11	214,544	217,483	220,463	223,483		
Urban ³	4	49,278	-		-		
Total	15				272,761	184,659	6,055
Kishoregonj							
Rural	6	153,425	155,527	157,658	159,818		
Urban ³	1	3,645	-		-		
Total	7				163,463	110,664	3,629
Chittagong							
Rural	19	548,491	556,005	563,623	571,344		
Urban ³	6	101,065	-		-		
Total	25				672,409	455,221	14,927
Sylhet							
Rural	15	393,194	398,581	404,041	409,577		
Urban ³	4	47,585	-		-		
Total	19				457,162	309,498	10,149
Mitford, Dhaka							
Rural	3	107,617	109,091	110,586	112,101		
Urban ³	8	199,249	-		-		
Total	11				311,350	210,784	6,912
All							
Rural	66	1,844,906	1,870,181	1,895,803	1,921,775	1301,042	42,663
Urban ³	25	435,236	-		435,236	294,655	9,662
Total	91	2,280,142			2,357,011	1,595,697	52,326

¹ Population census in Bangladesh was conducted in 2011

² Projected population considering 1.37% growth rate

³ Population in urban areas was counted through house-to-house visits in the mortality survey

⁴ Projected considering 67.7% of population are ≥ 14 years

⁵ Live-births are projected considering crude birth rate as 22.2/1000 population

Appendix 2: Estimation of HEV associated mortality in the catchment areas of six acute jaundice surveillance hospitals by recall period, 2014 (Survey period: November 2014–February 2017)

Mortality estimation (indicators)	Recall period		
	One year	Two years	Three years
<i>Estimated population and live births in the hospital catchment areas</i> (projected for 2014; Appendix 1)			
Population aged ≥ 14 years [P]	1,595,697	1,595,697	1,595,697
Live births [B]	52,326	52,326	52,326
<i>Surveillance hospitals</i>			
Maternal deaths			
Maternal deaths with acute jaundice [J_m]	27	27	27
HEV cases among maternal deaths with acute jaundice [E_m] (95% CI)	8 (4-13)	8 (4-13)	8 (4-13)
Non-maternal deaths			
Non-maternal deaths with acute jaundice [$\sum J_i$]	275	275	275
HEV cases among non-maternal deaths with acute jaundice [$\sum E_i$] (95% CI)	20 (13-30)	20 (13-30)	20 (13-30)
Stillbirths			
Stillbirths delivered by mothers with acute jaundice [S_j]	27	27	27
Stillbirths delivered by HEV positive mothers [S_e] (95% CI)	6 (3-11)	6 (3-11)	6 (3-11)
Neonatal deaths			
Neonatal deaths born to mothers with acute jaundice [N_j]	15	15	15
Neonatal death cases born to HEV positive mothers [N_e] (95% CI)	10 (6-13)	10 (6-13)	10 (6-13)

Mortality estimation (indicators)	Recall period		
	One year	Two years	Three years
Hospital catchment areas			
Maternal deaths with acute jaundice [D_m] (95% CI)	9 (4-17)	19 (11-30)	25 (16-37)
Non-maternal deaths with acute jaundice [$\sum D_i$] (95% CI)	214 (186-245)	437 (401-480)	587 (401-772)
Stillbirths delivered by mothers with acute jaundice [S] (95% CI)	28 (19-40)	51 (38-67)	57 (43-74)
Neonatal deaths born to mothers with acute jaundice [N] (95% CI)	21 (13-32)	35 (24-49)	53 (40-69)
HEV associated mortality			
HEV associated mortality per 100,000 population aged ≥ 14 years (95% CI)	1.0 (0.4-1.8)	1.1 (0.6-1.5)	0.93 (0.6-1.3)
Maternal mortality ratio due to HEV per 100,000 live-births (95% CI)	5.1 (1.2-15.8)	5.3 (1.7-13.7)	4.7 (1.6-11.4)
Stillbirth rate due to HEV per 100,000 live-births (95% CI)	11.9 (3.8-31.5)	10.8 (3.8-26.1)	8.1 (2.9-19.2)
Neonatal mortality rate due to HEV per 100,000 live-births (95% CI)	26.7 (10.3-52.0)	22.3 (9.7-39.4)	22.5 (10.5-37.5)
Estimated HEV associated annual deaths in Bangladesh			
Deaths among population aged ≥ 14 years	1102 (422-1901)	1114 (634-1584)	986 (599-1338)
Maternal deaths	177 (43-548)	186 (60-476)	163 (57-395)
Stillbirths	412 (131-1092)	375 (133-904)	279 (101-664)
Neonatal deaths	927 (359-1802)	780 (337-1367)	780 (365-1298)