

Ketamine treatment effects on DNA methylation and Epigenetic Biomarkers of aging

Kristin Dawson^{1†}, Athena May Jean M. Carangan^{2†}, Jessica Klunder³, Natalia Carreras-Gallo², Raghav Sehgal⁴, Samantha Megilligan¹, Benjamin C. Askins³, Nicole Perkins³, Tavis L. Mendez², Ryan Smith², Matthew Dawson², Michael Mallin², Albert T. Higgins-Chen^{5,6}, Varun B. Dwaraka^{2*}

Supplementary Material

Supplementary Tables

Supplementary Table 1. Effect of ketamine treatment series on non-epigenetic variables.

Mean and median values are calculated at baseline and after administration of ketamine. The final column shows the p-value of the difference between treatments. Significantly changed terms are bolded.

	Mean		Median		p-value	PC-Bonferoni
	Pre-	Post-	Pre-	Post-		
PCL-5	49.55	20.45	51	18	0.0000038	0.0000076
PHQ-9	16.85	7.4	18	7	0.0000949	0.0001898
A/G Ratio	1.913	1.969	1.9	1.9	0.1563	0.3126
Albumin	4.431	4.487	4.45	4.5	0.3253	0.6506
Alkaline Phosphatase	72.562	72.688	66.5	68	0.8764	1.000
ALT (SGPT)	19.188	17.875	18	16.5	0.1917	0.3834
Apolipoprotein B	82.25	83.188	83	82	0.8155	1.000
AST (SGOT)	19.875	18.875	18	18.5	0.0863	0.1726
Basophils (Absolute)	0.019	0.031	0	0	0.4237	0.8474
Basophils	0.75	0.812	1	1	0.7656	1.000
Bilirubin, Total	0.423	0.523	0.4	0.4	0.1747	0.3494
BUN	12.188	12	11	11	0.8243	1.000

BUN/Creatinine Ratio	14	13.75	12.5	13	0.7786	1.000
C-Reactive Protein, Cardiac	2.492	3.5	1.195	0.59	0.2412	0.4824
Calcium	9.456	9.456	9.6	9.45	0.4889	0.9778
Carbon Dioxide, Total	23.625	24.688	23	24	0.0776	0.1552
Chloride	102.25	102.125	102	102.5	0.8050	1.000
Cholesterol, Total	181.562	190.625	184	180.5	0.1704	0.3408
Cortisol	12.938	11.325	13.8	11.65	0.2744	0.5488
Creatinine	0.869	0.883	0.835	0.87	0.5925	1.000
eGFR	94.625	92.062	92	92.5	0.4887	0.9774
Eosinophils	2.688	2.688	2	2	0.4627	0.9254
Eosinophils (Absolute)	0.144	0.169	0.1	0.1	1	2
Folate (Folic Acid), Serum	9.875	9.088	10.35	7.95	0.6726	1.000
Free Testosterone (Direct)	4.486	3.764	2.25	2.3	0.975	1.000
Globulin, Total	2.369	2.331	2.45	2.25	0.609	1.000
Glucose	92	93.125	93.5	93	0.5516	1.000
HDL-C	57.5	56.562	64.5	59	0.842	1.000
HDL-P	34.425	34.444	34.5	35.3	1	1.000
Hematocrit	41.512	41.219	41.4	40.3	0.7173	1.000
Hemoglobin	13.85	13.85	13.9	13.6	0.9587	1.000
Hemoglobin A1c	5.4	5.388	5.4	5.5	0.5885	1.000
Homocysteine	8.988	9.488	8.85	9.25	0.1543	0.3086

Immature Granulocytes	0.071	0	0	0	1	1.000
Insulin	12.1	12.481	9	9.2	0.7057	1.000
LDL-C (NIH Calc)	104.812	113.688	99	105.5	0.0743	0.1486
LDL-P	1245.357	1258.071	1231	1175.5	0.4897	0.9794
LDL Size	21.2	21.4	21.3	21.4	0.0829	0.1658
LP-IR Score	59.125	51.25	57	52	0.1083	0.2166
Lymphocytes	34.312	34.688	34.5	35.5	0.6596	1.000
Lymphocytes (Absolute)	1.8	1.9	1.7	1.75	0.3301	0.6602
MCH	30.275	30.262	30.25	29.95	0.7365	1.000
MCHC	33.388	33.606	32.9	33.6	0.3254	0.6508
MCV	90.75	90.125	91	90.5	0.1411	0.2822
Monocytes	8.5	7.875	8	8	0.0857	0.1714
Monocytes (Absolute)	0.469	0.431	0.4	0.4	0.0723	0.1446
Neutrophils	53.688	53.938	52	53.5	0.8423	1.000
Neutrophils (Absolute)	3.138	3.038	3.05	2.6	0.7796	1.000
Platelets	261.312	273.188	260.5	277	0.2242	0.4484
Potassium	4.412	4.312	4.5	4.35	0.2038	0.4076
Protein, Total	6.8	6.819	6.7	6.8	0.7748	1.000
RBC	4.585	4.582	4.61	4.48	0.8971	1.000
RDW	12.656	12.525	12.7	12.4	0.2545	0.509
Small LDL-P	389.385	334	344	343	0.2459	0.4918
Sodium	140	139.75	140	140	0.7172	1.000
T4, Free (Direct)	1.204	1.22	1.185	1.245	0.7981	1.000

Triglycerides	106.812	113	92.5	98.5	0.6292	1.000
Triiodothyronine (T3), Free	3.212	3.175	3.2	3.15	0.7062	1.000
TSH	1.879	1.726	1.88	1.52	0.5282	1.000
Vitamin B12	581.938	581.188	564	612.5	0.7057	1.000
Vitamin D, 25-Hydroxy	37.594	37.225	34.05	35.35	0.9399	1.000
WBC	5.575	5.581	5.3	5.2	0.7366	1.000

Supplementary Table 2. Impact of Ketamine treatment on epigenetic age. The average change in epigenetic age acceleration (EAA) for 17 epigenetic clocks is shown, as measured at baseline and after administration of ketamine. The final column shows the p-value of the difference between treatments. Clocks exhibiting significant changes in their epigenetic age between timepoints are bolded.


	Glass Δ	Standard Error Difference	p-value	PCA Adj. Bonferroni
PC PhenoAge EAA	0.98	1.759	0.2	1.000
PC DNAmTL EAA	0.0042	0.028	0.7	1.000
PC GrimAge EAA	-0.34	0.660	0.29	1.000
GrimAge V2 EAA	-1.1		0.021	0.105
OMICmAge EAA	-0.98	0.693	0.0094	0.047
DunedinPACE	0.01		0.87	1.000
PhenoAge EAA	-2	0.043	0.024	0.120
Systems Age Blood EAA	-0.56	2.545	0.55	1.000
Systems Age Brain EAA	-1.48	3.691	0.41	1.000
Systems Age Inflammation EAA	-1.9	3.544	0.076	0.380
Systems Age Heart EAA	-0.82	3.057	0.37	1.000
Systems Age Hormone EAA	-0.42	1.613	0.78	1.000

Systems Age Immune EAA	-0.98	3.054	0.41	1.000
Systems Age Kidney EAA	-1.28	3.069	0.28	1.000
Systems Age Liver EAA	-1.38	2.999	0.33	1.000
Systems Age Lung EAA	-1.08	2.503	0.29	1.000
Systems Age Metabolic EAA	-2.94	4.070	0.12	0.600
Systems Age Musculoskeletal EAA	-2.78	3.555	0.11	0.550
SystemsAge EAA	-1.1	3.135	0.29	1.000

Supplementary Table 3. Comparison of immune cell subsets before and after Ketamine treatment. Mean and median values for 12 immune subsets estimated using DNA methylation are calculated at baseline and after administration of ketamine. The final column shows the p-value of the difference between timepoints. The significant differences are bolded.

	Mean		Median		p-value
	Pre-treatment	Post-treatment	Pre-treatment	Post-treatment	
CD4T Naive	0.083	0.073	0.084	0.069	0.18
CD4T Memory	0.046	0.029	0.05	0.025	0.038
CD8T Naive	0.029	0.014	0.013	0.004	0.17
CD8T Memory	0.054	0.062	0.044	0.052	0.35
B naive	0.033	0.023	0.033	0.018	0.12
B memory	0.014	0.013	0.011	0.013	0.93
Basophil	0.012	0.009	0.012	0.009	0.1
Regulatory T cells	0.033	0.038	0.03	0.034	0.19
Eosinophil	0.003	0.001	0	0	0.14
Natural Killer	0.028	0.023	0.019	0.025	0.62
Neutrophil	0.626	0.687	0.612	0.675	0.053
Monocyte	0.04	0.029	0.04	0.034	0.11

Supplementary Figure 1. Scatter Plot of Individual Scores at Baseline. The x-axis shows the treatment groups and the y-axis corresponds to the patients' self-reported score for each test at baseline. The colors correspond to the patients' diagnosis: MDD alone, PTSD alone, or comorbid MDD and PTSD. The red dashed line corresponds to the minimum score needed for a diagnosis of moderately severe depression based on the PHQ-9. The green dashed line corresponds to the minimum score needed for a diagnosis of PTSD.



Supplementary Figure 2. Correlation plot showing the strength of association between changes in non-epigenetic measures and epigenetic age from baseline to post-treatment. This correlation plot has all variables of interest in the x- and y-axis. The strength of correlation is denoted by the size and color of the circles in each cell. Coefficient values are listed in each cell, and asterisks have been used to denote significance. * for $p < 0.05$, ** for $p < 0.01$, and *** for $p < 0.001$.

