



Supplementary Materials




Supplementary Figure 1. Types of infections and their frequencies (n = 218, infections = 242)


Note. The same individual had multiple infections on a few occasions. *Abbreviations:* PUD, Peptic Ulcer Disease; URTI, Upper Respiratory Tract Infection; UTI, Urinary Tract Infection.



Supplementary Figure 2. Study flowchart of participant selection



Supplementary Figure 3. Distribution of HCI values (A) Distribution for individuals with infections (B) Distribution for individuals with Alzheimer's disease (AD) (C) Distribution for individuals with *APOE4*



Supplementary Figure 4. Correlation between continuous covariates with investigated brain metabolism measures.

Note. The darker shades indicate strong positive or negative correlations.

Supplementary Table 1. Regression estimates for predictors in the multivariate linear regression full model for HCl outcome

Variables	Estimates	95% CI	p
Infections (Yes)	0.15	0.03 -0.28	0.01*
Age	0.01	0.01, 0.02	<0.001***
Sex (Male)	0.17	0.08, 0.26	<0.001***
Education (Years)	-0.002	-0.02, 0.01	0.81
Marriage (Never)	-0.06	-0.29, 0.17	0.61
<i>APOE4</i>	0.32	0.25, 0.38	<0.001***
Race (White)	0.25	0.09, 0.42	0.002**
Smoking (Yes)	0.08	-0.01, 0.19	0.095
Alcohol (Yes)	-0.02	-0.27, 0.23	0.873
AD (Yes)	1.04	0.92, 1.16	<0.001***
Diabetes Medication (Yes)	0.19	-0.07, 0.45	0.159

Note. *p<0.05; **p<0.01; ***p<0.001.

Supplementary Table 2. Measures of variable importance computed using Random forest models

Variable	%IncMSE	IncNodepurity
AD	81.59	22222.41
<i>APOE4</i>	36.10	6778.73
Age	13.72	10642.90

Sex	12.31	1466.75
Race	9.67	1021.57
Smoking	4.54	1048.22
Infections	0.63	1014.08

Note. %IncMSE represents the increase in the regression mean-squared error when variables are randomly permuted. IncNodePurity measures the loss function based on the optimal node splits that maximize variance. Higher values for both metrics indicate more important variables.

Supplementary Table 3. Marginal means showing the model-predicted HCl stratified by sex

Infections	Sex	Mean age	Age range	Marginal means	SE	95% CI	p-value
No	Female	72.23	55.0 - 89.6	0.26	0.05	0.15, 0.37	<0.001***
Yes	Female	72.92	60.0 - 90.3	0.41	0.07	0.25, 0.56	<0.001***
No	Male	73.95	55.0 - 91.4	0.43	0.05	0.33, 0.54	<0.001***
Yes	Male	75.65	57.8 - 89.3	0.58	0.07	0.44, 0.73	<0.001***

Note. The means are averaged over the levels of age, sex, race, smoking, and AD.

Supplementary Table 4. Marginal means showing the model-predicted HCl stratified by APOE4

Infections	APOE4	Marginal means	SE	95% CI	p-value
No	0	0.03	0.05	-0.07, 0.13	0.53
Yes	0	0.18	0.07	0.03, 0.33	0.01*
No	1	0.38	0.05	0.27, 0.49	<0.001***
Yes	1	0.53	0.07	0.38, 0.68	<0.001***
No	2	0.62	0.08	0.47, 0.78	<0.001***
Yes	2	0.77	0.09	0.58, 0.97	<0.001***

Note. The means are averaged over the levels of age, sex, race, smoking, and AD.

Supplementary Table 5. Association of number of infections with HCl

Variables	Estimates	95% CI	p
Infections (n=1)	0.11	-0.01, 0.24	0.08

Infections (n>=2)	0.44	0.09, 0.80	0.01*
Age	0.01	0.006, 0.01	<0.001***
Sex (Male)	0.17	0.08, 0.25	<0.001***
APOE4	0.31	0.25, 0.38	<0.001***
Race (White)	0.25	0.09, 0.41	0.002**
Smoking (Yes)	0.08	-0.01, 0.18	0.09
AD (Yes)	1.03	0.92, 1.15	<0.001***

Note. *p<0.05; **p<0.01; ***p<0.001.

Supplementary Table 6. Regression estimates for predictors in the multivariate linear regression full model for SROI AD outcome

Variables	Estimates	95% CI	p
Infections (Yes)	-0.01	-0.02, -0.001	0.02*
Age	-0.003	-0.003, -0.002	<0.001***
Sex (Male)	-0.01	-0.01, -0.002	0.009**
Education (Years)	0.001	0.000, 0.002	0.043*
Marriage (Never)	0.01	-0.01, 0.03	0.26
APOE4	-0.02	-0.03, -0.02	<0.001***
Race (White)	-0.01	-0.02, 0.004	0.18
Smoking (Yes)	-0.01	-0.02, 0.001	0.10
Alcohol (Yes)	0.01	-0.01, 0.03	0.52
AD (Yes)	-0.08	-0.09, -0.07	<0.001***
Diabetes Medication (Yes)	-0.03	-0.05, -0.005	0.01**

Note. *p<0.05; **p<0.01; ***p<0.001.

Supplementary Table 7. Regression estimates for predictors in the multivariate linear regression full model for SROI MCI outcome

Variables	Estimates	95% CI	p
Infections (Yes)	-0.013	-0.026, -0.0005	0.04*
Age	-0.004	-0.005, -0.004	<0.001***
Sex (Male)	-0.022	-0.032, -0.013	<0.001***
Education (Years)	0.001	-0.0005, 0.002	0.19
Marriage (Never)	0.007	-0.016, 0.031	0.52
<i>APOE4</i>	-0.022	-0.029, -0.016	<0.001***
Race (White)	-0.005	-0.022, 0.010	0.49
Smoking (Yes)	-0.006	-0.016, 0.0038	0.22
Alcohol (Yes)	-0.003	-0.028, 0.022	0.81
AD (Yes)	-0.090	-0.102, -0.078	<0.001***
Diabetes Medication (Yes)	-0.035	-0.061, -0.008	<0.01**

Note. *p<0.05; **p<0.01; ***p<0.001.