

Supplementary Tables

Supplementary Table 1 – Genetic associations of reticular pseudodrusen identified in previous studies

Source	CFH Y402H†	ARMS2	HTRA1	CFH I62V†	C3	C2/CFB‡	VEGFA	LIPC	SKIV2L
Klein et al., (Beaver Dam Eye Study), 2008 ³⁰	↑	--	--	--	--	--	--	--	--
Smith et al., 2011 ³¹	↓	↑	--	--	--	--	--	--	--
Ueda-Arakawa et al., 2013 ³²	0	↑	--	0	--	--	--	--	--
Puche et al., 2013 ³³	↑	↑	↑	--	↑	--	--	--	--
Joachim et al., (Blue Mountain Eye Study), 2014 ³⁴	↑	↑	--	--	--	--	--	--	--
Yoneyama et al., 2014 ³⁵	--	↑	--	0	--	--	--	--	--
Boddu et al., 2014 ³⁶	0	0	--	--	--	--	--	--	--

Continued on next page

Source	CFH Y402H†	ARMS2	HTRA1	CFH I62V†	C3	C2/CFB‡	VEGFA	LIPC	SKIV2L
Finger et al. (Melbourne Collaborative Cohort Study), 2016 ³⁷	↑	↑	↑	↑	--	--	--	--	--
Buitendijk et al., (Rotterdam Study), 2016 ³⁸	↑	↑	--	↓	↑	0	↑	0	↓
Wu et al., 2016 ³⁹	0	↑	--	--	--	--	--	--	--
Lin et al., (CATT trials), 2018 ⁴⁰	0	↑	↑	0	0	0	--	0	--
Domalpally et al., 2019 ⁴¹	0	↑	--	--	0	0	0	--	--
Dutheil et al., (Alienor Study), 2020 ⁴²	↑	↑	--	--	0	0	--	↑	--

Continued on next page

Source	CFH Y402H†	ARMS2	HTRA1	CFH I62V†	C3	C2/CFB‡	VEGFA	LIPC	SKIV2L
Altay et al. (EUGENDA Study), 2021 ⁴³	--	↑	--	↓	--	--	--	--	↓
Thomson et al., 2022 ⁹	0	↑	--	--	0	0	--	0	--

↑ - increased risk, ↓ - decreased risk, 0 no risk, -- SNP not investigated in study

Gene variants were only added to the list if they appeared in more than one study. Otherwise a description of findings is in the text.

†SNP rs800292 refers to CFH I62V and SNP s1061170 to CFH Y402H

‡C2 and CFB are in linkage disequilibrium. They are shown together or separately in different studies

Supplementary Table 2. Candidate approach to identify genetic associations with number of (i) reticular pseudodrusen and (ii) drusen.

	Rs identifier	chr:pos[hg19]	EA/OA (EAF)	Beta (95% CI)	P	Nearest gene	Advanced or early AMD locus
Reticular pseudodrusen							
	rs10922109	1:196704632	C/A (0.61)	0.09 (0.04; 0.13)	1.8e-04	<i>CFH</i>	Advanced
	rs3750846	10:124215565	C/T (0.21)	0.16 (0.1; 0.21)	2.1e-09	<i>ARMS2-HTRA1</i>	Advanced
Drusen							
	rs10922109	1:196704632	C/A (0.66)	0.32 (0.26; 0.39)	5.6e-25	<i>CFH</i>	Advanced
	rs10033900	4:110659067	T/C (0.48)	0.1 (0.04; 0.16)	4.5e-04	<i>CFI</i>	Advanced
	rs62358361	5:39327888	T/G (0.02)	0.52 (0.28; 0.76)	1.5e-05	<i>C9</i>	Advanced
	rs3750846	10:124215565	C/T (0.25)	0.29 (0.22; 0.35)	8.9e-19	<i>ARMS2-HTRA1</i>	Advanced
	rs2230199	19:6718387	C/G (0.22)	0.15 (0.08; 0.22)	2.5e-05	<i>C3</i>	Advanced

Supplementary Table 3. Colocalization results between GWAS findings for number of pure reticular pseudodrusen and age-related macular degeneration (Finngen) at the *CFH* and *ARMS2-HTRA1* loci. Maximum probabilities are highlighted in bold.

Gene symbol	Lead RSID	PP.H0	PP.H1	PP.H2	PP.H3	PP.H4
<i>CFH</i>	rs1061170	7.2e-269	1.3e-268	7.4e-02	0.13	0.79
<i>ARMS2-HTRA1</i>	rs11200630	-	-	1.8e-06	4.2e-03	1.00

Supplementary Table 4. Genome-wide significant associations with number of mixed reticular pseudodrusen and mixed drusen.

	Rs identifier	chr:pos[hg19]	EA/OA (EAF)	Beta (95% CI)	P	Nearest gene
Reticular pseudodrusen						
	rs9970075	1:196673430	T/G (0.49)	0.38 (0.31; 0.45)	2.3e-26	<i>CFH</i>
	1:196884765_CA_C	1:196884765	CA/C (0.91)	0.39 (0.26; 0.53)	3.8e-09	<i>CFHR4</i>
	rs201396317	10:124213674	A/C (0.24)	0.32 (0.24; 0.4)	9.9e-16	<i>ARMS2</i>
Drusen						
	rs10922102	1:196668287	C/T (0.49)	0.45 (0.39; 0.51)	6.8e-51	<i>CFH</i>
	rs141073581	1:196851265	A/G (0.96)	0.48 (0.31; 0.65)	2.1e-08	<i>CFHR4</i>
	rs9427662	1:196946775	T/C (0.92)	0.35 (0.23; 0.47)	3.4e-09	<i>CFHR5</i>
	rs429608	6:31930462	G/A (0.87)	0.27 (0.18; 0.36)	9.3e-10	<i>SKIV2L</i>
	rs61871744	10:124203787	C/T (0.27)	0.34 (0.28; 0.41)	1.2e-26	<i>ARMS2</i>

Supplementary Table 5. Genome-wide significant associations with presence or absence of pure drusen.

	Rs identifier	chr:pos[hg19]	EA/OA (EAF)	OR (95% CI)	P	Nearest gene
Drusen						
	1:196661838_TTTTC_T	1:196661838	TTTTC/T (0.44)	3.32 (2.7; 4.08)	9.3e-34	<i>CFH</i>
	rs61871744	10:124203787	C/T (0.24)	2.54 (2.06; 3.13)	2.4e-19	<i>ARMS2</i>

Supplementary Table 6 – Characterization of previous genetic studies for reticular pseudodrusen

Source	Imaging modalities							Minimal number of RPD required	Late AMD only	Genes explored	Pure RPD	Mean age
	CFP	NIR	SD-OCT	FAF	RF	FA	ICG					
Klein et al., (Beaver Dam Eye Study), 2008 ³⁰	x							NA	No	AMD partial	0	61.8
Smith et al., 2011 ³¹	x	x		x				NA	No	AMD partial	0	83
Ueda-Arakawa et al., 2013 ³²	x	x	x	x		x	x	5	Yes	AMD partial	0	73.9
Puche et al., 2013 ³³		x	x					NA	No	AMD partial	0	77.4
Joachim et al., (Blue Mountain Eye Study), 2014 ³⁴	x							NA	No	AMD partial	Unclear	69.5

Continued on next page

Source	Imaging modalities							Minimal number of RPD required	Late AMD only	Genes explored	Pure RPD	Mean age
	CFP	NIR	SD-OCT	FAF	RF	FA	ICG					
Yoneyama et al., 2014 ³⁵	x	x	x	x	x			NA	Yes	AMD partial	0	74.9
Boddu et al., 2014 ³⁶		x	x					NA	No	AMD partial	30	87*
Finger et al. (Melbourne Collaborative Cohort Study), 2016 ³⁷	x							NA	No	AMD partial	6	76
Buitendijk et al., (Rotterdam Study), 2016 ³⁸	x	x						NA	No	AMD partial	14	82.1
Wu et al., 2016 ³⁹		x	x					5	No	AMD partial	0	76.2

Continued on next page

Source	Imaging modalities							Minimal number of RPD required	Late AMD only	Genes explored	Pure RPD	Mean age
	CFP	NIR	SD-OCT	FAF	RF	FA	ICG					
Lin et al., (CATT trials), 2018 ⁴⁰	x				x	x		NA	No	AMD partial	0	78
Domalpally et al., 2019 ⁴¹	x			x				5	No	AMD	0	79
Dutheil et al., (Alienor Study), 2020 ⁴²	x	x	x	x				NA	No	AMD partial	22	81.9
Altay et al. (EUGENDA Study), 2021 ⁴³		x	x					NA	No	AMD	9	80.7
Thomson et al., 2022 ⁹		x	x	x				NA	No	AMD partial	29	81.3


AMD = age-related macular degeneration, CFP = Colour fundus photography, FA = Fluorescein angiography, FAF = Fundus autofluorescence, ICG = Indocyanine green, NIR = Near infrared photography, RF = Red-free, RPD = reticular pseudodrusen, SD-OCT = Spectral domain optical coherence tomography

* - Median age. Mean age was not provided.


AMD partial = only some of known AMD-associated genetic loci were explored

Supplementary figures

Supplementary figure S1. Locus zoom plots for lead variants from the genome-wide association study on number of reticular pseudodrusen.



Supplementary figure S2. Manhattan and quantile-quantile plots of continuous GWAS results for mixed RPD with or without drusen) and mixed drusen with or without RPD).



Supplementary figure S3. Manhattan and quantile-quantile plots of case-control GWAS results for pure RPD (in the absence of drusen) and pure drusen (in the absence of RPD)

