

Characteristic	N = 5,008
District	
Butha-Buthe	2,486 (49.6%)
Mokhotlong	2,522 (50.4%)
Number of household members, median (IQR)	3 (2, 4)
International Wealth Index, median (IQR)	26 (17, 36)
Monthly household income (USD), median (IQR)¹	42.4 (18.6, 63.6)
Unknown	7
Main source of household income	
Salary/wages ²	1,289 (25.7%)
Informal business (e.g. street vendor)	1,109 (22.1%)
Formal business (e.g. retail trade)	37 (0.7%)
Farming	1,213 (24.2%)
Rentals / interest	61 (1.2%)
Grants / pension	1,109 (22.1%)
Refused to say	190 (3.8%)
Access to health facility	
Easy access	2,460 (49.1%)
Hard access ³	2,548 (50.9%)
Difficulties accessing health care⁴	1,120 (22.4%)
Difficulties accessing medications⁵	821 (16.4%)
Household Food Insecurity Access category	
Food secure	2,385 (47.6%)
Mildly food insecure	167 (3.3%)
Moderately food insecure	965 (19.3%)
Severely food insecure	1,489 (29.7%)
Unknown	2
Drinking water	
Running (piped) water inside the dwelling	15 (0.3%)
Running (piped) water outside dwelling but inside yard	64 (1.3%)
Neighbor's tap	43 (0.9%)
Public / community tap	4,075 (81.4%)
Borehole	548 (10.9%)
Dam/ river	216 (4.3%)
Rainwater tank	41 (0.8%)
Bottled water	0 (0.0%)
Refused to say	6 (0.1%)
Toilet	
Own flush toilet	21 (0.4%)
Flush toilet shared with another household	12 (0.2%)
Own ventilated improved pit latrine	628 (12.5%)
Own pit latrine with slab	1,993 (39.8%)

Characteristic	N = 5,008
Open pit latrine	179 (3.6%)
Composting toilet	68 (1.4%)
None/Bush/Field	1,959 (39.1%)
Pit latrine shared with another household	141 (2.8%)
Refused to say	7 (0.1%)
Electricity available	663 (13.2%)
Cooking fuel	
Electricity	106 (2.1%)
LPG	208 (4.2%)
Biogas	29 (0.6%)
Paraffin	147 (2.9%)
Wood	4,210 (84.1%)
Coal	4 (0.1%)
Straw, shrubs, grass	16 (0.3%)
Animal dung	286 (5.7%)
Refused to say	2 (0.0%)
Heating fuel	
Electricity	68 (1.4%)
Gas	41 (0.8%)
Paraffin	323 (6.4%)
Wood	3,198 (63.9%)
Coal	23 (0.5%)
Animal dung	1,308 (26.1%)
Crop waste	13 (0.3%)
Refused to say	34 (0.7%)
People per bedroom	
1-2	2,527 (50.5%)
3-4	1,880 (37.5%)
5 or more	599 (12.0%)
Other / unknown / refused to say	2 (0.0%)
Wall material	
Mud	3,099 (61.9%)
Mud and cement	1,042 (20.8%)
Corrugated iron	230 (4.6%)
Bare brick	245 (4.9%)
Finished / plaster	348 (6.9%)
Other	42 (0.8%)
Refused to say	2 (0.0%)
Roof material	
Natural	3,099 (61.9%)
Rudimentary (wood planks, cardboard)	242 (4.8%)

Characteristic	N = 5,008
Finished (metal, corrugated, wood, asbestos, ceramic / clay tiles, roofing shingles)	1,548 (30.9%)
Other	116 (2.3%)
Refused to say	3 (0.1%)
Floor material	
Earth	2,511 (50.1%)
Wood planks	30 (0.6%)
Cement	1,462 (29.2%)
Vinyl tiles	203 (4.1%)
Ceramic tiles	232 (4.6%)
Carpet	549 (11.0%)
Other	19 (0.4%)
Refused to say	2 (0.0%)
Agricultural land ownership	3,127 (62.4%)
Clock	1,059 (21.1%)
Water pump	83 (1.7%)
Grain grinder	700 (14.0%)
Sewing machine	1,093 (21.8%)
Color TV	1,827 (36.5%)
Black/white TV	1,620 (32.3%)
Video player / DVD player	212 (4.2%)
Tape player / CD player	150 (3.0%)
Camera, video camera	39 (0.8%)
Refrigerator	259 (5.2%)
Freezer	66 (1.3%)
Washing machine	12 (0.2%)
Dishwasher	7 (0.1%)
Electric or gas stove	1,063 (21.2%)
Kerosene stove	1,878 (37.5%)
Mobile phone	2,767 (55.3%)
Car	135 (2.7%)
Bicycle	26 (0.5%)
Landline or fix phone	21 (0.4%)
Cattle	0 (0, 2)
Horses	0 (0, 0)
Donkeys	0 (0, 1)
Sheep	0 (0, 7)
Goats	0 (0, 0)
Chickens	0 (0, 5)
Pigs	0 (0, 0)
Rabbits	0 (0, 0)

Characteristic	N = 5,008
¹ Collected in Lesotho Loti, converted using the bilateral conversion rate on May 31, 2024 [62]	
² Salaries and wages of household members or other people (i.e. of family members living outside the household)	
³ Defined as needing to cross a mountain or river or travel >10 km to the nearest health facility.	
⁴ Defined as at least one household member not being able to consult a healthcare provider despite needing it during the last 12 months.	
⁵ Defined as at least one household member experiencing difficulties obtaining required medication.	

Supplementary table 1 Overview of all collected household level socioeconomic indicators. IQR: interquartile range

Characteristic	Female, n(%) 4560 (58%)	Male, n(%) 3314 (42%)	Total, n(%) 7874 (100%)
Age (years), median (IQR)	41 (27, 60)	39 (28, 54)	40 (27, 58)
18-39 years	2,148 (47.1%)	1,662 (50.2%)	3,810 (48.4%)
40-64 years	1,529 (33.5%)	1,185 (35.8%)	2,714 (34.5%)
≥65 years	883 (19.4%)	467 (14.1%)	1,350 (17.1%)
District			
Butha-Buthe	2,283 (50.1%)	1,592 (48.0%)	3,875 (49.2%)
Mokhotlong	2,277 (49.9%)	1,722 (52.0%)	3,999 (50.8%)
Relationship status			
Single	655 (14.4%)	902 (27.2%)	1,557 (19.8%)
In a committed relationship	2,690 (59.0%)	2,024 (61.1%)	4,714 (59.9%)
Separated or divorced	87 (1.9%)	123 (3.7%)	210 (2.7%)
Widowed	1,119 (24.5%)	245 (7.4%)	1,364 (17.3%)
Unknown/refused to say	9 (0.2%)	20 (0.6%)	29 (0.4%)
Education			
No schooling	276 (6.1%)	684 (20.7%)	960 (12.2%)
Primary school	2,623 (57.5%)	1,879 (56.7%)	4,502 (57.2%)
Secondary/high school	1,448 (31.8%)	624 (18.8%)	2,072 (26.3%)
Tertiary/higher/post high school	209 (4.6%)	121 (3.7%)	330 (4.2%)
Unknown/refused to say	3 (0.1%)	4 (0.1%)	7 (0.1%)
Missing	1	2	3
Work status			
Working for pay	444 (9.7%)	572 (17.3%)	1,016 (12.9%)
Self-employed	703 (15.4%)	709 (21.4%)	1,412 (17.9%)
Working on own plot or looking after livestock	469 (10.3%)	1,075 (32.4%)	1,544 (19.6%)
Helping another family member with their business, without pay	33 (0.7%)	30 (0.9%)	63 (0.8%)
Full-time student	90 (2.0%)	38 (1.1%)	128 (1.6%)
Homemaker (looking after children/others/home)	1,734 (38.0%)	184 (5.6%)	1,918 (24.4%)
Long term sick or disabled	53 (1.2%)	41 (1.2%)	94 (1.2%)
Retired	205 (4.5%)	136 (4.1%)	341 (4.3%)
Unemployed	702 (15.4%)	461 (13.9%)	1,163 (14.8%)
Unknown/refused to say	127 (2.8%)	68 (2.1%)	195 (2.5%)
Body mass index (kg/m²)			
Underweight (<18.5)	250 (5.5%)	383 (11.7%)	633 (8.1%)
Normal weight (18.5–24.9)	1,678 (37.2%)	2,165 (66.1%)	3,843 (49.4%)
Overweight (25.0-29.9)	1,323 (29.3%)	548 (16.7%)	1,871 (24.0%)
Obese (≥30)	1,259 (27.9%)	181 (5.5%)	1,440 (18.5%)
Missing	50	37	87
Abdominal circumference (cm)	86 (78, 97)	80 (75, 87)	83 (76, 93)
Missing	400	368	768

Characteristic	Female, n(%) 4560 (58%)	Male, n(%) 3314 (42%)	Total, n(%) 7874 (100%)
Physical activity			
Low physical activity	625 (13.8%)	360 (11.0%)	985 (12.6%)
Moderate physical activity	701 (15.5%)	345 (10.5%)	1,046 (13.4%)
High physical activity	3,201 (70.7%)	2,575 (78.5%)	5,776 (74.0%)
Missing	33	34	67
Fruit intake (servings/day)			
	0.14 (0.00, 0.86)	0.14 (0.00, 0.86)	0.14 (0.00, 0.86)
Missing	1	2	3
Vegetable intake (servings/day)			
	1.14 (0.57, 2.14)	0.86 (0.29, 2.00)	1.00 (0.43, 2.00)
Missing	1	2	3
Portions of vegetables and fruits/day			
<5	4,043 (88.7%)	2,962 (89.4%)	7,005 (89.0%)
≥5	516 (11.3%)	350 (10.6%)	866 (11.0%)
Missing	1	2	3
Adding salt during cooking			
Always	3,008 (66.0%)	2,226 (67.2%)	5,234 (66.5%)
Often	130 (2.9%)	123 (3.7%)	253 (3.2%)
Sometimes	773 (17.0%)	523 (15.8%)	1,296 (16.5%)
Rarely	509 (11.2%)	360 (10.9%)	869 (11.0%)
Never	131 (2.9%)	68 (2.1%)	199 (2.5%)
Unknown/refused to say	8 (0.2%)	12 (0.4%)	20 (0.3%)
Missing	1	2	3
Adding salt while eating			
Always	631 (13.8%)	523 (15.8%)	1,154 (14.7%)
Often	206 (4.5%)	219 (6.6%)	425 (5.4%)
Sometimes	1,295 (28.4%)	1,028 (31.0%)	2,323 (29.5%)
Rarely	1,246 (27.3%)	1,011 (30.5%)	2,257 (28.7%)
Never	1,175 (25.8%)	521 (15.7%)	1,696 (21.5%)
Unknown/refused to say	6 (0.1%)	10 (0.3%)	16 (0.2%)
Missing	1	2	3
Importance of lowering salt intake			
Very important	4,000 (87.7%)	2,697 (81.4%)	6,697 (85.1%)
Somewhat important	145 (3.2%)	164 (5.0%)	309 (3.9%)
Not at all important	110 (2.4%)	133 (4.0%)	243 (3.1%)
Unknown/refused to say	304 (6.7%)	318 (9.6%)	622 (7.9%)
Missing	1	2	3
Perceiving high salt intake as potential health problem			
Yes	3,775 (82.8%)	2,504 (75.6%)	6,279 (79.8%)
No	519 (11.4%)	502 (15.2%)	1,021 (13.0%)
Unknown/refused to say	265 (5.8%)	306 (9.2%)	571 (7.3%)
Missing	1	2	3

Characteristic	Female, n(%) 4560 (58%)	Male, n(%) 3314 (42%)	Total, n(%) 7874 (100%)
Eating processed food high in salt			
Always	380 (8.3%)	298 (9.0%)	678 (8.6%)
Often	217 (4.8%)	223 (6.7%)	440 (5.6%)
Sometimes	1,460 (32.0%)	1,048 (31.6%)	2,508 (31.9%)
Rarely	1,921 (42.1%)	1,383 (41.8%)	3,304 (42.0%)
Never	560 (12.3%)	346 (10.4%)	906 (11.5%)
Unknown/refused to say	21 (0.5%)	14 (0.4%)	35 (0.4%)
Missing	1	2	3
Perceived quantity of salt intake			
Far too much	86 (1.9%)	78 (2.4%)	164 (2.1%)
Too much	193 (4.2%)	186 (5.6%)	379 (4.8%)
Just the right amount	2,879 (63.1%)	2,108 (63.6%)	4,987 (63.4%)
Too little	1,177 (25.8%)	786 (23.7%)	1,963 (24.9%)
Far too little	140 (3.1%)	63 (1.9%)	203 (2.6%)
Unknown/refused to say	84 (1.8%)	91 (2.7%)	175 (2.2%)
Missing	1	2	3
Fruit juices			
None per week	2,745 (60.2%)	2,172 (65.6%)	4,917 (62.5%)
One to two times a week	1,458 (32.0%)	942 (28.4%)	2,400 (30.5%)
Three to four times a week	225 (4.9%)	105 (3.2%)	330 (4.2%)
Five or more times a week	118 (2.6%)	78 (2.4%)	196 (2.5%)
Refused to say	13 (0.3%)	15 (0.5%)	28 (0.4%)
Missing	1	2	3
Fizzy sugar sweetened beverages			
None per week	3,169 (69.5%)	2,115 (63.9%)	5,284 (67.1%)
One to two times a week	1,227 (26.9%)	1,020 (30.8%)	2,247 (28.5%)
Three to four times a week	122 (2.7%)	131 (4.0%)	253 (3.2%)
Five or more times a week	23 (0.5%)	32 (1.0%)	55 (0.7%)
Refused to say	18 (0.4%)	14 (0.4%)	32 (0.4%)
Missing	1	2	3
Candy			
None per week	2,506 (55.0%)	2,082 (62.9%)	4,588 (58.3%)
One to two times a week	1,624 (35.6%)	971 (29.3%)	2,595 (33.0%)
Three to four times a week	299 (6.6%)	174 (5.3%)	473 (6.0%)
Five or more times a week	115 (2.5%)	70 (2.1%)	185 (2.4%)
Refused to say	15 (0.3%)	15 (0.5%)	30 (0.4%)
Missing	1	2	3
Biscuits			
None per week	3,215 (70.5%)	2,454 (74.1%)	5,669 (72.0%)
One to two times a week	1,135 (24.9%)	707 (21.3%)	1,842 (23.4%)

Characteristic	Female, n(%) 4560 (58%)	Male, n(%) 3314 (42%)	Total, n(%) 7874 (100%)
Three to four times a week	140 (3.1%)	107 (3.2%)	247 (3.1%)
Five or more times a week	55 (1.2%)	28 (0.8%)	83 (1.1%)
Refused to say	14 (0.3%)	16 (0.5%)	30 (0.4%)
Missing	1	2	3
Chocolate			
None per week	3,767 (82.6%)	2,796 (84.4%)	6,563 (83.4%)
One to two times a week	671 (14.7%)	427 (12.9%)	1,098 (13.9%)
Three to four times a week	87 (1.9%)	57 (1.7%)	144 (1.8%)
Five or more times a week	20 (0.4%)	17 (0.5%)	37 (0.5%)
Refused to say	14 (0.3%)	15 (0.5%)	29 (0.4%)
Missing	1	2	3
Crisps			
None per week	2,072 (45.4%)	1,760 (53.1%)	3,832 (48.7%)
One to two times a week	1,847 (40.5%)	1,178 (35.6%)	3,025 (38.4%)
Three to four times a week	456 (10.0%)	258 (7.8%)	714 (9.1%)
Five or more times a week	168 (3.7%)	103 (3.1%)	271 (3.4%)
Refused to say	16 (0.4%)	13 (0.4%)	29 (0.4%)
Missing	1	2	3
Fries/chips			
None per week	3,036 (66.6%)	2,232 (67.4%)	5,268 (66.9%)
One to two times a week	1,254 (27.5%)	884 (26.7%)	2,138 (27.2%)
Three to four times a week	204 (4.5%)	134 (4.0%)	338 (4.3%)
Five or more times a week	50 (1.1%)	47 (1.4%)	97 (1.2%)
Refused to say	15 (0.3%)	15 (0.5%)	30 (0.4%)
Missing	1	2	3
Makoonya (traditional fried bread bun)			
None per week	2,260 (49.6%)	1,638 (49.5%)	3,898 (49.5%)
One to two times a week	1,802 (39.5%)	1,350 (40.8%)	3,152 (40.0%)
Three to four times a week	366 (8.0%)	218 (6.6%)	584 (7.4%)
Five or more times a week	117 (2.6%)	92 (2.8%)	209 (2.7%)
Refused to say	14 (0.3%)	14 (0.4%)	28 (0.4%)
Missing	1	2	3
Meat			
None per week	1,952 (42.8%)	1,303 (39.3%)	3,255 (41.4%)
One to two times a week	2,024 (44.4%)	1,481 (44.7%)	3,505 (44.5%)
Three to four times a week	458 (10.0%)	388 (11.7%)	846 (10.7%)
Five or more times a week	110 (2.4%)	128 (3.9%)	238 (3.0%)
Refused to say	15 (0.3%)	12 (0.4%)	27 (0.3%)
Missing	1	2	3
Smoking status			

Characteristic	Female, n(%) 4560 (58%)	Male, n(%) 3314 (42%)	Total, n(%) 7874 (100%)
Current Smoker	1,015 (22.3%)	1,579 (47.9%)	2,594 (33.1%)
Never Smoker	3,500 (77.0%)	1,657 (50.2%)	5,157 (65.8%)
Ex-smoker quit smoking less than 5 years ago	21 (0.5%)	35 (1.1%)	56 (0.7%)
Ex-smoker quit smoking 5 years ago or longer	8 (0.2%)	28 (0.8%)	36 (0.5%)
Missing	16	15	31
Frequency of alcohol consumption			
Every day/ 7 days per week	71 (1.6%)	158 (4.8%)	229 (2.9%)
5 to 6 days per week	43 (0.9%)	102 (3.1%)	145 (1.8%)
3 to 4 days per week	93 (2.0%)	183 (5.5%)	276 (3.5%)
1 to 2 days per week	287 (6.3%)	505 (15.3%)	792 (10.1%)
1 to 3 days per month	139 (3.1%)	245 (7.4%)	384 (4.9%)
Less than 1 to 3 days per month	180 (4.0%)	195 (5.9%)	375 (4.8%)
Never	3,729 (82.1%)	1,914 (58.0%)	5,643 (71.9%)
Missing	18	12	30
First-degree relative with hypertension	976 (21.4%)	374 (11.3%)	1,350 (17.1%)
First-degree relative with diabetes	257 (5.6%)	90 (2.7%)	347 (4.4%)
History of myocardial infarction	62 (1.4%)	11 (0.3%)	73 (0.9%)
History of chronic kidney disease	16 (0.4%)	5 (0.2%)	21 (0.3%)
Current or previous TB diagnosis	169 (3.7%)	238 (7.2%)	407 (5.2%)
Severe vision impairment/blindness	35 (0.8%)	7 (0.2%)	42 (0.5%)
Peripheral arterial disease	27 (0.6%)	5 (0.2%)	32 (0.4%)
Heart failure	32 (0.7%)	5 (0.2%)	37 (0.5%)
History of stroke	20 (0.4%)	13 (0.4%)	33 (0.4%)
History of neuropathy	39 (0.9%)	8 (0.2%)	47 (0.6%)
Chronic lung disease (COPD/asthma)	16 (0.4%)	11 (0.3%)	27 (0.3%)
Diabetic foot syndrome	6 (0.1%)	1 (0.0%)	7 (0.1%)
Self-reported HIV status			
HIV positive	782 (17.1%)	406 (12.3%)	1,188 (15.1%)
Tested negative ≤ 12 months ago	2,435 (53.4%)	1,503 (45.4%)	3,938 (50.0%)
Tested negative > 12 months ago	846 (18.6%)	684 (20.6%)	1,530 (19.4%)
Unknown	453 (9.9%)	695 (21.0%)	1,148 (14.6%)
Refused to say	44 (1.0%)	26 (0.8%)	70 (0.9%)
ART intake among those HIV positive (n=1188)			
Yes	771 (98.6%)	398 (98.0%)	1,169 (98.4%)
No	11 (1.4%)	8 (2.0%)	19 (1.6%)
Missing	3,778	2,908	6,686
Lipid lowering medication			
Yes	17 (0.4%)	3 (0.1%)	20 (0.3%)
No	4,543 (99.6%)	3,311 (99.9%)	7,854 (99.7%)
Aspirine			

Characteristic	Female, n(%)	Male, n(%)	Total, n(%)
	4560 (58%)	3314 (42%)	7874 (100%)
Yes	386 (8.5%)	103 (3.1%)	489 (6.2%)
No	4,174 (91.5%)	3,211 (96.9%)	7,385 (93.8%)

Supplementary table 2 Overview of complete individual level demographic, clinical and behavioural risk factor data of adult participants.