perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license .

# The impact of exercise training on the brain and cognition in T2DM, and its physiological mediators: a systematic review

- 3
- 4 Jitske Vandersmissen<sup>1</sup>, Ilse Dewachter<sup>2</sup>, Koen Cuypers<sup>1,3\*</sup>, Dominique Hansen<sup>1,4\*</sup>
- REVAL Rehabilitation Research Center, Faculty of Rehabilitation Sciences, Hasselt University, 3590
   Diepenbeek, Belgium. Jitske.vandersmissen@uhasselt.be
- 7 2. Biomedical Research Institute, BIOMED, Hasselt University, 3590 Diepenbeek, Belgium.
- Movement Control and Neuroplasticity Research Group, Department of Movement Sciences, Group
   Biomedical Sciences, KU Leuven, 3000 Leuven, Belgium.
- 10 4. Heart Centre Hasselt, Jessa Hospital, 3500 Hasselt, Belgium.
- 11 \* Shared authorship

# 12 Corresponding author:

- 13 Jitske Vandersmissen Jitske.vandersmissen@uhasselt.be
- 14 REVAL Rehabilitation Research Centre, Faculty of Rehabilitation Sciences, Hasselt University
- 15 Wetenschapspark 7, 3590 Diepenbeek, Belgium
- 16 Abstract
- 17 Background: Type 2 diabetes (T2DM) affects brain structure and function, and is associated with an
- 18 increased risk of dementia and mild cognitive impairment. It is known that exercise training has a
- 19 beneficial effect on cognition and the brain, at least in healthy people, but the impact of exercise
- 20 training on cognition and the brain remains to be fully elucidated in patients with T2DM.

21 Methods: This paper systematically reviews studies that evaluate the effect of exercise training on 22 cognition in T2DM, and aims to indicate the most beneficial exercise modality for improving or 23 preserving cognition in this patient group. In addition, the possible physiological mediators and targets 24 involved in these improvements are narratively described in the second part of this review. Papers 25 published up until the end of June 2024 were searched by means of the electronic database PubMed. 26 Studies directly investigating the effect of any kind of exercise training on the brain or cognition in 27 patients with T2DM, or animal models thereof, were included, with the exception of human studies 28 assessing cognition only at one time point, and studies combining exercise training with other 29 interventions (e.g. dietary changes, cognitive training, etc.).

- Results: For the systematic part of the review, 24 papers were found to be eligible. 20 out of 24 papers 30 31 (83.3%) showed a significant positive effect of exercise training on cognition in T2DM, of which four 32 studies only showed a moderate significant effect. Four papers (16.7%) did not show a significant effect 33 of exercise on cognition in T2DM, but two of them did show a positive trend. Similar effects were found 34 for resistance and endurance exercise, with both possibly requiring a minimal intensity to reach 35 cognitive improvement. In addition, BDNF, lactate, leptin, adiponectin, GSK3β, GLP-1, the AMPK/SIRT1 36 pathway, and the PI3K/Akt pathway were identified as plausible mediators directly from studies 37 investigating the effect of exercise training on the brain in T2DM.
- **Conclusion:** Overall, exercise training beneficially affects cognition and the brain in T2DM, with resistance and endurance exercise having similar effects. However, additional studies comparing the effect of different exercise intensities are needed to determine the optimal exercise intensity for cognitive improvement. Furthermore, we were able to define several mediators involved in the effect of exercise training on cognition in T2DM, but further research is necessary to unravel the entire NOTE: This preprint reports new research that has not been certified by peer review and should not be used to guide clinical practice.

This review demonstrates a positive effect of exercise training on the brain in T2DM, but 44

45 simultaneously emphasizes the need for additional studies on this topic. BDNF, lactate, leptin, 46 adiponectin, GSK3B, GLP-1, the AMPK/SIRT1 pathway, and the PI3K/Akt pathway were identified as

47 factors mediating the effect of exercise on the brain in T2DM.

# 48 1. Introduction

## 49 1.1 Cognitive decline in aging

50 Cognitive decline is inherent to aging. As we age, synaptic plasticity reduces, neural mitochondrial 51 function declines, epigenetic changes occur, etc., all contributing to decreased cognitive abilities [1-5]. 52 Especially fluid intelligence, defined as the ability to solve problems, process new information, and 53 learn new things, declines with aging. Fluid intelligence consists of several domains, including executive 54 function, psychomotor ability, memory, and processing speed [6, 7].

Aging is associated with a decline in both grey and white matter volume, accompanied by ventricular enlargement and cortical thinning, as well as a decline in brain function [7, 8]. Amyloid β plaques are not only found in Alzheimer's disease (AD) patients, but also in 20-30% of healthy adults, and are speculated to contribute to this neuronal loss [6, 9]. Also neuronal volume and the number of neuronal connections decreases due to a decline in the complexity of dendrite arborization, decreased neuritic spines, and reduced dendrite length [6, 10]. In addition, demyelination and decreased white matter integrity contribute to functional impairment of the brain [8].

## 62 1.2 Cognitive dysfunction in T2DM

63 T2DM patients are prone to more severe cognitive decline than what occurs during normal aging. A 64 large fraction (45%) of T2DM patients experience cognitive dysfunction and have an increased risk of 65 dementia [11-16]. Subjects with diabetes have a relative risk of 1.46 for AD, and a relative risk of 1.51 for any form of dementia [17]. Especially executive functioning, memory, attention, and information 66 processing speed have been shown to be affected in T2DM [14, 18, 19]. This cognitive impairment is 67 68 related to the affected brain structure and metabolism in T2DM. Decreased regional grey matter 69 volume as well as altered intrinsic activity in the default mode network have been demonstrated in 70 these patients [20-22]. Multiple pathophysiologies have been suggested as the cause of this, including 71 cardiovascular complications, chronic low-grade inflammation, and hyperglycemia [14, 15, 18].

72 Several studies have demonstrated a link between T2DM and AD, also called type 3 diabetes. These 73 studies focus on the fact that insulin signalling also plays an important role in the brain, and that T2DM 74 and AD seem to share a number of pathophysiological processes such as amyloid  $\beta$  plaques, disturbed 75 cerebral glucose metabolism, tau hyperphosphorylation, and inflammation [18, 23-26]. Insulin and 76 insulin-like-growth-factor (IGF)-1 are important for neuronal survival and brain function. Numerous 77 insulin receptors (IR) and IGF-1 receptors (IGF-1R) can be found in the brain, with a large amount being 78 present in the hippocampus. One of the major downstream pathways of the IR is the PI3K/Akt 79 pathway, which plays an important role in the regulation of brain function, and in the inactivation of 80 GSK3 $\beta$ , which is an enzyme able to phosphorylate tau at pathological tau epitopes [23, 27, 28]. 81 Reduced cerebral IR activation and insulin levels have been demonstrated in AD [25, 29], highlighting 82 similar processes in AD and T2DM. Moreover, insulin-degrading enzyme is capable of clearing amyloid 83  $\beta$  peptides that aggregate into amyloid plaques, a key pathological hallmark of AD. In T2DM, 84 competitive binding of high insulin concentrations and amyloid  $\beta$  peptides with this enzyme could be 85 involved in decreased amyloid  $\beta$  clearance, thereby potentially contributing to increased accumulation 86 of amyloid  $\beta$  peptides, and formation of amyloid  $\beta$  plaques [15, 30, 31].

#### 87 1.3 Positive effect of exercise on cognitive function

In healthy people, exercise has a positive effect on cognitive function [32-34]. This effect is mediated
 by multiple processes. Muscle contraction during exercise releases myokines, which stimulate the
 production of neurotrophic factors such as brain-derived neurotrophic factor (BDNF), promoting

perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license .

neurogenesis. Additionally, the exercise-induced release of anti-inflammatory factors contributes to
balancing the brain's redox status, counteracting many pathological processes [35-38]. Research shows
that exercise reduces the age-related decrease in hippocampal volume [39-41], which, in combination
with increased levels of neurotrophic factors, contributes to the maintenance of memory and
neuroplasticity [36, 42, 43]. Moreover, there is a positive association between cardiorespiratory fitness
levels and whole-brain and white matter volume in patients with early-stage AD [44]. In this way,
exercise reduces the risk of dementia and neurodegenerative diseases [36, 45-47].

98 Lactate has been identified as an important contributor to the positive effect of exercise on cognitive 99 function. During anaerobic exercise, lactate is released from contracting muscles, eventually leading 100 to increased lactate levels in the brain after crossing the blood-brain-barrier (BBB). This results in an 101 increased expression of brain plasticity genes such as BDNF, Arc, and c-fos, contributing to 102 neurogenesis [48-50]. In addition, lactate also binds to hydroxycarboxylic acid receptor 1 (HCAR1), 103 which leads to an increase in vascular endothelial growth factor (VEGF), stimulating angiogenesis and 104 thereby contributing to increased cognitive function [50-52]. Also the myokine irisin has been shown 105 to positively influence cognition. Synaptic plasticity and memory in amyloid pathology mimicking AD 106 mouse models can be rescued by boosting brain levels of FNDC5/irisin, and peripheral overexpression 107 of FNDC5/irisin rescues memory impairment [53]. Cathepsin b is another myokine which has been 108 associated with improved cognition. It is known to play a role in both neurogenesis and angiogenesis 109 [54, 55]. Cathepsin b levels increase in mouse and human plasma in response to exercise, which is positively correlated with memory, while cathepsin b-knockout mice do not show any cognitive 110 111 improvements following running exercise [56]. This suggests that cathepsin b is mandatory for the 112 positive effect of exercise on cognition.

113 1.4 The association of physical activity with cognitive function in T2DM

Because of the above-mentioned positive effects of exercise on cognitive function, and to get more insight into whether this translates to T2DM, some studies have explored the association between physical activity (PA) and cognition in T2DM.

117 For example, one study found significantly higher cognitive scores in an active group of T2DM patients compared to a sedentary group of T2DM patients [57]. They also found a significant negative 118 correlation (r = -0.2, P = 0.03) between cognition and BMI in the sedentary group, and a significant 119 120 positive correlation (r = 0.55, P = 0.01) between cognition and minutes of weekly exercise in the active group. Another study found that active elderly T2DM patients had a significantly slower rate of 121 122 cognitive decline compared to non-active/sedentary elderly T2DM patients, more specifically in the 123 global cognition (p = 0.005), executive functioning (p = 0.014), and attention/working memory (p = 0.014) 124 0.01) domains [58]. These findings were confirmed by multiple other studies [59-61].

- 125 In a study investigating the influence of the BDNF Val66Met polymorphism on the association between 126 cognition and PA in diabetic patients, it was found that carriers of the Met-allele showed significantly 127 higher scores of words recall (p < 0.001), mental status (p = 0.004), and total cognition, (p = 0.04), and 128 had a significantly higher education level. Overall, PA was associated with better total cognition, words 129 recall, and mental status, no matter the intensity. This association was strongest in the Met/Met carrier 130 group, and stronger for females than for males within this group. They also found that light to moderate PA showed greater association with cognitive domains than moderate to vigorous PA. These 131 findings thus suggest that female Met/Met diabetic carriers cognitively benefit the most from PA, and 132 133 that light to moderate PA has the most influence on the brain [61].
- Another study examined the individual and joint influence of diabetes status, apolipoprotein E (APOE)
   ε4, and PA on the risk of dementia and cognitive impairment without dementia (CIND) in cognitively

perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license .

- 136 normal older adults. The risk of dementia and CIND was higher in diabetic patients and APOE  $\varepsilon$ 4 carriers
- 137 who reported low levels of moderate to vigorous PA. The risk was even found to be nearly 10-fold
- higher for physically inactive diabetic APOE ɛ4 carriers. Higher levels of PA were associated with lower 138
- 139 dementia/CIND risk [60].

140 Overall, these findings suggest that higher levels of PA are associated with better cognitive function and less cognitive decline in T2DM. However, based on these studies alone, one cannot assume 141 142 causality. In addition, there is a possibility of reverse causality, where reduced cognitive function contributes to lower levels of PA. Therefore, several studies have investigated the acute effect of 143 144 exercise on cognitive function in T2DM.

145 1.5 The acute effect of exercise on cognitive function in T2DM

146 The acute effect of exercise on cognition in T2DM has mainly been investigated in the domain of 147 executive function.

148 One study found that both endurance and resistance training acutely improved inhibitory control and 149 response time in T2DM patients [62]. In addition, moderate-intensity integrated concurrent exercise 150 (ICE), consisting of both endurance and resistance training, acutely improved all three aspects of 151 executive function (inhibition, conversion and refresh function) in cognitively normal hospitalised T2DM patients, while resistance exercise improved inhibition, and endurance exercise only caused 152 153 significant improvements in the refresh function. The ICE-induced improvements in executive function were accompanied by a simultaneous increase in cerebral blood flow in the dorsolateral prefrontal 154 155 cortex (DLPFC), the frontal pursuit area (FPA), and the orbitofrontal cortex (OFC), while resistance 156 exercise showed a corresponding activation of DLPFC and FPA, and endurance exercise increased 157 cerebral perfusion in the FPA, OFC and Broca region [63]. These results suggest that an exercise-158 induced increase in cerebral blood flow in certain areas can elicit specific cognitive improvements. This 159 knowledge can be used to customise exercise programs in T2DM patients in order to obtain the desired 160 effect.

161 These studies already give some insight into the effect of exercise on cognition in T2DM. However, 162 with the aim to incorporate exercise in T2DM treatment for cognitive improvement, it is necessary to 163 know the effect of *long-term exercise training* on the brain, which can be different from the acute 164 effect. Therefore, this review will focus on pre-post intervention studies. This review only describes RCTs investigating the effect of exercise training on cognition and brain structure in T2DM and animal 165 166 models thereof, with special emphasis on the mediators and targets involved in this. So far, this is the first review of this kind. The knowledge provided by this review could more reliably provide insights in 167 the mechanisms of cognitive decline and cognitive improvement in T2DM, and could be used to design 168 169 more specific exercise programs to target the brain in T2DM patients.

170 2. Methods

171 This review consists of two parts, the first of which is a systematic review of the papers investigating the effect of exercise training on cognition in T2DM. The second part narratively describes studies that 172 173 show beneficial brain changes or brain changes in combination with cognitive improvement in T2DM 174 as a result of exercise training, and aims to identify the involved mediators by pooling studies with 175 similar findings.

#### 176 2.1 Search methods

177 The papers described in the systematic part of this review were found by entering the following search 178 string into PubMed: (exercise OR "physical activity" OR "aerobic exercise" OR "aerobic training" OR

"resistance exercise" OR "resistance training" OR "combined exercise" OR "combined training" OR 179 "endurance exercise" OR "endurance training") AND (T2DM OR diabetes OR "type 2 diabetes") AND 180 (memory OR "executive function\*" OR "processing speed" OR attention OR brain OR cognition OR 181 "cognitive function"). The filters "randomised controlled trial" and "clinical trial" were applied. 182 183 Relevant related articles suggested by PubMed when accessing articles found by entering the aforementioned search string were also considered. Reference lists of reviews in line with the present 184 one [64-72] were examined as well. The current review discusses articles published up until the end of 185 186 June 2024. Duplicate articles and studies of poor quality were removed (Fig. 1). For the narrative part of the review, only studies directly investigating the effect of exercise on the brain in T2DM were 187

188 considered to identify plausible mediators and pathways.



# 189

190 Fig. 1 Prisma flow chart of study selection procedure. Reason 1: Does not investigate the effect of exercise training; Reason 191 2: Does not investigate the effect of exercise training on cognition; Reason 3: Does not investigate the effect of exercise 192 training on cognition in T2DM; Reason 4: Not conducted in T2DM; Reason 5: Does not investigate the effect on cognition; 193 Reason 6: Does not investigate the effect on cognition in T2DM; Reason 7: Does not investigate the effect of exercise training 194 in T2DM; Reason 8: Does not investigate the effect of exercise training on its own (diet, cognitive training, ...); Reason 9: Article 195 not available in English. RCT = randomised controlled trial; T1DM = type 1 diabetes. From: Page MJ, McKenzie JE, Bossuyt PM, 196 Boutron I, Hoffmann TC, Mulrow CD, et al. The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. 197 BMJ 2021;372:n71. doi: 10.1136/bmj.n71.

#### 198 2.2 Eligibility

199 The PICO (population, intervention, comparison, outcome) method was used to determine whether 200 articles were eligible for inclusion in this review. In order to be included in the systematic part of this review, the articles had to be written in English, and had to investigate the effect of any kind of exercise 201 202 training on cognition in T2DM patients or T2DM animal models. Articles discussing only type 1 diabetes 203 were excluded, as well as articles only investigating depression, anxiety, or other mental disorders in 204 T2DM. Also studies investigating the effect of exercise training in combination with another 205 intervention (e.g. dietary changes, cognitive training, etc.) were excluded. Studies in which cognitive function was only assessed at one time point in human subjects were excluded since these make it 206 impossible to accurately assess the effect of exercise training on cognition (pre- and post-intervention 207 208 assessments mandatory).

perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license .

#### 209 2.3 Quality control

210 Quality control of the included human studies was conducted by means of the PEDro scale, which assesses internal validity [73]. With this tool, a maximal score of 10 can be reached, indicating optimal 211 212 methodological quality. However, it was impossible for the majority of the studies included in this review to reach a maximal score of 10, since in exercise interventions it is not feasible to blind the 213 214 participants and the therapists who administered the therapy. Because of this, a score of >4/10 instead 215 of >6/10 was required for inclusion. Quality of the animal studies was assessed by means of the 216 Collaborative Approach to Meta-Analysis and Review of Animal Data from Experimental Studies (CAMARADES) checklist [74]. Animal studies with a score of <6/10 were excluded. If nothing was 217 218 mentioned about the criterion, it was indicated as not fulfilled.

# 219 Table 1 PEDro characteristics of included human studies

	1	2	3	4	5	6	7	8	9	10	11	Total
Leischik et al. (2021)	V	V	٧	٧	Х	Х	Х	٧	٧	٧	٧	7/10
Anderson- Hanley et al. (2012)	V	V	х	v	Х	Х	Х	Х	V	V	٧	5/10
Ploydang et al. (2023)	V	٧	Х	V	Х	Х	Х	٧	٧	٧	٧	6/10
Wang et al. (2023)	V	٧	V	v	Х	Х	٧	Х	٧	٧	V	7/10
Zhao et al. (2022)	V	v	٧	Х	٧	٧	٧	٧	٧	٧	٧	9/10
Yamamoto et al. (2021)	V	٧	Х	v	Х	Х	Х	٧	٧	v	٧	6/10
Teixeira et al. (2019)	V	٧	Х	٧	Х	Х	Х	٧	٧	Х	٧	5/10
Silveira- Rodrigues et al. (2021)	V	х	Х	V	Х	Х	Х	٧	V	V	٧	5/10
Espeland et al. (2017)	V	٧	Х	٧	Х	٧	٧	٧	٧	V	٧	8/10
Silveira- Rodrigues et al. (2023)	V	Х	Х	v	Х	Х	Х	٧	V	V	٧	5/10
Ghodrati et al. (2023)	V	V	Х	٧	Х	Х	Х	٧	٧	٧	٧	6/10
Martinez- Velilla et al. (2021)	٧	٧	х	V	Х	Х	٧	٧	V	V	٧	7/10
Callisaya et al. (2017)	V	v	٧	٧	٧	Х	٧	٧	٧	٧	٧	8/10
Ghahfarrokhi et al. (2024)	V	V	٧	٧	Х	Х	v	٧	٧	٧	٧	8/10
Shellington et al. (2018)	V	V	٧	V	Х	Х	Х	Х	٧	٧	٧	5/10
Chen et al. (2023)	V	٧	Х	v	Х	Х	٧	٧	٧	٧	٧	7/10
Cai et al. (2019)	V	Х	х	٧	Х	Х	Х	٧	٧	٧	٧	5/10
Coelho Junior et al. (2018)	٧	Х	Х	v	Х	Х	Х	٧	٧	٧	٧	5/10

220

perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license .

221 v = fulfilled, X = not fulfilled, 1 = Eligibility criteria specified, 2 = Randomisation, 3 = Allocation concealed, 4 = Groups similar

at baseline, 5 = Blinding of subjects, 6 = blinding of therapists, 7 = Blinding of assessors, 8 = Measures obtained for more

than 85% of initial subjects, 9 = Intention-to-treat analysis, 10 = Between-group statistical comparisons, 11 = point measures
 and measures of variability

## 225 Table 2 CAMARADES characteristics of included animal studies

	1	2	3	4	5	6	7	8	9	10	Total
Parsa et al. (2021)	٧	٧	٧	Х	٧	٧	٧	Х	٧	V	8/10
Shekarchian et al. (2023)	٧	٧	٧	х	Х	٧	٧	х	Х	V	6/10
Jesmin et al. (2022)	٧	٧	٧	х	Х	٧	٧	Х	Х	V	6/10
Lang et al. (2020)	٧	٧	٧	V	٧	٧	٧	х	٧	V	9/10
Shima et al. (2017)	٧	٧	٧	х	Х	٧	٧	х	٧	V	7/10
Shima et al. (2023)	٧	V	Х	Х	Х	٧	٧	Х	٧	V	6/10

226

227 v = fulfilled, X = not fulfilled, 1 = Published in peer-reviewed journal, 2 = Control of temperature, 3 = Randomisation, 4 = Control of temperature, 3 = Randomisation, 3 = Randomisation,

Allocation concealed, 5 = Blinding of assessors, 6 = No anaesthetics with marked intrinsic properties, 7 = Use of animals with
 diabetes, 8 = Sample size calculation, 9 = Compliance with regulatory requirements, 10 = Statement regarding conflict of
 interest

230 Interest

#### 231 **3. Results**

## 232 3.1 Systematic review on the effect of exercise training on cognition in T2DM

#### 233 <u>3.1.1 Endurance exercise training</u>

234

# 235 Human studies

236 237 Endurance exercise as simple as walking can already significantly improve cognition in T2DM patients. 238 In a study by Leischik et al. (2021), T2DM patients were randomised into a walking group, pedometer 239 group or control group. The walking group had to walk for 40 min, 3 times a week. The pedometer 240 group had to reach 10 000 steps per day. Both the walking group and the pedometer group, the latter 241 of which managed to reach an average of 8700 steps per day, showed significant improvements in 242 attention and non-verbal memory. Verbal memory also significantly improved in the walking group 243 [75]. Besides normal walking, also aquatic Nordic walking performed for 60 min 3 times a week for 12 244 weeks, has proven to cause significantly increased Montreal Cognitive Assessment (MoCA) scores in 245 T2DM patients. However, no significant improvements were seen in the MMSE, trail making test part 246 B, or Stroop colour and word test [76].

247

248 Multiple other studies have demonstrated the positive effect of endurance exercise training on 249 cognition in T2DM. One study found that 6 months of endurance exercise significantly improves 250 executive function in older adults with glucose intolerance, including prediabetic and newly diagnosed 251 diabetic patients. Memory, however, showed no significant improvements [77]. Another study 252 investigated the relationship between 3 months of endurance exercise and executive function in 10 253 T2DM patients and 10 age-matched controls. Executive function was tested at three different time 254 points: a pre-test to familiarise the participants with the tests, and to avoid practice effects in the post-255 exercise test, a pre-exercise test one month after the pre-test, and a post-exercise test. Differences 256 between the pre-exercise and post-exercise test were assessed. A significantly improved performance 257 was observed in the diabetic group for the Colour Trails 2 test, in which the participants needed to 258 connect numbered dots in numerical order while alternating the colour of the dots. The time to

perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license .

259 complete the task decreased from 117 seconds to 92 seconds in the diabetic group, as opposed to a 260 non-significant decrease from 103 seconds to 99 seconds in the healthy control group. No significant 261 improvements were found for the Digit Span Backwards or Stroop C tests [78]. Additionally, in a study by Wang et al. (2023), the effect of endurance training on hippocampal volume and cognition was 262 263 tested in 82 T2DM patients. The patients were divided into a control group and a training group. After 264 one year of endurance training, they found significantly increased MMSE and MoCA scores in the 265 training group compared to the control group, suggesting that endurance exercise can improve 266 cognitive function in T2DM [79].

- 267
- 268 Animal studies

269

270 Also animal studies show a predominantly positive effect of endurance exercise training on cognition. 271 Shekarchian et al. (2023) found that streptozotocin- and high fat diet-induced diabetic C57BL/6 J mice 272 that received 4 weeks of swimming training scored significantly better on tests for working, spatial, 273 and recognition memory compared to non-exercising diabetic mice [80]. Another study showed that 274 swimming training for 12 weeks in streptozotocin-nicotinamide-induced type 2 diabetic rats improved 275 exploratory behaviour, locomotor activity, passive avoidance memory, and non-spatial cognitive 276 memory compared to their sedentary counterparts. However, this improvement was found to be 277 insignificant [81].

278

279 The Morris water maze test has oftentimes been used to demonstrate the effect of endurance exercise 280 on cognition in animal models of T2DM. In this test, the animal is placed in a circular swimming pool 281 where a platform is hidden somewhere under the water surface. After a few practice rounds, the time 282 it takes the animal to find the platform, and the length of the path that is followed, are measured. 283 These measures give an indication of spatial learning and memory [82]. For example, one study looked 284 into the effect of 4 months of both light and moderate intensity treadmill running on memory in pre-285 symptomatic Otsuka-Long-Evans-Tokushima fatty (OLETF) rats. Before the exercise intervention, the 286 OLETF rats spent considerably less time in the quadrant area where the platform was located during 287 the learning phase, compared to control Long-Evans Tokushima (LETO) rats. After the intervention, 288 both exercise intensities showed to have improved memory function in the OLETF rats, since both 289 escape latency and swim length were shortened [83]. Lang et al. (2020) also used the Morris water 290 maze test to test the effect of 8 weeks of moderate-intensity treadmill exercise on memory in T2DM 291 mice. The exercising T2DM mice showed a significantly reduced escape latency on day 4-5 compared 292 to the non-exercising T2DM mice, and their number of platform crossings significantly increased over 293 time [84]. Another similar study was performed where the effect of 4 weeks of moderate-intensity 294 treadmill running in OLETF rats was investigated. They found that both swim path length and escape latency of the exercising OLETF rats were reduced at trials on days 2-4, and that their time spent in the 295 296 platform area significantly improved after the exercise intervention [85]. Another study by Shima et al. 297 (2023) assessed the effect of 4 weeks of light-intensity running on a forced exercise wheel bed in ob/ob 298 mice. However, here they did not find an effect of exercise on the swim distance, escape latency, or 299 speed. They did find that the times of crossing the target platform during the probe test in the 300 exercised ob/ob mice did not differ significantly from that in the control C57BL/6 mice, and that it was greater than in the sedentary ob/ob mice, but not significantly [86]. 301

302

Based on these studies, endurance exercise conducted for 40-60 min, 3-5 times per week enables cognitive improvements in T2DM patients. Especially memory shows to be sensitive to endurance exercise-induced improvement. However, two out of five human studies discussed here only found significant improvements in MoCA and/or MMSE scores. Since the MMSE and MoCA were designed as screening tools to diagnose mild cognitive impairment (MCI) [87, 88], they cannot be considered very reliable for detecting changes in cognitive function.

- 309
- 310 <u>3.1.2 Resistance exercise training</u>

perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license .

311

#### 312 Human studies

313

314 The beneficial effect of resistance exercise on cognition in T2DM patients was shown by Zhao et al. 315 (2022). They examined the effect of 12 months of power training on executive function, 316 attention/speed, memory, and global cognition in older adults with T2DM, and aimed to determine 317 whether there is an association between cognitive improvements and improvements in muscle 318 strength, body composition, and/or endurance. 103 patients were divided into a power training group 319 or a sham low-intensity group that performed the same exercises as the power training group, but 320 without added weight. Cognition of both groups improved over time, with increased scores in the Trails 321 A, Trails B, word list recall, and word list memory. In addition, improved memory was associated with 322 both increased skeletal muscle mass and reduced body fat mass, and improvement in Trails B minus A 323 in the power training group was associated with increases in knee extension strength [89]. Yamamoto 324 et al. (2021) failed to find similar results. In their study, 60 T2DM patients aged 72.9 ± 2.4 years were 325 divided into a control group, a resistance training group, and a resistance training group with leucine 326 (an amino acid promoting muscle synthesis) supplementation. The training groups performed daily 327 bodyweight resistance exercises and exercises with elastic bands every day at home, for a total period 328 of 48 weeks. Cognitive function was assessed by means of the MMSE at baseline and after the 329 intervention period. The MMSE score in the control group was significantly decreased to 27.5 ± 2.6 330 after 48 weeks, while the score in the training groups had not significantly changed. This caused the 331 MMSE score at 48 weeks to be significantly higher in the training groups than in the control group [90].

332

333 Although both studies suggest a positive effect of resistance exercise training on cognition in T2DM, a 334 sufficient number of studies to draw a definite conclusion is lacking. Moreover, one study only assessed 335 cognitive function by means of the MMSE, limiting the information on cognitive function, and thus the 336 reliability of the results.

337

339

#### 338 3.1.3 Endurance vs. resistance exercise training

- 340 Human studies
- 341

342 To determine the exercise modality most effective in improving cognition in T2DM, some studies 343 compared the effect of endurance and resistance exercise training. Teixeira et al. (2029) investigated 344 the effect of either endurance or resistance exercise on cognitive function in T2DM patients or patients 345 with arterial hypertension. The patients were randomised into a resistance exercise group or an 346 endurance exercise group. Both groups exercised at moderate intensity for 12 weeks. Cognitive 347 function was assessed before and after the intervention by means of the mental test and training 348 system (MTTS), consisting of the cognitrone (attention and concentration), the determination test 349 (reaction time), and the visual pursuit test (selective attention). The group with both T2DM and 350 hypertension, but not the group with only hypertension, showed a significantly improved performance 351 in the cognitrone, but no significantly improved reaction time. No differences were observed between 352 the endurance and resistance exercise group. These results show that both endurance and resistance 353 exercise are capable of increasing attention and concentration in T2DM patients with arterial 354 hypertension [91].

355

356 This study shows that endurance and resistance exercise training have a comparable positive effect on 357 cognitive function in T2DM, however, additional evidence confirming these results is needed.

358

359 3.1.4 Combined endurance and resistance exercise training

- 360 361 Human studies
- 362

perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license .

363 Several studies have explored the effect of combined exercise training, consisting of both resistance 364 and endurance exercise, sometimes in combination with flexibility or balance exercises, on cognition 365 in T2DM patients. One study found significantly improved inhibitory control, working memory, 366 cognitive flexibility, and attention after 8 weeks of combined exercise training in T2DM patients, 367 compared to a control group [92]. The same research team also examined the effect of 8 weeks of 368 combined exercise on plasma BDNF levels, executive function and long-term memory in T2DM 369 patients. They observed significantly improved executive function following the combined exercise, 370 while BDNF levels were not found to be significantly changed [93]. Espeland et al. (2027) conducted 371 an exploratory analysis of data from the Lifestyle Interventions and Independence for Elders (LIFE) trial, 372 which was a randomised controlled clinical trial of exercise intervention consisting of walking, 373 resistance training, and flexibility exercises in sedentary non-demented T2DM patients and healthy 374 subjects. Cognitive function was tested at baseline and 2 years after randomisation. They found that 375 cognitive function, more specifically global cognitive function and delayed memory, significantly 376 improved in the diabetic participants of the intervention group only, suggesting a beneficial effect of 377 combined training on cognition in T2DM [94]. Additionally, one study determined the effect of 12 378 weeks of combined exercise training, consisting of endurance, resistance, and balance exercises, on 379 cognition in women with T2DM. Cognition was tested by means of the MoCA, the digit symbol 380 substitution test, and the forward digit span test. After the 12-week intervention, the exercise group 381 scored significantly higher on the MoCA compared to the control group, and, with an increase of 3.1, 382 improved significantly compared to baseline. However, neither of the groups showed improvement in 383 the other tests [95].

384

385 In a study by Martinez-Velilla et al. (2021), 103 acutely hospitalised elderly T2DM patients were 386 randomised to an exercise group or a control group. Cognitive function was assessed by means of the 387 MMSE at baseline and at discharge. The exercise intervention consisted of a combination of resistance, 388 balance, and walking exercises. The median length of stay, and thus of the intervention, was 8 days. 389 There was no difference in MMSE score between both groups at baseline. However, after the 390 intervention, the exercise group scored on average 1.6 points higher than the control group (23.7 vs. 391 22.1), which was found to be significant [96]. In addition, a pilot study looked into the effect of 6 392 months of a progressive endurance- and resistance-training program on the brain and cognition in 393 T2DM. 50 T2DM patients were randomised into an intervention group and a control group. The 394 intervention group performed 6 months of endurance and progressive resistance training. The control 395 group received upper and lower limb stretching of light intensity and a gentle movement program, 396 which were performed in the same volume, frequency and setting as in the intervention group. The 397 intervention group showed improved hippocampal and total brain volumes, improved white matter 398 integrity, and less decline in white matter volume. They also showed a better global cognitive score, 399 and better performance on the Digit Symbol Coding Test, Rey Complex Copy test, Stroop C-D, Trail 400 Making Test A and B, Hopkins verbal learning test (intermediate and recognition scores), and 401 Controlled Oral Word Association Test compared to the control group [97]. Another pilot study 402 examined the effect of 6 weeks of high-intensity low-volume (HIFT) vs. low-intensity high-volume (LIFT) 403 functional training on cognition in cognitively impaired elderly T2DM patients. The HIFT group 404 exercised three times a week at 100-120% of the lactate threshold, while the LIFT group exercised five 405 times a week at 70-75% of the lactate threshold. The MMSE was used to diagnose cognitive 406 impairment (MMSE < 23). Processing speed, learning, memory, and attention were assessed by means 407 of the Symbol Digit Modalities Test (SDMT), California Verbal Learning Test Second Edition (CVLT-II), 408 Brief Visuospatial Memory Test-Revised (BVMT-R), and Stroop tests respectively. After the 409 intervention, MMSE, Stroop, SDMT, CVLT-II and BVMT-R scores had improved significantly in the HIFT 410 group, while only MMSE and Stroop scores, had improved significantly in the LIFT group. However, the 411 only cognitive score change that was significantly different from the control group that did not receive 412 an exercise intervention, was the change in Stroop scores in the HIFT group [98]. 413

perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license .

- 414 Overall, most studies suggest a positive effect of combined exercise training on cognitive function in
   415 T2DM. However, mixed findings indicate the need for additional studies.
- 416

417 <u>3.1.5 Other types of exercise</u>

418

419 *Human studies*420

- 421 Besides the well-known exercise forms such as running and cycling, some studies have explored the 422 effect of unconventional exercise training on cognition in T2DM.
- For example, one study compared the effect of 36 weeks of Tai Chi Chuan, a mind-body exercise, and 36 weeks of fitness walking on global cognitive function in T2DM patients. Both interventions significantly increased MoCA scores, with a significantly larger effect of Tai Chi Chuan compared to fitness walking [99].
- The effect of low-intensity Qigong exercise on cognitive function in older T2DM patients was also investigated [100]. Qigong is an ancient Chinese exercise for mind-body integration and is often used in the prevention and treatment of chronic metabolic diseases [101]. The participants in the exercise group practiced Kinect-based Kaimai-style Qigong for 12 weeks. Based on MMSE scores at baseline and after the intervention, the Qigong group showed significant improvements in cognitive function
- 432 compared to the control group [100].
- 433 Coelho Junior et al. (2018) investigated whether multicomponent exercise (MCE) improves executive 434 function in T2DM. 72 T2DM patients performed MCE for 26 weeks. MCE consisted of 12 different 435 exercises mimicking activities of daily live gestures, performed for 1 minute each, with each exercise 436 immediately followed by a 2 minute walk. Although the patients showed an increased mobility after 437 the intervention, they did not show improvements in executive function [102].
- 438
- Finally, the effect of a 24-week square-stepping exercise (SSE) program on cognition in T2DM patients older than 49 years was also investigated. The researchers assessed global cognitive function, delineated into 4 domains: memory, reasoning, concentration and planning. The SSE group showed greater improvement in planning scores compared to the control group. However, the improvement in this specific domain might be related to the nature of the training program, where the participants are required to memorise patterns [103].
- These findings suggest that unconventional exercise, often involving the mind to a larger extent than conventional exercise, can contribute to improved cognition in T2DM.

447 Table 3 Characteristics of included studies

	Author	Year	Age	Population	N intervention	Control group	N controls
Endurance exercise training	Leischik et al.	2021	59.1 (control); 60.4 (walking); 56.4 (pedometer)	T2DM patients walking or T2DM patients counting 10 000 steps/day with pedometer	36 (18 per group)	T2DM patients walking 30 min/day	16
ti unning	Parsa et al.	2021	6-8 weeks	Male diabetic albino Wistar rats with swimming training, P. Psyllium, or both	30 (10 per group)	Healthy sedentary control group, sedentary diabetic group	20 (10 per group)
	Anderson- Hanley et al.	2012	72.8/73.7	T2DM patients	10	Age-matched non-diabetic controls	10
	Ploydang et al.	2023	60-75	T2DM patients	17	Non-exercising control T2DM	17
	Wang et al.	2023	60-75	T2DM patients receiving endurance training	40	T2DM patients without training	42
	Shekarchian et al.	2023	10-11 weeks	Exercising adult male C57BL/6 J T2DM mice	10	Non-exercising adult male C57BL/6 J T2DM and control mice + exercising control mice	30 (10 per group)
	Jesmin et al.	2022	8 weeks	Otsuka-Long-Evans-Tokushima fatty (OLETF) rats	26	Long-Evans Tokushima (LETO) rats (sedentary, light intensity exercise, and moderate intensity exercise) + sedentary OLETF rats	38 (13, 7, 5 and 13 respectively)
	Lang et al.	2020	10 weeks	Exercising C57BL/ 6 T2DM mice (E)	9	Control C57BL/6 mice (C), non- exercising T2DM mice (T), and exercising T2DM mice with nicotinamide treatment (NAM)	27 (9 per group)
	Shima et al.	2017	26 weeks	Exercising Otsuka-Long-Evans-Tokushima fatty (OLETF) rats	6 to 8	Exercising and non-exercising Long- Evans Tokushima (LETO) rats + non- exercising OLETF rats	6 to 8 per group
	Shima et al.	2023	8 weeks	Exercising ob/ob mice	5	Sedentary ob/ob mice and sedentary C57BL/6 mice	10 (5 per group)
Resistance exercise training	Zhao et al.	2022	67.9 ± 5.5	T2DM patients with power training	49	T2DM patients with sham low-intensity training (same machines: 3 sets of 8 reps with empty bar)	54
-	Yamamoto et al.	2021	72.9 ± 2.4	T2DM patients performing resistance exercise or resistance exercise + leucine supplementation	18 (R) 17 (RL)	Control group of T2DM patients	17
Endurance vs. resistance exercise training	Teixeira et al.	2019	55 ± 12	Patients with T2DM and/or systemic arterial hypertension performing resistance training	6	Patients with T2DM and/or systemic arterial hypertension performing endurance training	7
Combined endurance and resistance	Silveira- Rodrigues et al.	2021	63.9 ± 7.7 (training) 63.3 ± 7.8 (control)	T2DM patients	16	T2DM control group	15

exercise training							
	Espeland et al.	2017	70-89	Sedentary non-demented T2DM patients and non-diabetic adults receiving physical activity intervention	735	Sedentary non-demented T2DM patients and non-diabetic adults receiving health education	741
	Silveira- Rodrigues et al.	2023	63±8	T2DM patients performing combined training	17	Control T2DM patients not exercising	18
	Ghodrati et al.	2023	55-70	Women with T2DM receiving combined training	13	T2DM women not receiving intervention	9
	Martinez-Velilla et al.	2021	86 ± 5 (control) 87 ± 4 (intervention)	Acutely hospitalised elderly T2DM patients in exercise group	54	control acutely hospitalised elderly T2DM patients	49
	Callisaya et al.	2017	66.2	T2DM patients receiving intervention	24	T2DM patients who received upper and lower limb stretching of light intensity and a gentle movement program, which were performed in the same volume, frequency and setting as the intervention group	23
	Ghahfarrokhi et al.	2024	67.5±5.8	Elderly, cognitively impaired T2DM patients performing HIFT or LIFT	32 (16 per group)	Elderly, cognitively impaired T2DM patients without intervention	16
Other exercise types	Shellington et al.	2018	71.2 (6.9) (control) 65.9 (5.2) (SSE)	T2DM patients with self-reported cognitive complaints performing SSE	12	T2DM patients with self-reported cognitive complaints without intervention	13
	Chen et al.	2023	67.55 [5.02]	T2DM patients performing Tai Chi Chuan	107	T2DM patients performing fitness walking + control group receiving only self-management education	110 (fitness walking) 111 (control)
	Cai et al.	2019	older adults	T2DM patients performing Qigong exercise	27	Control T2DM group without intervention	28
	Coelho Junior et al.	2018	64.4 ± 5.7 (control) 66.0 ± 3.2 (T2DM)	T2DM patients performing MCE	72	Age-matched healthy controls performing MCE	72

# 

	Author	Type of exercise training	Duration of exercise training	Duration of each session	Frequency of exercise training	Intensity of exercise training	Outcomes (cognitive tests)	Main results
Endurance exercise training	Leischik et al.	Walking	12 weeks	40 min	3x/week	Ş	FAIR-Test 2, verbal memory, non-verbal memory	Walking and counting steps led to enhanced cognitive performance pace and concentration
	Parsa et al.	Swimming training	12 weeks	40 min	5 days/week	?	Cognitive memory: novel object recognition test and elevated plus maze Aversive memory and learning: passive avoidance learning test	Swimming training alone insignificantly improved exploratory behaviour, locomotor activity, and passive avoidance memory, and non- spatial cognitive memory compared to sedentary diabetic rats
	Anderson-Hanley et al.	Endurance exercise on stationary bike or cybercycle	3 months	45 min	5x/week	Moderate to vigorous	Executive function: Colour trails 2 test, Stroop C test, Digit span backwards test	Significant improvements in colour trails 2 test in exercise group
	Ploydang et al.	Aquatic Nordic walking	12 weeks	60 min	3x/week	First 6 weeks: 40- 50% HRR Last 6 weeks: 50- 60% HRR	MoCA, MMSE, trail making test part B, Stroop colour and word test	Significantly increased MoCA scores in Nordic walking group. No significant improvements in MMSE, Stroop test, or trail making test.
	Wang et al.	Aerobic dancing	1 year	60 min	3x/week	?	MMSE + MoCA	Aerobic training group had significantly increased MMSE and MoCA scores compared to the control group
	Shekarchian et al.	Swimming	4 weeks	Week 1-2: 3x10min week 3-4: 6x10min	5 days/week	?	Working memory (Y maze), recognition memory (Novel object recognition), spatial memory, (Morris water maze)	Exercise improved all memory aspects in exercising T2DM mice compared to non-exercising T2DM mice
	Jesmin et al.	Treadmill running	4 months	30min/day	5 days/week	Light intensity: OLETF 7.0 m/min, LETO 10.0 m/min; moderate intensity: OLETF 12.5 m/min, LETO 20 m/min	Morris water maze	Shortened swim length and escape latency in both light and moderate exercise groups → improved memory
	Lang et al.	Treadmill running	8 weeks	37min/day	5 days/week	Moderate intensity	Morris water maze	Compared with T group, the escape latency of E group was significantly reduced on day 4– 5, treadmill exercise significantly increased the number of platform crossings in E group

	Shima et al.	Treadmill running	4 weeks	30min/day	5 days/week	Moderate- intensity (OLETF rats, 12.5 m/min; LETO rats, 20 m/min)	Morris water maze	Escape latency and swim path length of exercising OLETF rats were shortened, without alteration of the speed of swimming, at trials on days 2, 3 and 4. The time spent by OLETF rats in the platform area significantly improved after 4 weeks of exercise
	Shima et al.	Running on a forced exercise wheel bed	4 weeks	30min/day	5 days/week	Light intensity	Morris water maze	No effect of exercise on the escape latency, swim distance or speed in ob/ob mice. The times of crossing the target platform during the probe test in C57BL/6 mice were significantly greater than that in sedentary ob/ob mice, but not than that in exercised ob/ob mice
Resistance exercise training	Zhao et al.	Whole-body, machine based power training (3 sets of 8 reps of lateral pulldown, chest press, upper back, leg press, knee extension, and knee flexion, hip extension and hip abduction)	12 months	Ş	3 days/week	High intensity (15-18 on Borg scale, 80% 1RM)	Memory: word list memory, word list recall, and word list recognition subtests of the Consortium to Establish a Registry for Alzheimer's Disease Attention/speed: Trail Making Test Part A (Trails A) Executive function: Trail Making Test Part B (Trails B) and the Trail Making Test B minus A (Trails B minus A) Global cognitive function: MMSE	Cognition of both groups improved over time, with increased scores in the Trails A, Trails B, word list recall, and word list memory
	Yamamoto et al.	Body weight resistance training + exercises with elastic bands: tube fly, front raise, hammer curl, leg extension, calf raise, and squat. Performed 20 times each	48 weeks	?	Daily	?	MMSE	No significant changes in MMSE score
Endurance vs resistance exercise training	Teixeira et al.	Endurance exercise: elliptical, bike, treadmill, or upper-body cycle ergometer Resistance exercise	12 weeks	40 min	3x/week	Endurance: 60% HRmax resistance: 11-12 Borg scale	Mental test and training system (MTTS): cognitrone (attention and concentration), determination test (reaction time), visual pursuit test (selective attention)	The group with both T2DM and hypertension showed a significantly increased number of reactions, but no significantly improved reaction time. No differences were observed between the endurance and resistance exercise group.
Combined endurance and resistance	Silveira-Rodrigues et al.	6 resistance exercises (reverse grip lat pulldown, leg press, bench press machine, calf raises	8 weeks	?	3x/week	?	MoCA	Significantly improved inhibitory control, working memory, cognitive flexibility, and

exercise training		machine, dumbbell shoulder press, and abdominal crunches) + 15 min of treadmill walking, with increasing weight and repetition number over 8 weeks						attention in the exercise group compared to the control group
	Espeland et al.	Walking, resistance training, and flexibility exercises	2 years	50 min	3-4x/week	Moderate intensity	Modified MiniMental State Exam (3MSE) (global cognitive function); the Wechsler Adult Intelligence Scale-III Digit Symbol Coding test (DSC) (psychomotor speed, attention, and working memory); Hopkins Verbal Learning Test-Revised (HVLT-D) (delayed recall) Computer tests (processing speed and executive function): 1-back and 2-back tasks, the Eriksen Flanker task, and a task-switching paradigm	Global cognitive function and delayed memory, significantly improved only in T2DM patients who received training
	Silveira-Rodrigues et al.	Combined training: endurance exercise + resistance exercise	8 weeks	40-60min	3x/week	?	Executive function: Trail Making Test, Stroop Colour Task, and Digit Span Long-term memory: simplified Taylor Complex Figure Test	Executive function was significantly improved in exercise group
	Ghodrati et al.	Combined training: endurance + resistance + balance exercises	12 weeks	65 min	3x/week	Endurance: 55-75% Hrmax resistance: 65-85% 1RM	MoCA, digit symbol substitution test, forward digit span test	Intervention group scored significantly higher on MoCA than control group, and increased score with 3.1 compared to baseline. No improvement in digit symbol substitution test or forward digit span test.
	Martinez-Velilla et al.	Combination of resistance, balance, and walking exercises. Squats rising from a chair, leg press, and bilateral knee extension, seated bench press. semitandem foot standing, line walking, stepping practice, walking with small obstacles, proprioceptive exercises on unstable surfaces, altering the base of support, and weight transfer from one leg to the other, knee extension and flexion, hip abduction, walking along corridor	8 days	20 min	2x/day 5-7 days/week	?	MMSE	After the intervention, the exercise group scored on average 1.6 points higher than the control group (23.7 vs. 22.1), significant
	Callisaya et al.	Combination of endurance and resistance exercise	6 months	1 hour	3x/week	Started at low to moderate, progressed	MRI to measure cortical thickness, total brain volume, hippocampal volume and white matter volume, and white	The intervention group showed improved hippocampal and total brain volumes, improved white matter integrity, and less

						moderate to vigorous	Cognitive function: Victoria Stroop test, the Trail Making Test (shifting score B-A), the Digit Symbol Coding Test, the digit span subtest of the Wechsler Adult Intelligence Scale (third edition), Controlled Oral Word Association Test, the three part Hopkins Verbal Learning Test (revised), and the Rey Complex Figure copy and delay	They also showed a better global cognitive score, and better performance on the Digit Symbol Coding Test, Rey Complex Copy test, Stroop C-D, Trail Making Test A and B, Hopkins intermediate and recognition, and Controlled Oral Word Association Test
	Ghahfarrokhi et al.	Endurance exercises, upper and lower body strength, balance exercises and maintaining posture, and hip control exercises and mid-body stability	6 weeks	30-35 min (HIFT) 40-45 min (LIFT)	3x/week (HIFT) 5x/week (LIFT)	HIFT: 100- 120% lactate threshold LIFT: 70- 75% lactate threshold	MMSE, Symbol Digit Modalities Test (SDMT), CVLT-II, Brief Visuospatial Memory Test- Revised (BVMT-R), Stroop tests	MMSE, Stroop, SDMT, CVLT-II and BVMT-R scores improved significantly in the HIFT group. MMSE and Stroop scores improved significantly in the LIFT group. Only change in Stroop scores in the HIFT group was significantly different from the control group.
Other exercise types	Shellington et al.	Square-stepping exercise (SSE) program	24 weeks	60 min (45 min SSE)	2x/week	Low intensity	12-item computer-based Cambridge Brain Sciences (CBS) cognitive battery to test memory, concentration, planning, reasoning + combined global cognitive function score	The SSE group showed greater improvement in planning scores compared to the control group, no other improvements
	Chen et al.	24-form simplified tai chi chuan	36 weeks	60 min	3x/week	Moderate intensity	MoCA	Both interventions significantly increased MoCA scores, with a significantly larger effect of Tai Chi Chuan compared to fitness walking
	Cai et al.	Kinect-based Kaimai-style Qigong	12 weeks	30 min	3x/week	Low intensity	MMSE	Qigong group showed significant improvements in cognitive function compared to the control group
	Coelho Junior et al.	Multicomponent exercise: 12 different exercises mimicking activities of daily life gestures	26 weeks	36 min	2x/week	Moderate to hard	Timed up-and-go cognitive test	No improvements in executive function in T2DM

to

matter microstructure

decline in white matter volume.

perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license .

# 453 **3.2** Reviewing targets and mediators of the effect of exercise training on the brain in T2DM

454 To understand why certain exercise trainings cause cognitive improvement in T2DM, and others do 455 not, there is a need for studies investigating the underlying biomedical mechanisms of the effect of 456 exercise on the brain and cognition in T2DM. This knowledge can contribute to the development of 457 pharmacological therapies and exercise programs specifically designed for improving brain health and 458 cognition in T2DM in the most efficient way. The following section discusses such studies, and aims to 459 identify the different targets and mediators involved. Although there is much more extensive literature 460 available on the possible mediators of the effect of exercise on the brain in T2DM, this review will 461 solely focus on the studies in T2DM that have actually demonstrated an involvement of certain 462 mediators.

463

## 464 465

# 54 <u>3.2.1 AD-related pathological markers: amyloid β peptides and hyperphosphorylated tau</u>

466 Increasing evidence and literature points to converging pathways and pathogenetic processes in T2DM 467 and AD. Here we focus on studies showing effects on amyloid  $\beta$  and tau in T2DM due to exercise 468 training. We refer to extensive reviews regarding T2DM and AD for further reading [24, 104, 105].

- 469
- 470 Leptin

471 Several studies have indicated that exercise is negatively associated with cerebral amyloid  $\beta$  and 472 hyperphosphorylated tau in T2DM, and contributes to memory maintenance [106, 107]. The aspartyl 473 protease  $\beta$ -site A $\beta$ PP-cleaving enzyme 1 (BACE1) is responsible for catalysing the rate-limiting step of 474 amyloid  $\beta$  production. It has been shown that the adipocytokine leptin reduces BACE1 activity and 475 expression, thereby limiting the production of amyloid  $\beta$  [108, 109]. This is supported by Rezaei et al. 476 (2023), who suggested leptin as a possible mediator of exercise-induced prevention of memory 477 impairment in T2DM after finding elevated serum and hippocampal levels of this hormone together 478 with decreased hippocampal levels of BACE1, amyloid  $\beta$  and hyperphosphorylated tau after 8 weeks 479 of HIIT training in T2DM rats [110].

480

481 GSK3B

482 The maintenance of cognitive function in exercising T2DM rats treated with dexamethasone, 483 demonstrated by De Sousa et al. (2020), was according to the authors presumably mediated by the 484 observed lesser inhibition of the activation of hippocampal IRS-1 and higher concentration of GSK3β 485 phosphorylated on serine-9 (Ser-9) [111]. GSK3 $\beta$  is namely inactivated by Ser-9 phosphorylation [112], 486 which prevents GSK3ß activation, and subsequently tau phosphorylation and associated neurotoxic 487 effects [113, 114]. Accordingly, Rezaei et al. (2023) also found decreased hippocampal GSK3β 488 dephosphorylation together with decreased hippocampal levels of amyloid  $\beta$  and phosphorylated tau 489 in HIIT-trained T2DM rats [110]. GSK3β plays an important role in neurodegeneration in T2DM. Both 490 glucolipotoxicity and insulin resistance contribute to GSK3β overactivation, leading to β-catenin 491 phosphorylation and subsequent proteasomal degradation. This results in the inhibition of the 492 expression of reactive oxygen species (ROS) scavenging enzymes, which leads to more oxidative stress 493 and consequently the disruption of mitochondrial structure, function and axonal trafficking. Moreover, 494 GSK3β overactivation also stimulates tau hyperphosphorylation, leading to microtubule destabilisation 495 and thus disruption of axonal mitochondrial trafficking, as well as to a decrease in mitochondrial 496 complex I and thus impaired energy production [114]. By inhibiting the activation of GSK3 $\beta$  through 497 Ser-9 phosphorylation, exercise could thus contribute to the prevention of cognitive decline in T2DM.

498

# 499 Adiponectin

500 In the study of Rezaei et al. (2023), the T2DM rats showed decreased serum and hippocampal levels of 501 insulin and adiponectin at baseline, as well as decreased levels of insulin receptors, adiponectin 502 receptors and AMPK in the hippocampus, and increased hippocampal GSK3β and 503 hyperphosphorylated tau. HIIT training counteracted these findings. Both serum insulin and

perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license .

504 adiponectin were significantly increased by HIIT, as well as the levels of hippocampal insulin and 505 adiponectin receptors. HIIT training also increased hippocampal AMPK phosphorylation, and 506 decreased hippocampal tau phosphorylation and GSK3β dephosphorylation. The authors suggest that 507 the increased levels of adiponectin can contribute to the preservation of hippocampal volume and 508 function through the stimulation of synaptic plasticity, as it has been shown by Pousti et al. (2018) and 509 Weisz et al. (2017) respectively that adiponectin modulates synaptic plasticity in the hippocampal 510 dentate gyrus and regulates hippocampal synaptic transmission [110, 115, 116]. Another study shows 511 memory and learning impairments together with reduced AMPK phosphorylation and increased GSK3β 512 activation in adiponectin-knockout mice [117], suggesting that the neuroprotective effects of 513 adiponectin are mediated via AMPK phosphorylation and GSK3<sup>β</sup> inactivation, which is in line with the 514 study of Rezaei et al. (2023). The exercise-induced decreased GSK3β Ser-9 dephosphorylation means 515 that there is a higher ratio of inactive, phosphorylated GSK3 $\beta$ , preventing tau phosphorylation. In 516 addition, the adiponectin-mediated AMPK phosphorylation could also be neuroprotective by inhibiting 517 the inflammatory response of microglia to amyloid  $\beta$  [118].

519 <u>3.2.2 Brain volume</u>

520

518

521 Several studies have demonstrated that exercise increases hippocampal volume in T2DM patients. One 522 study showed that a higher step count, measured over 7 days, was associated with a larger 523 hippocampal volume in T2DM patients [59]. In addition, one year of endurance training in T2DM 524 patients with normal cognition significantly increased hippocampal volume, and prevented a decline 525 in MMSE and MoCA scores. Hippocampal volume in the training group was significantly increased 526 compared to baseline, as well as compared to the control group [79]. Another study showed a higher 527 hippocampal CA1 and CA3 neuronal density in diabetic Sprague-Dawley (SD) rats after 6 weeks of 528 exercise on a running wheel, probably contributing to the observed improved cognitive function [119]. 529 Similarly, a pilot study in which T2DM patients performed 6 months of a progressive endurance- and 530 resistance-training program showed improved hippocampal and total brain volumes, improved white 531 matter integrity, and less decline in white matter volume in the intervention group compared to the 532 control group [97].

- 533
- 534 <u>3.2.3 Neuronal survival</u>

535 536 BDNF

537 BDNF is one of the most important neurotrophic factors. By binding the TrkB receptor, it regulates and 538 promotes cell survival, neuroplasticity, and neurogenesis in the central nervous system [120, 121]. As 539 already mentioned previously, BDNF plays an important role in the effect of exercise on cognitive 540 function. A study investigating the effect of 4 weeks of swimming training on working, spatial, and 541 recognition memory in diabetic C57BL/6 J mice, shows that higher PA is associated with increased 542 hippocampal and prefrontal BDNF levels, together with an improvement in all memory aspects [80]. 543 Similarly, Jesmin et al. (2022) showed significantly increased hippocampal BDNF levels together with 544 improved memory function in OLETF rats after 4 months of endurance exercise [83]. This shows that 545 also under diabetic conditions, BDNF is an important mediator of exercise effects on cognition.

546

Another study looked into the effect of 12 weeks of swimming training on BDNF/TrkB signalling and apoptosis in the cerebral cortex of male diabetic C57BL/6JNarl mice. The exercised diabetic mice showed less neural apoptosis compared to the control diabetic mice, accompanied by an increased activity of the BDNF/TrkB signaling pathway, and a decreased activity of the Fas/FasL-mediated and mitochondria-initiated apoptotic pathways. These findings indicate that exercise promotes neuronal survival in diabetic mice through the BDNF pathway [122].

554 <u>3.2.4 Neuroplasticity and neurogenesis</u>

555

553

perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license .

556 One study determined the effect of 7 weeks of treadmill exercise on neuroblast differentiation in the 557 subgranular zone of the dentate gyrus (SZDG) in Zucker diabetic fatty (ZDF) rats and Zucker lean control 558 (ZLC) rats. At 23 weeks old, the rats started to exercise for 7 weeks at 12-16m/min. Compared to the 559 non-exercising groups, the neuroblasts of both the exercising diabetic and exercising control rats 560 showed a significant increase in tertiary dendrites. However, the number of neuroblasts only increased 561 in the control rats. This shows that endurance exercise in diabetic rats can stimulate neuroplasticity, 562 but not necessarily neuroproliferation [123]. A similar study tested the effect of 5 weeks of treadmill 563 exercise on cell differentiation and proliferation in the SZDG of the same ZDF rat model. Proliferation 564 was detected by means of the proliferation marker Ki67, and progenitor differentiation into neurons 565 was detected by means of the differentiation marker doublecortin (DCX). In this study, the rats started 566 exercising at 6 weeks old, and ran at 22m/min. Compared to the sedentary ZDF rats, the exercised rats 567 showed a significant increase in Ki67 positive cells and DCX-immunoreactive structures, indicating both 568 increased proliferation and increased differentiation [124]. This confirms that exercise can stimulate 569 neuronal differentiation in the diabetic SDZG, as shown by Hwang et al. (2010) [123]. The fact that, this 570 time, exercise was able to stimulate neuronal proliferation as well, might indicate that exercise has to 571 be performed at a certain intensity rather than a certain volume to achieve this effect.

572 573 BDNF

574 Increased hippocampal BDNF levels in combination with increased dendritic spine density on the 575 secondary and tertiary dendrites of dentate granule neurons were found after voluntary wheel running 576 in *db/db* mice [125]. This suggests that BDNF contributes to exercise-induced neuroplasticity in 577 diabetic mouse models.

578

# 579 PI3K/Akt and AMPK/SIRT1

580 It was demonstrated that endurance exercise can upregulate synaptic plasticity-associated proteins in 581 the hippocampus of T2DM rats, presumably via the activation of PI3K/Akt/mTOR and AMPK/SIRT1 signalling pathways and inhibition of the NF $\kappa$ B/NLRP3/IL-1 $\beta$  signalling pathway [126]. Under non-582 583 diabetic conditions, the PI3K/Akt/mTOR pathway is activated by insulin, resulting in the formation of 584 dendritic spines and excitatory synapses in hippocampal neurons [127]. In T2DM, insulin resistance 585 prevents this activation, contributing to reduced synaptic plasticity. Exercise thus could, to a certain 586 extent, take over this role of insulin. The aforementioned BDNF also activates the PI3K/Akt pathway, 587 constituting another mediator of the exercise-induced increase in synaptic plasticity-associated 588 proteins [120]. Activation of the AMPK/SIRT1 pathway, on the other hand, contributes to synaptic 589 plasticity by increasing insulin sensitivity and glucose uptake, regulating BDNF expression, promoting 590 neuronal survival, etc. [128-132]. During exercise, AMPK activates SIRT1, after which SIRT1 591 deacetylates PGC-1 $\alpha$ , resulting in its activation. PGC-1 $\alpha$  is important for mitochondrial remodelling and 592 biogenesis, contributing to synaptic plasticity [133]. Its activation leads to the upregulation of FNDC5, 593 which in turn crosses the BBB and upregulates hippocampal BDNF expression [120].

594

595 Another study also found an increase in the synaptic plasticity proteins synaptophysin (SYN) and N-596 methyl-D-aspartate receptor (NMDAR) in the prefrontal cortex of diabetic SD rats after 4 weeks of 597 treadmill training. They demonstrated that endurance exercise increased the level of phosphorylated 598 PI3K, suggesting an increased activity of the PI3K/Akt pathway, and increased insulin sensitivity. In 599 addition, they found a slight increase in the phosphorylation and a slight decrease in the acetylation 600 of FOXO1, which is a transcription factor that promotes NF-kB activity and consequently the expression 601 of proinflammatory factors. FOXO1 phosphorylation enables its cytoplasmic retention and thus 602 prevents the transcription of its target inflammatory genes [134]. Since FOXO is a target of Akt, the 603 increased PI3K/Akt activity might be the cause of the increased FOXO1 phosphorylation [135]. Nuclear 604 NF-kB acetylation was found to be decreased as well in the rats that performed endurance exercise, 605 reducing its DNA binding and transcriptional activity [134, 136]. These findings were confirmed by a 606 study showing that 8 weeks of moderate intensity treadmill exercise in diabetic C57BL/ 6 mice

perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license .

607 improves cognitive dysfunction and increases the release of BDNF through the activation of the 608 SIRT1/PGC-1 $\alpha$ /FNDC5/BDNF pathway [84].

609

610 <u>3.2.5 Cerebral blood flow</u>

611

612 Zhao et al. (2023) investigated the effect of a 2-month moderate-intensity exercise program on brain 613 oxygenation in sedentary older T2DM patients with cognitive impairment. They also looked into the 614 effect of the exercise program on the hemodynamic responses to the Mini-Cog test, which assesses 615 short-term memory and visuo-constructive abilities. The exercise program consisted of a combination 616 of endurance and resistance exercise. They found that the exercise program improved the very low-617 frequency oscillations (<0.05 Hz) during the Mini-Cog test, as assessed by near-infrared spectroscopy. 618 Since very-low frequency oscillations are crucial for small-vessel function, this indicates that combined 619 exercise has the power to improve cerebral vascular function in T2DM. In addition, the exercise 620 program significantly reduced the oxyhemoglobin (oxyHb) activation in the right superior frontal 621 region during the Mini-Cog test, thereby reducing the overactivation usually observed in T2DM [137]. 622

- 623 <u>3.2.6 Glycometabolism</u>
- 624

# 625 Lactate

626 Since dysregulated peripheral glycometabolism is a hallmark of T2DM, Shima et al. (2017) 627 hypothesised that hippocampal glycometabolic dysfunction might contribute to memory impairment 628 in T2DM [85]. As it has already been demonstrated in healthy animals that moderate exercise enhances 629 memory function and hippocampal glycogen levels [138, 139], they wanted to determine if this was 630 also the case in a T2DM animal model. At baseline, OLETF rats had higher hippocampal levels of 631 glycogen and lower MCT2 expression compared to LETO control rats. They also showed impaired 632 memory. After 4 weeks of treadmill running, the OLETF rats showed even higher hippocampal glycogen 633 levels, in addition to normalised hippocampal MCT2 expression together with improved memory 634 function. Since MCT2 is an important lactate transporter, this suggests that lactate plays a key role in 635 the effect of exercise on glycometabolism and memory in T2DM [85]. In line with this, it was 636 demonstrated that 4 months of treadmill running prevented the progression of cognitive decline in 637 presymptomatic OLETF rats through improved hippocampal MCT2 expression, again confirming the 638 role of lactate in the positive effect of exercise on cognition [83]. Another study by Shima et al. (2023) 639 showed the same effect in an advanced stage T2DM mouse model. Ob/ob mice were subjected to 4 640 weeks of light-intensity exercise. The exercise program resulted in improved hippocampal MCT2 levels, 641 accompanied by improved hippocampal memory retention [86]. As Soya et al. (2019) mention in their 642 review discussing these findings, the upregulated MCT2 levels enable an increased uptake of L-lactate 643 into neurons, where it can serve as an energy substrate, and act as a signalling molecule to induce the 644 expression of neuroplasticity genes, eventually contributing to the increased memory function 645 observed in these studies [140].

646

# 647 <u>3.2.7 Inflammation</u>

648

As inflammation is a very broad process, involving many different players, we would like to emphasize again that the pathways and mediators discussed in this review are only the ones that clearly have been shown to be involved in the effect of exercise on the brain in T2DM. For more information about the involvement of inflammation in T2DM, and the effect of exercise on it, we refer to more exhaustive reviews [71, 141].

654

# 655 AMPK/SIRT1

The AMPK/SIRT1 pathway is not only important for rescuing cognitive function by increasing BDNF
 levels, but also by counteracting inflammation. Treadmill exercise in diabetic C57BL/ 6 mice for 8 weeks

reduced activation of hippocampal proinflammatory microglia M1, as well as the hippocampal levels

perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license .

659 of proinflammatory factors IL-1 $\beta$ , IL-6, TNF- $\alpha$ , and increased the expression levels of anti-inflammatory 660 factors IL-10, TGF-β1. This co-occurred with an activation of the SIRT1/NF-κB pathway, which is known 661 to be responsible for counteracting inflammation, suggesting the importance of this pathway in 662 exercise-mediated anti-inflammatory actions in T2DM [84]. A different study also reported a reduction 663 in hippocampal IL-1 $\beta$  and TNF- $\alpha$ , accompanied by cognitive improvement, after T2DM SD rats 664 performed 6 weeks of endurance exercise on a running wheel. However, although we can assume that 665 the AMPK/SIRT1 pathway is involved here as well, the authors did not elaborate on the pathways 666 responsible for the observed reduction in pro-inflammatory factors [119].

667

# 668 <u>3.2.8 Insulin resistance</u>

669

670 T2DM-associated insulin resistance is not only present in the periphery, but also in the brain, where it contributes to the cognitive deficits observed in T2DM patients via multiple mechanisms [128, 142-671 672 144]. For example, insulin resistance and the consequent hyperinsulinemia have been associated with 673 increased tau hyperphosphorylation, increased deposition together with reduced breakdown of 674 amyloid  $\beta$ , dysfunctional IGF-1 receptors (IGFRs), etc. This all contributes to neurodegeneration and 675 cognitive impairment [142]. Studies have shown that the insulin-sensitising effect of exercise is also 676 manifested in the brain, which gives exercise the power to prevent or decrease these cognitive deficits 677 [145, 146]. One possible way in which this effect of exercise could be mediated, is by the afore-678 mentioned decrease in TNF- $\alpha$ . TNF- $\alpha$  activates JNK, which is a kinase that causes insulin resistance via serine phosphorylation of IRS-1 [71, 147]. IRS-1 is important for the signal transduction of insulin, 679 680 ultimately leading to glycogen synthesis and glucose transporter 4 (GLUT-4) translocation. Serine 681 phosphorylation of IRS-1 inhibits this insulin signal transduction and thus results in insulin resistance 682 [148-150]. It could thus be hypothesised that by decreasing TNF- $\alpha$ , exercise could increase insulin 683 sensitivity in the brain.

684

685 GLP-1

686 Park et al. (2019) investigated the effect of resistance exercise training on GLP-1R levels in the 687 hypothalamus of OLETF rats. The exercise training comprised 12 weeks of ladder climbing. As a result 688 of the resistance training, the rats showed increased levels of hypothalamic GLP-1R mRNA, protein 689 kinase A (PKA), protein kinase B (PKB), glucose transporter 2 (GLUT-2), and decreased hypothalamic 690 levels of protein kinase C-iota (PKC-i) [151]. All these findings point in the direction of improved 691 glycaemic control. PKB, also known as Akt, plays an important role in the PI3K/Akt pathway 692 downstream of the IR, as mentioned before in this review [23, 27, 126]. PKA, on the other hand, is 693 essential for the regulation of metabolism and triglyceride storage [152, 153], while PKC-L is known to 694 cause metabolic abnormalities in T2DM [154, 155]. GLP-1R, the main focus of this study, is a main 695 receptor involved in T2DM. It is responsible for lowering blood glucose levels by, among other things, 696 stimulating insulin secretion and suppressing glucagon secretion [156]. In addition, GLP-1R has been 697 suggested to play a neuroprotective role in diabetes-related neurodegeneration by increasing insulin 698 sensitivity [157]. Moreover, a different study has shown that recombinant human GLP-1 can reduce 699 oxidative stress by activating PKA, which was found to be elevated in the study of Park et al. (2019), 700 and by inhibiting PKC, which was found to be decreased, ultimately reversing diabetic nephropathy 701 [158]. The increase in GLP-1 mRNA found by Park et al. (2019) thus indicates decreased oxidative stress 702 and increased insulin sensitivity in the T2DM rats after resistance exercise training.

703

705

704 <u>3.2.9 Mitochondrial function</u>

Mitochondrial dysfunction is an important contributor to T2DM pathology [159, 160]. Multiple studies have already demonstrated mitochondrial dysfunction in the T2DM brain, where it contributes to neurodegeneration and cognitive dysfunction [161, 162]. It is also known that exercise positively influences mitochondrial function [163-165]. Studies show that exercise can restore mitochondrial

perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license .

function in the muscle of T2DM patients [166-169]. However, little is known about the effect of exercise on mitochondrial function in the brain in T2DM.

712

713 Insulin sensitivity

One study investigated the effect of endurance exercise training in combination with metformin on 714 715 mitochondrial function in male C57BL/6J mice with brain insulin resistance induced by a high-fat diet. 716 They observed that the mitochondria in brain regions rich in insulin receptors produced less ATP and 717 showed reduced activity of oxidative enzymes. The mice also showed elevated ROS production and 718 reduced activity of antioxidant enzymes, accompanied by higher rates of mitochondrial fission, and 719 accumulation of damaged mitochondrial proteins. Endurance exercise training together with 720 metformin improved insulin sensitivity. This resulted in reduced ROS emission, less hippocampal 721 mitochondrial fission, less mitochondrial protein oxidation, and increased ATP production in astrocytes 722 and primary cortical neurons. The reduction in ROS emission and increased ATP production were 723 counteracted by intranasal administration of the insulin receptor antagonist S961, proving that these 724 mitochondrial ameliorations were mediated by increased insulin sensitivity [170]. Since metformin was 725 also administered in this study, the observed effects are not fully attributable to exercise, however, it 726 is clear that metformin and exercise have a synergistic effect, and that exercise can certainly be an 727 added value in the treatment of T2DM.



728

729 Fig. 2 Schematic representation of mediators and targets involved in the effect of exercise training on the brain in T2DM. 730 Sharp blue arrows indicate stimulation or increase. Blunt blue arrows indicate inhibition. A $\beta$  = amyloid  $\beta$ ; AMPK = AMP-731 activated Protein Kinase; ATP = Adenosine Triphosphate; BDNF = Brain-Derived Neurotrophic Factor; FNDC5 = Fibronectin 732 type III Domain-Containing protein 5; FOXO1 = Forkhead box protein O1; GLP-1R = Glucagon-Like Peptide 1 Receptor; 733 GSK3 $\beta$  = Glycogen Synthase Kinase-3 beta; IRS-1 = Insulin Receptor Substrate 1; IL-1 $\beta$  = Interleukin 1beta; IL-6 = Interleukin 734 6; MCT2 = Monocarboxylate Transporter 2 ; mTOR = mammalian Target Of Rapamycin; Nf-κB = Nuclear Factor kappa-light-735 chain-enhancer of activated B-cells; PGC-1 $\alpha$  = Peroxisome proliferator-activated receptor Gamma Coactivator 1-alpha; PI3K 736 = Phosphoinositide 3-Kinase; PKA = Protein Kinase A; PKC = Protein Kinase C; p-tau = phosphorylated tau; ROS = Reactive 737 Oxygen Species; Sirt1 = Sirtuin 1; TNF- $\alpha$  = Tumor Necrosis Factor alpha

perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license .

#### 738 4. Discussion

739 In the systematic part of this review, 24 studies investigating the effect of exercise training on cognition 740 in T2DM were included. Although mixed results were found, most studies (20 out of 24) demonstrated 741 a significant positive effect of exercise training on cognition, which is consistent with findings of 742 previous reviews on this topic [65, 67]. 7 out of 10 studies investigating the effect of endurance 743 exercise training were able to show cognitive improvement, indicating the effectiveness of this training 744 modality. Even endurance exercise as simple as walking showed to already be successful in improving 745 cognition in T2DM. Out of the two studies investigating the effect of resistance exercise on cognition 746 in T2DM, one study failed to show cognitive improvement. This lack of effect could be attributed to 747 the lower intensity or lack of weights in this exercise intervention, although an effect was observed in 748 the sham low-intensity group of the study by Zhao et al. (2022), where also no added weights were 749 used. Additional studies are required to clarify this. Studies comparing endurance exercise training to 750 resistance exercise training found similar results for both exercise forms. Consequently, exercise 751 interventions in T2DM patients aimed at improving cognition do not have to be limited to one type of 752 exercise, and can be adapted to the patient's capacities. In addition, all included studies investigating 753 the effect of combined (endurance + resistance (+ flexibility/balance)) exercise training were able to 754 show significant cognitive improvements. Being in line with the exercise recommendations of the 755 American College of Sports Medicine for older adults [171], this finding favours combined exercise to 756 reach cognitive improvement in T2DM. Finally, tai chi chuan and Kaimai style Qigong showed to 757 improve cognition in T2DM. The beneficial effect of these exercise forms might be attributed to the 758 fact that they are motorically more complex, and require a more extensive involvement of the mind.

Based on this review, one can thus conclude that both endurance and resistance exercise training, as well as exercise training involving the mind, are effective at improving cognition in T2DM. Combined exercise training is presumably most desirable, although studies comparing combined exercise training to endurance or resistance exercise training alone are lacking. It also seems that a minimal intensity is required in resistance exercise training to elicit significant improvements. However, because no studies directly compare the effect of different exercise training intensities on cognition in T2DM, the optimal exercise intensity for cognitive improvement in T2DM is yet to be determined.

766 The mechanisms underlying cognitive improvement following exercise in T2DM have been explored in 767 the second, narrative part of this review. Both the AMPK/Sirt1 pathway and PI3K/Akt/mTOR pathway 768 have been identified as key players. The studies included in this review show that exercise-mediated 769 activation of these pathways leads to beneficial brain changes in T2DM, such as increased 770 neuroplasticity, brain volume, synaptic plasticity, dendritic spines, insulin sensitivity, mitochondrial 771 remodelling and biogenesis, and ATP production, as well as decreased inflammation, oxidative stress, 772 and amyloid  $\beta$  and p-tau production. BDNF, lactate, leptin, adiponectin, GSK3 $\beta$  and GLP-1 were 773 identified as the most important factors mediating these changes. Although other factors such as IGF-774 1 have been suggested to play a role in the positive effect of exercise on cognition [172], no studies in 775 T2DM have directly demonstrated this. Much more research is needed to identify the full spectrum of 776 involved mediators and pathways, investigate interactions between different cell types, etc. In 777 addition, it needs to be considered that the majority of the studies described in this review were 778 conducted in rodents, and are yet to be confirmed in humans.

This review has various limitations that could interfere with the reliable interpretation of its findings. First of all, the fact that the studies included in this review use a variety of cognitive tests, makes it difficult to compare the different studies. Moreover, several studies merely use the MMSE or MoCA to test cognition, which are diagnostic tools designed to screen for MCI, and not to detect (subtle) changes in cognition [87, 88]. There is a need for a standardised cognitive test battery in T2DM to

perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license .

784 increase the comparability of future studies and draw more reliable conclusions. In addition, the age, 785 medication use, and ethnicity of the patient population varied across studies. Since cognition is heavily 786 influenced by age [173], and medication use [174, 175] and genetics [61] strongly affect the response 787 to exercise, comparability of the included studies may have been limited by these inconsistencies. 788 Regarding the narrative part of this review, an important limitation is that only studies directly 789 demonstrating an involvement of certain mediators in the effect of exercise training on the brain in 790 T2DM have been considered. This means the present review is a non-exhaustive presentation of the 791 involved mediators and pathways. Finally, publication bias might have skewed the results of this review 792 towards a positive effect of exercise on cognition in T2DM.

793 In order to define an exercise program that is specifically tailored to cognitive improvement in T2DM, 794 future studies should directly compare exercise of different intensities and volumes, as well as 795 different exercise modalities. In addition, as already mentioned, a standardised cognitive test should 796 be used that is most sensitive to picking up exercise-induced cognitive changes in T2DM. Furthermore, 797 medication use, disease stage, age, and genetics should be taken into account.

798 5. Conclusion

799 Overall, it can be concluded that exercise has a positive influence on cognition and brain structure in 800 T2DM. There are few studies that fail to find a positive correlation between exercise training and 801 cognition in T2DM, which can often be attributed to a small sample size or exercise of too low intensity. 802 From this review, we can assume that resistance and endurance exercise training have similar effects 803 on cognition in T2DM, and that unconventional exercise training involving the mind, such as tai chi, is 804 also capable of inducing cognitive improvements. However, since there is a lack of studies directly 805 comparing different exercise intensities, little can be said about which exercise intensity would elicit 806 the best cognitive results in T2DM. Future studies should look into this in order to make exercise 807 programs for cognitive preservation in T2DM patients as targeted and efficient as possible.

- 808 6. Declarations
- 809 Author contributions
- 810 Jitske Vandersmissen conducted the literature search and wrote the manuscript. Dominique Hansen,
- 811 Koen Cuypers, and Ilse Dewachter revised the manuscript and provided feedback.
- 812 **Competing interests**
- 813 The authors confirm that there were no conflicts of interest.
- 814 The authors have no relevant financial or non-financial interests to disclose.
- 815 Funding
- 816 Internal funding of Hasselt University (23OWB01BOF) was received to assist with the preparation of 817 this manuscript.
- 818
- 819 7. References
- 820
- 821 1. Harman, M.F. and M.G. Martin, Epigenetic mechanisms related to cognitive decline during 822 aging. J Neurosci Res, 2020. 98(2): p. 234-246.
- 2. Yang, Y., et al., Cognitive Decline Associated with Aging. Adv Exp Med Biol, 2023. 1419: p. 25-823 824 46.

	porpolally.	
It is made available under a	CC-BY-NC-ND 4.0	International license .

825	3.	Grimm, A. and A. Eckert, <i>Brain aging and neurodegeneration: from a mitochondrial point of</i>
820 827	4.	Todorova, V. and A. Blokland, <i>Mitochondria and Synaptic Plasticity in the Mature and Aging</i>
828		Nervous System. Curr Neuropharmacol, 2017. 15(1): p. 166-173.
829 830	5.	Bergado, J.A. and W. Almaguer, <i>Aging and synaptic plasticity: a review</i> . Neural Plast, 2002. <b>9</b> (4): p. 217-32.
831 832	6.	Harada, C.N., M.C. Natelson Love, and K.L. Triebel, <i>Normal cognitive aging</i> . Clin Geriatr Med, 2013. <b>29</b> (4): p. 737-52.
833	7.	Oschwald, J., et al., Brain structure and cognitive ability in healthy gaing: a review on
834		longitudinal correlated change. Rev Neurosci, 2019. <b>31</b> (1): p. 1-57.
835	8.	Blinkouskaya, Y., et al., Brain aging mechanisms with mechanical manifestations. Mech
836		Ageing Dev, 2021. <b>200</b> : p. 111575.
837	9.	Geula, C., et al., Aging renders the brain vulnerable to amyloid beta-protein neurotoxicity. Nat
838		Med, 1998. <b>4</b> (7): p. 827-31.
839	10.	Dickstein, D.L., et al., <i>Dendritic spine changes associated with normal aging</i> . Neuroscience,
840		2013. <b>251</b> : p. 21-32.
841	11.	Antal, B., et al., Type 2 diabetes mellitus accelerates brain aging and cognitive decline:
842		Complementary findings from UK Biobank and meta-analyses. Elife, 2022. 11.
843	12.	Malik, A., et al., Cognitive Impairment in Type 2 Diabetes Mellitus. Cureus, 2022. 14(2): p.
844		e22193.
845	13.	Roy, S., et al., Cognitive Function and Control of Type 2 Diabetes Mellitus in Young Adults. N
846		Am J Med Sci, 2015. <b>7</b> (5): p. 220-6.
847	14.	Zilliox, L.A., et al., Diabetes and Cognitive Impairment. Curr Diab Rep, 2016. 16(9): p. 87.
848	15.	Biessels, G.J., et al., Dementia and cognitive decline in type 2 diabetes and prediabetic stages:
849		<i>towards targeted interventions.</i> Lancet Diabetes Endocrinol, 2014. <b>2</b> (3): p. 246-55.
850	16.	Chakraborty, A., et al., Age Related Prevalence of Mild Cognitive Impairment in Type 2
851		Diabetes Mellitus Patients in the Indian Population and Association of Serum Lipids With
852		Cognitive Dysfunction. Front Endocrinol (Lausanne), 2021. 12: p. 798652.
853	17.	Cheng, G., et al., Diabetes as a risk factor for dementia and mild cognitive impairment: a
854		meta-analysis of longitudinal studies. Intern Med J, 2012. <b>42</b> (5): p. 484-91.
855	18.	Sato, N. and R. Morishita, Brain alterations and clinical symptoms of dementia in diabetes:
856		abeta/tau-dependent and independent mechanisms. Front Endocrinol (Lausanne), 2014. 5: p.
857		143.
858	19.	Karvani, M., et al., Neurocognitive impairment in type 2 diabetes mellitus. Hormones
859		(Athens), 2019. <b>18</b> (4): p. 523-534.
860	20.	Li, Z.Y., et al., Changes of brain function in patients with type 2 diabetes mellitus measured by
861		different analysis methods: A new coordinate-based meta-analysis of neuroimaging. Front
862		Neurol, 2022. <b>13</b> : p. 923310.
863	21.	Liu, J., et al., Reduced Gray Matter Volume in Patients with Type 2 Diabetes Mellitus. Front
864		Aging Neurosci, 2017. <b>9</b> : p. 161.
865	22.	Cui, Y., et al., Aberrant functional connectivity of default-mode network in type 2 diabetes
866		patients. Eur Radiol, 2015. 25(11): p. 3238-46.
867	23.	Tumminia, A., et al., Type 2 Diabetes Mellitus and Alzheimer's Disease: Role of Insulin
868		Signalling and Therapeutic Implications. Int J Mol Sci, 2018. <b>19</b> (11).
869	24.	Michailidis, M., et al., Alzheimer's Disease as Type 3 Diabetes: Common Pathophysiological
870		Mechanisms between Alzheimer's Disease and Type 2 Diabetes. Int J Mol Sci, 2022. <b>23</b> (5).
8/1	25.	Steen, E., et al., Impaired insulin and insulin-like growth factor expression and signaling
8/2		mechanisms in Alzheimer's diseaseis this type 3 diabetes? J Alzheimers Dis, 2005. 7(1): p.
8/3	26	
875 875	26.	Femminelia, G.D., et al., Antidiabetic Drugs in Alzheimer's Disease: Mechanisms of Action and Future Perspectives   Diabetes Res. 2017, <b>2017</b> , p. 7420796

876 877	27.	Blazquez, E., et al., Insulin in the brain: its pathophysiological implications for States related with central insulin resistance, type 2 diabetes and Alzheimer's disease. Front Endocrinol
878		(Lausanne), 2014. <b>5</b> : p. 161.
879	28.	Hooper, C., R. Killick, and S. Lovestone, The GSK3 hypothesis of Alzheimer's disease. J
880		Neurochem, 2008. <b>104</b> (6): p. 1433-9.
881	29.	Talbot, K., et al., Demonstrated brain insulin resistance in Alzheimer's disease patients is
882		associated with IGF-1 resistance, IRS-1 dysregulation, and cognitive decline. J Clin Invest,
883		2012. <b>122</b> (4): p. 1316-38.
884	30.	Farris, W., et al., Insulin-degrading enzyme regulates the levels of insulin, amyloid beta-
885		protein, and the beta-amyloid precursor protein intracellular domain in vivo. Proc Natl Acad
886		Sci U S A, 2003. <b>100</b> (7): p. 4162-7.
887	31.	Sousa, L., et al., Insulin-degrading enzyme: an ally against metabolic and neurodegenerative
888		diseases. J Pathol, 2021. 255(4): p. 346-361.
889	32.	Song, D. and D.S.F. Yu, Effects of a moderate-intensity aerobic exercise programme on the
890		cognitive function and quality of life of community-dwelling elderly people with mild cognitive
891		impairment: A randomised controlled trial. Int J Nurs Stud, 2019. 93: p. 97-105.
892	33.	Baker, L.D., et al., Effects of aerobic exercise on mild cognitive impairment: a controlled trial.
893		Arch Neurol, 2010. <b>67</b> (1): p. 71-9.
894	34.	Gomez-Pinilla, F. and C. Hillman, The influence of exercise on cognitive abilities. Compr
895		Physiol, 2013. <b>3</b> (1): p. 403-28.
896	35.	Lu, Y., et al., Treadmill Exercise Exerts Neuroprotection and Regulates Microglial Polarization
897		and Oxidative Stress in a Streptozotocin-Induced Rat Model of Sporadic Alzheimer's Disease. J
898		Alzheimers Dis, 2017. <b>56</b> (4): p. 1469-1484.
899	36.	Wang, R. and R.M.D. Holsinger, Exercise-induced brain-derived neurotrophic factor
900		expression: Therapeutic implications for Alzheimer's dementia. Ageing Res Rev, 2018. 48: p.
901		109-121.
902	37.	Izquierdo, M., et al., Cytokine and hormone responses to resistance training. Eur J Appl
903		Physiol, 2009. <b>107</b> (4): p. 397-409.
904	38.	de Sousa, C.V., et al., The Antioxidant Effect of Exercise: A Systematic Review and Meta-
905		Analysis. Sports Med, 2017. 47(2): p. 277-293.
906	39.	Demnitz, N., et al., Hippocampal maintenance after a 12-month physical activity intervention
907		in older adults: The REACT MRI study. Neuroimage Clin, 2022. 35: p. 102762.
908	40.	ten Brinke, L.F., et al., Aerobic exercise increases hippocampal volume in older women with
909		probable mild cognitive impairment: a 6-month randomised controlled trial. Br J Sports Med,
910		2015. <b>49</b> (4): p. 248-54.
911	41.	Morris, J.K., et al., Aerobic exercise for Alzheimer's disease: A randomized controlled pilot
912		<i>trial.</i> PLoS One, 2017. <b>12</b> (2): p. e0170547.
913	42.	Choi, S.H., et al., Combined adult neurogenesis and BDNF mimic exercise effects on cognition
914		in an Alzheimer's mouse model. Science, 2018. <b>361</b> (6406).
915	43.	Erickson, K.I., et al., Exercise training increases size of hippocampus and improves memory.
916		Proc Natl Acad Sci U S A, 2011. <b>108</b> (7): p. 3017-22.
917	44.	Burns, J.M., et al., Cardiorespiratory fitness and brain atrophy in early Alzheimer disease.
918		Neurology, 2008. <b>71</b> (3): p. 210-6.
919	45.	Lautenschlager, N.T., et al., Effect of physical activity on cognitive function in older adults at
920		risk for Alzheimer disease: a randomized trial. JAMA, 2008. <b>300</b> (9): p. 1027-37.
921	46.	Buchman, A.S., et al., Total daily physical activity and the risk of AD and cognitive decline in
922		older adults. Neurology, 2012. 78(17): p. 1323-9.
923	47.	Xu, L., et al., Brain metabolism in Alzheimer's disease: biological mechanisms of exercise.
924		Transl Neurodegener, 2023. <b>12</b> (1): p. 33.
925	48.	Yang, J., et al., Lactate promotes plasticity gene expression by potentiating NMDA signaling in
926		neurons. Proc Natl Acad Sci U S A, 2014. <b>111</b> (33): p. 12228-33.

	perpetuity.
It is made available under a	CC-BY-NC-ND 4.0 International license

927 928 920	49.	El Hayek, L., et al., Lactate Mediates the Effects of Exercise on Learning and Memory through SIRT1-Dependent Activation of Hippocampal Brain-Derived Neurotrophic Factor (BDNF). J
929 930 931	50.	Brinkmann, C., et al., <i>Effects of Cycling and Exergaming on Neurotrophic Factors in Elderly</i> <i>Type 2 Diabetic Men - A Preliminary Investigation</i> . Exp Clin Endocrinol Diabetes, 2017. <b>125</b> (7):
932 933 934	51.	p. 436-440. Morland, C., et al., <i>Exercise induces cerebral VEGF and angiogenesis via the lactate receptor</i> HCAB1 Nat Commun. 2017. <b>8</b> : p. 15557
935 936	52.	Morland, C., et al., <i>The lactate receptor, G-protein-coupled receptor 81/hydroxycarboxylic acid receptor 1: Expression and action in brain.</i> J Neurosci Res, 2015. <b>93</b> (7): p. 1045-55.
937 938	53.	Lourenco, M.V., et al., <i>Exercise-linked FNDC5/irisin rescues synaptic plasticity and memory defects in Alzheimer's models</i> . Nat Med, 2019. <b>25</b> (1): p. 165-175.
939 940	54.	Gokce, E. and N. Gun, <i>The Relationship Between Exercise, Cathepsin B, and Cognitive Functions: Systematic Review.</i> Percept Mot Skills, 2023. <b>130</b> (4): p. 1366-1385.
941 942	55.	Huang, B., K. Chen, and Y. Li, <i>Aerobic exercise, an effective prevention and treatment for mild cognitive impairment.</i> Front Aging Neurosci, 2023. <b>15</b> : p. 1194559.
943 944	56.	Moon, H.Y., et al., Running-Induced Systemic Cathepsin B Secretion Is Associated with Memory Function. Cell Metab, 2016. <b>24</b> (2): p. 332-40.
945 946 947	57.	Madarshahian, F., M. Hassanabadi, and M. Koshniat Nikoo, <i>Cognitive status and foot self care</i> practice in overweight diabetics, engaged in different levels of physical activity. J Diabetes Metab Disord, 2014. <b>13</b> (1): p. 31.
948 949	58.	Rabinowitz, Y., et al., <i>Physical Activity Is Associated with Slower Cognitive Decline in Older Adults with Type 2 Diabetes</i> . J Prev Alzheimers Dis, 2023. <b>10</b> (3): p. 497-502.
950 951 952	59.	Zabetian-Targhi, F., et al., <i>The Association Between Physical Activity Intensity, Cognition, and Brain Structure in People With Type 2 Diabetes.</i> J Gerontol A Biol Sci Med Sci, 2021. <b>76</b> (11): p. 2047-2053.
953 954	60.	Shih, I.F., et al., <i>Physical activity modifies the influence of apolipoprotein E epsilon4 allele and type 2 diabetes on dementia and cognitive impairment among older Mexican Americans.</i>
955 956 957	61.	Liu, T., et al., The Influence of the BDNF Val66Met Polymorphism on the Association of Regular Physical Activity With Cognition Among Individuals With Diabetes. Biol Res Nurs,
958 959 960 961	62.	Silveira-Rodrigues, J.G., et al., Acute bouts of aerobic and resistance exercise similarly alter inhibitory control and response time while inversely modifying plasma BDNF concentrations in middle-aged and older adults with type 2 diabetes. Exp Brain Res, 2023. <b>241</b> (4): p. 1173-
962 963 964 965	63.	Wang, H., W. Tang, and Y. Zhao, Acute effects of different exercise forms on executive function and the mechanism of cerebral hemodynamics in hospitalized T2DM patients: a within-subject study. Front Public Health, 2023. <b>11</b> : p. 1165892.
966 967 968	64.	Zhao, R.R., A.J. O'Sullivan, and M.A. Fiatarone Singh, <i>Exercise or physical activity and cognitive function in adults with type 2 diabetes, insulin resistance or impaired glucose tolerance: a systematic review.</i> Eur Rev Aging Phys Act, 2018. <b>15</b> : p. 1.
969 970 971	65.	Wang, R., et al., <i>The effect of physical activity interventions on cognition function in patients with diabetes: A systematic review and meta-analysis.</i> Diabetes Metab Res Rev, 2021. <b>37</b> (7): p. e3443.
972 973 974	66.	Cooke, S., et al., <i>Effects of exercise, cognitive, and dual-task interventions on cognition in type 2 diabetes mellitus: A systematic review and meta-analysis.</i> PLoS One, 2020. <b>15</b> (5): p. e0232958.
975 976 977	67.	Cai, Y.H., et al., <i>Effect of Exercise on the Cognitive Function of Older Patients With Type 2 Diabetes Mellitus: A Systematic Review and Meta-Analysis.</i> Front Hum Neurosci, 2022. <b>16</b> : p. 876935.

978 979	68.	Dyer, A.H., et al., Non-pharmacological interventions for cognition in patients with Type 2 diabetes mellitus: a systematic review OIM 2020 <b>113</b> (3): p. 155-161
980	69.	Callisaya, M. and K. Nosaka, Effects of Exercise on Type 2 Diabetes Mellitus-Related Cognitive
981		Impairment and Dementia. J Alzheimers Dis, 2017. <b>59</b> (2): p. 503-513.
982	70.	Yi, S.S., <i>Effects of exercise on brain functions in diabetic animal models</i> . World J Diabetes,
905	71	2015. 0(4). p. 505-97.
984	/1.	Bertram, S., K. Brixius, and C. Brinkmann, <i>Exercise for the diabetic brain. now physical</i>
985		training may neip prevent dementia and Alzneimer's disease in 12Divi patients. Endocrine,
986	70	2016. <b>53</b> (2): p. 350-63. Dedelelii N., et el., Effecte ef Denular Dhusiael Activity en the Constitute Devicements of Ture 2.
987 988	72.	Diabetic Patients: A Systematic Review. Metab Syndr Relat Disord, 2017. <b>15</b> (10): p. 481-493.
989	73.	Maher, C.G., et al., Reliability of the PEDro scale for rating quality of randomized controlled
990		<i>trials.</i> Phys Ther, 2003. <b>83</b> (8): p. 713-21.
991	74.	Macleod, M.R., et al., Pooling of animal experimental data reveals influence of study design
992		and publication bias. Stroke, 2004. <b>35</b> (5): p. 1203-8.
993	75.	Leischik, R., et al., Exercise Improves Cognitive Function-A Randomized Trial on the Effects of
994		Physical Activity on Cognition in Type 2 Diabetes Patients. J Pers Med, 2021. 11(6).
995	76.	Ploydang, T., et al., Nordic Walking in Water on Cerebrovascular Reactivity and Cognitive
996		Function in Elderly Patients with Type 2 Diabetes. Med Sci Sports Exerc, 2023.
997	77.	Baker, L.D., et al., Aerobic exercise improves cognition for older adults with glucose
998		intolerance, a risk factor for Alzheimer's disease. J Alzheimers Dis, 2010. 22(2): p. 569-79.
999	78.	Anderson-Hanley, C., et al., Neuropsychological benefits of stationary bike exercise and a
1000		cybercycle exergame for older adults with diabetes: an exploratory analysis. J Diabetes Sci
1001		Technol, 2012. <b>6</b> (4): p. 849-57.
1002	79.	Wang, Y., et al., Aerobic Training Increases Hippocampal Volume and Protects Cognitive
1003		Function for Type 2 Diabetes Patients with Normal Cognition. Exp Clin Endocrinol Diabetes,
1004		2023.
1005	80.	Shekarchian, M., M. Peeri, and M.A. Azarbayjani, <i>Physical activity in a swimming pool</i>
1006		attenuates memory impairment by reducing glutamate and inflammatory cytokines and
1007		increasing BDNF in the brain of mice with type 2 diabetes. Brain Res Bull, 2023. <b>201</b> : p.
1008		110725.
1009	81.	Parsa, H., et al., Swimming training and Plantago psyllium ameliorate cognitive impairment
1010		and glucose tolerance in streptozotocin-nicotinamide-induced type 2 diabetic rats. J Physiol
1011		Sci, 2021. <b>71</b> (1): p. 37.
1012	82.	Vorhees, C.V. and M.T. Williams, <i>Morris water maze: procedures for assessing spatial and</i>
1013		related forms of learning and memory. Nat Protoc, 2006. 1(2): p. 848-58.
1014	83.	Jesmin, S., et al., Long-term light and moderate exercise intervention similarly prevent both
1015		hippocampal and alycemic dysfunction in presymptomatic type 2 diabetic rats. Am J Physiol
1016		Endocrinol Metab, 2022. <b>322</b> (3): p. E219-E230.
1017	84.	Lang, X., et al., Treadmill exercise mitigates neuroinflammation and increases BDNF via
1018		activation of SIRT1 signaling in a mouse model of T2DM. Brain Res Bull. 2020. <b>165</b> : p. 30-39.
1019	85.	Shima, T., et al., Moderate exercise ameliorates dysregulated hippocampal alycometabolism
1020		and memory function in a rat model of type 2 diabetes. Diabetologia. 2017. <b>60</b> (3): p. 597-606.
1021	86.	Shima, T., et al., Light-intensity exercise improves memory dysfunction with the restoration of
1022		hippocampal MCT2 and miRNAs in type 2 diabetic mice. Metab Brain Dis. 2023. <b>38</b> (1): p. 245-
1023		254.
1024	87.	Nasreddine, Z.S., et al., The Montreal Coanitive Assessment, MoCA: a brief screening tool for
1025		<i>mild coanitive impairment</i> . J Am Geriatr Soc. 2005. <b>53</b> (4): n. 695-9.
1026	88.	Arevalo-Rodriguez, L. et al., Mini-Mental State Examination (MMSE) for the early detection of
1027		dementia in people with mild cognitive impairment (MCI). Cochrane Database Syst Rev. 2021
1028		<b>7</b> (7): p. CD010783.

1029	89.	Zhao, R.R., et al., Effect of High-Intensity Power Training on Cognitive Function in Older Adults
1030		With Type 2 Diabetes: Secondary Outcomes of the GREAT2DO Study. J Gerontol A Biol Sci
1031		Med Sci, 2022. <b>77</b> (10): p. 1975-1985.
1032	90.	Yamamoto, Y., et al., Effects of resistance training using elastic bands on muscle strength with
1033		or without a leucine supplement for 48 weeks in elderly patients with type 2 diabetes. Endocr
1034		J, 2021. <b>68</b> (3): p. 291-298.
1035	91.	Teixeira, R.B., et al., Evaluating the effects of exercise on cognitive function in hypertensive
1036		and diabetic patients using the mental test and training system. World J Biol Psychiatry,
1037		2019. <b>20</b> (3): p. 209-218.
1038	92.	Silveira-Rodrigues, J.G., et al., Combined exercise training improves specific domains of
1039		cognitive functions and metabolic markers in middle-aged and older adults with type 2
1040		<i>diabetes mellitus.</i> Diabetes Res Clin Pract, 2021. <b>173</b> : p. 108700.
1041	93.	Silveira-Rodrigues, J.G., et al., Combined Training Improves Executive Functions Without
1042		Changing Brain-Derived Neurotrophic Factor Levels of Middle-Aged and Older Adults with
1043		Type 2 Diabetes. Exp Clin Endocrinol Diabetes, 2023. <b>131</b> (6): p. 345-353.
1044	94.	Espeland, M.A., et al., Effects of Physical Activity Intervention on Physical and Cognitive
1045		Function in Sedentary Adults With and Without Diabetes. J Gerontol A Biol Sci Med Sci, 2017.
1046		<b>72</b> (6): p. 861-866.
1047	95.	Ghodrati, N., et al., Effect of Combined Exercise Training on Physical and Cognitive Function in
1048		Women With Type 2 Diabetes, Can J Diabetes, 2023, <b>47</b> (2); p. 162-170.
1049	96.	Martinez-Velilla, N., et al., Effects of a Tailored Exercise Intervention in Acutely Hospitalized
1050		Oldest Old Diabetic Adults: An Ancillary Analysis. J Clin Endocrinol Metab. 2021. <b>106</b> (2): p.
1051		e899-e906.
1052	97.	Callisava, M.L., et al., Feasibility of a multi-modal exercise program on cognition in older
1053		adults with Type 2 diabetes - a nilot randomised controlled trial. BMC Geriatr. 2017. <b>17</b> (1): n
1054		237
1055	98.	Ghahfarrokhi, M.M., et al., Feasibility and preliminary efficacy of different intensities of
1056	50.	functional training in elderly type 2 diabetes natients with cognitive impairment: a nilot
1057		randomised controlled trial BMC Geriatr 2024 <b>24</b> (1): p. 71
1058	99	Chen Y et al. Effects of Tai Chi Chuan on Cognitive Function in Adults 60 Years or Older With
1059	55.	Type 2 Diabetes and Mild Cognitive Impairment in China: A Randomized Clinical Trial JAMA
1060		Netw Open 2023 $6(A)$ : n e237004
1061	100	Cai H et al Effect of Low-Intensity Kinect-Based Kaimai-Style Olaona Exercise in Older
1062	100.	Adults With Type 2 Diabetes   Gerontol Nurs 2019 <b>45</b> (2): n 42-52
1063	101	Li X et al. Effects of fitness gigong and tai chi on middle-aged and elderly natients with type
1064	101.	2 diabetes mellitus PLoS One 2020 <b>15</b> (12): n e02/3989
1065	102	Coelbo Junior H L et al Multicomponent Evercise Improves Hemodynamic Parameters and
1066	102.	Mobility, but Not Maximal Walking Speed Transfer Canacity, and Executive Function of Older
1067		Type II Diabetic Patients Riomed Res Int 2018 <b>2018</b> : n 4832851
1068	103	Shellington F.M. et al. Results From a Feasibility Study of Saugre-Stepping Eversise in Older
1060	105.	Adults With Type 2 Dighetes and Self-Reported Cognitive Complaints to Improve Global
1005		Cognitive Europing Con   Disbetes 2018 <b>42</b> (6): p. 602-612 e1
1070	104	Stanciu G.D. et al. Link Retween Dighetes and Alzheimer's Disease due to the Shared
1071	104.	Amulaid Aggregation and Deposition Involving both Neurodegenerative Changes and
1072		Neurovascular Damages 1 Clin Med 2020 <b>9</b> (6)
1073	105	Li L and C Holscher Common nathological processes in Alzheimer disease and tune 2
1074	105.	diabatas: a raview Brain Das Day 2007 EC/21 a 294 402
1075	106	unuscies, a review, bidin res rev, 2007. 30(2). 4. 304-402. He X E et al. Voluntary Evercica Promotos Clymphatic Classence of Amylaid Bata and
1077	100.	ne, n.r., et al., voluntary exercise Promotes Giymphalic Clearance of Amylola Bela ana Reduces the Activation of Astrophyse and Microsofia in Acad Mico. Front Mol Nourses: 2017
1070		<b>10</b> n 144
7019		τυ. μ. 144.

1079 1080	107.	Xu, L., et al., <i>Treadmill exercise promotes E3 ubiquitin ligase to remove amyloid beta and P-tau and improve cognitive ability in APP/PS1 transgenic mice.</i> J Neuroinflammation, 2022.
1081	100	19(1): p. 243.
1082	108.	Marwarna, G. and O. Ghribi, Leptin signaling and Alzheimer's disease. Am J Neurodegener
1083		Dis, 2012. 1(3): p. 245-65.
1084	109.	Marwarha, G., et al., Leptin attenuates BACE1 expression and amyloid-beta genesis via the
1085		activation of SIRT1 signaling pathway. Biochim Biophys Acta, 2014. <b>1842</b> (9): p. 1587-95.
1086	110.	Rezaei, M.H., et al., Leptin Signaling Could Mediate Hippocampal Decumulation of Beta-
1087		Amyloid and Tau Induced by High-Intensity Interval Training in Rats with Type 2 Diabetes. Cell
1088		Mol Neurobiol, 2023.
1089	111.	De Sousa, R.A.L., et al., High-intensity resistance training induces changes in cognitive
1090		function, but not in locomotor activity or anxious behavior in rats induced to type 2 diabetes.
1091		Physiol Behav, 2020. <b>223</b> : p. 112998.
1092	112.	Fang, X., et al., Phosphorylation and inactivation of glycogen synthase kinase 3 by protein
1093		<i>kinase A</i> . Proc Natl Acad Sci U S A, 2000. <b>97</b> (22): p. 11960-5.
1094	113.	Hernandez, F., J.J. Lucas, and J. Avila, GSK3 and tau: two convergence points in Alzheimer's
1095		disease. J Alzheimers Dis, 2013. <b>33 Suppl 1</b> : p. S141-4.
1096	114.	Rolev, K.D., X.S. Shu, and Y. Ying, <i>Targeted pharmacotherapy against neurodegeneration and</i>
1097		neuroinflammation in early diabetic retinopathy. Neuropharmacology, 2021. <b>187</b> : p. 108498.
1098	115.	Pousti, F., et al., Adiponectin modulates synaptic plasticity in hippocampal dentate gyrus.
1099		Neurosci Lett. 2018. <b>662</b> : p. 227-232.
1100	116.	Weisz, F., et al., The role of adiponectin receptors in the regulation of synaptic transmission in
1101		the hippocampus. Synapse. 2017. <b>71</b> (5).
1102	117.	Ng. R.C., et al., Chronic adiponectin deficiency leads to Alzheimer's disease-like coanitive
1103		impairments and patholoaies through AMPK inactivation and cerebral insulin resistance in
1104		aged mice. Mol Neurodegener. 2016. <b>11</b> (1): p. 71.
1105	118.	lian, M., et al., Adiponectin suppresses amyloid-beta oligomer (AbetaO)-induced
1106		inflammatory response of microalia via AdinoR1-AMPK-NF-kannaB signaling nathway. I
1107		Neuroinflammation, 2019, <b>16</b> (1): p. 110.
1108	119	Mehta BK KK Singh and S Baneriee Effect of exercise on type 2 dighetes-associated
1109	115.	cognitive impairment in rats. Int I Neurosci 2019 <b>129</b> (3): p. 252-263
1110	120	Colucci-D'Amato I. I. Speranza and E. Volnicelli. Neurotrophic Eactor BDNE. Physiological
1111	120.	Functions and Theraneutic Potential in Depression Neurodegeneration and Brain Cancer Int L
1117		Mol Sci 2020 <b>21</b> (20)
1112	121	Jamali A. S. Shahrhanian and S. Morteza Tayehi. <i>The Effects of Exercise Training on the</i>
1111	121.	Brain-Derived Neurotrophic Eactor (BDNE) in the Patients with Type 2 Diabetes: A Systematic
1115		Review of the Randomized Controlled Trials   Diabetes Metab Disord 2020 19(1): n 633-
1116		642
1117	100	045. Chang S.M. and S.D. Loo, Exercise Training Enhances PDNE/TrkP Signaling Bathway and
1110	122.	Inhibits Anontosis in Diabatic Carabral Cartay. Int I Mal Sci. 2022 <b>22</b> (12)
1110	172	Humana, LK, et al. Effects of ano and trondmill everyise in chronic diabatic stages on
1120	125.	nwallg, I.K., et al., Ejjects of uge und tredumin exercise in chronic didbetic stuges on
1120	174	Neurobiast all perentiation in a rat model of type 2 alabetes. Brain Res, 2010. <b>1341</b> . p. 63-71.
1121	124.	ii, S.S., et al., Ejjects of the dentate gurus in a rat model of type II diabetes. Neurosham Pos
1122		subgranular zone of the dentate gyras in a rat model of type in alabetes. Neurochem Res,
1123	105	2009. <b>34</b> (6): p. 1039-46.
1124	125.	Strananan, A.M., et al., <i>Voluntary exercise and caloric restriction ennance nippocampal</i>
1125		<i>denartic spine density and BDNF levels in diabetic mice</i> . Hippocampus, 2009. <b>19</b> (10): p. 951-
1120	120	01.
112/	126.	Li, J., et al., iviecnanisms of Aerobic Exercise Opregulating the Expression of Hippocampal
1128		Synaptic Plasticity-Associated Proteins in Diabetic Rats. Neural Plast, 2019. <b>2019</b> : p. 7920540.

1129 1130 1131	127.	Lee, C.C., C.C. Huang, and K.S. Hsu, <i>Insulin promotes dendritic spine and synapse formation by the PI3K/Akt/mTOR and Rac1 signaling pathways.</i> Neuropharmacology, 2011. <b>61</b> (4): p. 867-79
1132 1133 1134	128.	Barone, E., et al., <i>The interplay among oxidative stress, brain insulin resistance and AMPK dysfunction contribute to neurodegeneration in type 2 diabetes and Alzheimer disease.</i> Free Badic Biol Med. 2021. <b>176</b> : p. 16-33
1135	129.	Xu, J., et al., Brain SIRT1 Mediates Metabolic Homeostasis and Neuroprotection. Front
1137 1138 1139 1140	130.	Banerjee, J., A. Bruckbauer, and M.B. Zemel, Activation of the AMPK/Sirt1 pathway by a leucine-metformin combination increases insulin sensitivity in skeletal muscle, and stimulates glucose and lipid metabolism and increases life span in Caenorhabditis elegans. Metabolism, 2016. <b>65</b> (11): p. 1679-1691.
1141 1142	131.	Wang, R., et al., <i>Deciphering therapeutic options for neurodegenerative diseases: insights from SIRT1</i> . J Mol Med (Berl), 2022. <b>100</b> (4): p. 537-553.
1143 1144 1145	132.	Lin, L., et al., Aerobic Exercise Improves Type 2 Diabetes Mellitus-Related Cognitive Impairment by Inhibiting JAK2/STAT3 and Enhancing AMPK/SIRT1 Pathways in Mice. Dis Markers, 2022. <b>2022</b> : p. 6010504.
1146 1147 1148	133.	Yang, X., et al., <i>The diabetes medication canagliflozin promotes mitochondrial remodelling of adipocyte via the AMPK-Sirt1-Pgc-1alpha signalling pathway.</i> Adipocyte, 2020. <b>9</b> (1): p. 484-494.
1149 1150 1151	134.	Wang, Q., et al., <i>Aerobic Exercise Improves Synaptic-Related Proteins of Diabetic Rats by</i> Inhibiting FOXO1/NF-kappaB/NLRP3 Inflammatory Signaling Pathway and Ameliorating PI3K/Akt Insulin Signaling Pathway. J Mol Neurosci, 2019. <b>69</b> (1): p. 28-38.
1152 1153	135.	Hemmings, B.A. and D.F. Restuccia, <i>PI3K-PKB/Akt pathway</i> . Cold Spring Harb Perspect Biol, 2012. <b>4</b> (9): p. a011189.
1154 1155	136.	Greene, W.C. and L.F. Chen, <i>Regulation of NF-kappaB action by reversible acetylation</i> . Novartis Found Symp, 2004. <b>259</b> : p. 208-17; discussion 218-25.
1156 1157 1158	137.	Zhao, F., M. Tomita, and A. Dutta, <i>Operational Modal Analysis of Near-Infrared Spectroscopy</i> <i>Measure of 2-Month Exercise Intervention Effects in Sedentary Older Adults with Diabetes</i> <i>and Cognitive Impairment</i> . Brain Sci, 2023. <b>13</b> (7).
1159 1160	138.	van Praag, H., et al., <i>Running enhances neurogenesis, learning, and long-term potentiation in mice.</i> Proc Natl Acad Sci U S A, 1999. <b>96</b> (23): p. 13427-31.
1161 1162	139.	Matsui, T., et al., <i>Brain glycogen supercompensation following exhaustive exercise</i> . J Physiol, 2012. <b>590</b> (3): p. 607-16.
1163 1164 1165	140.	Soya, M., et al., <i>Dysregulation of Glycogen Metabolism with Concomitant Spatial Memory</i> <i>Dysfunction in Type 2 Diabetes: Potential Beneficial Effects of Chronic Exercise</i> . Adv Neurobiol, 2019. <b>23</b> : p. 363-383.
1166 1167	141.	Karstoft, K. and B.K. Pedersen, <i>Exercise and type 2 diabetes: focus on metabolism and inflammation.</i> Immunol Cell Biol, 2016. <b>94</b> (2): p. 146-50.
1168 1169 1170	142.	Dutta, B.J., et al., <i>Inside the diabetic brain: Insulin resistance and molecular mechanism associated with cognitive impairment and its possible therapeutic strategies.</i> Pharmacol Res, 2022. <b>182</b> : p. 106358.
1171 1172	143.	Arnold, S.E., et al., <i>Brain insulin resistance in type 2 diabetes and Alzheimer disease: concepts and conundrums.</i> Nat Rev Neurol, 2018. <b>14</b> (3): p. 168-181.
1173 1174	144.	Zhao, H., et al., <i>Insulin Resistance Is a Risk Factor for Mild Cognitive Impairment in Elderly Adults with T2DM</i> . Open Life Sci, 2019. <b>14</b> : p. 255-261.
1175 1176	145.	Kullmann, S., et al., <i>Exercise restores brain insulin sensitivity in sedentary adults who are overweight and obese</i> . JCI Insight, 2022. <b>7</b> (18).
1177 1178	146.	Malin, S.K., et al., <i>Brain insulin resistance and cognitive function: influence of exercise</i> . J Appl Physiol (1985), 2022. <b>133</b> (6): p. 1368-1380.
1179 1180	147.	Li, H. and X. Yu, <i>Emerging role of JNK in insulin resistance</i> . Curr Diabetes Rev, 2013. <b>9</b> (5): p. 422-8.

1181 1182	148.	Sykiotis, G.P. and A.G. Papavassiliou, <i>Serine phosphorylation of insulin receptor substrate-1: a</i>
1102	140	Tapti 1 E ot al. Alteration in inculin action: role of IPS 1 sering phosphorylation in the
1103	149.	ratic, J.F., et al., Alteration in insulin action. Tole of INS-1 Serine phosphorylation in the
1104 110E	150	Perioregulation of insulin signalling. I Coll Sci. 2001. <b>114</b> (Dt 9): p. 1420.20
1185	150.	Bevan, P., <i>Insulin signalling</i> . J Cell Sci, 2001. <b>114</b> (Pt 8): p. 1429-30.
1180	151.	Park, S.H., et al., Resistance Exercise Training Attenuates the Loss of Endogenous GLP-1
118/		Receptor in the Hypothalamus of Type 2 Diabetic Rats. Int J Environ Res Public Health, 2019.
1188		<b>16</b> (5).
1189 1190	152.	Enns, L.C. and W. Ladiges, <i>Protein kinase A signaling as an anti-aging target.</i> Ageing Res Rev, 2010. <b>9</b> (3): p. 269-72.
1191	153.	Chen, C.J., et al., <i>Protein Kingse A and Lipid Metabolism</i> , Sheng Li Ke Xue Jin Zhan, 2016.
1192		<b>47</b> (5): p. 321-9.
1193	154	Saian, M.P. and R.V. Farese. Insulin signalling in henatocytes of humans with type 2 diabetes:
1194		excessive production and activity of protein kinase (-iota (PKC-iota) and dependent processes
1195		and reversal by PKC-iota inhibitors. Diabetologia 2012 <b>55</b> (5): p. 1446-57
1196	155	Sajan M.P. et al. Correction of metabolic abnormalities in a rodent model of obesity
1107	155.	metabolic syndrome, and type 2 diabetes mellitus by inhibitors of benatic protein kinase C-
1100		inter Motabolism 2012 <b>61</b> (4): n 450.60
1100	156	Noior LL CLD 1 recentor agonists for individualized treatment of tune 2 diabetes mellitus
1200	150.	Net Poy Endocrinol 2012 9(12): p 728.42
1200	157	Nat Rev Endocrinioi, 2012. <b>o</b> (12). p. 720-42.
1201	157.	Chefig, D., et al., The Role of Glucugon-Like Peptide-1 Receptor Agonists (GLP-1 RA) In
1202	150	Via We at al. Protain kingan C and protain kingan A are involved in the protaction of
1203	158.	Yin, w., et al., Protein kindse C and protein kindse A dre involved in the protection of
1204		recombinant numan giucagon-like peptide-1 on giomeruli and tubules in diabetic rats. J
1205	450	Diabetes Investig, 2019. <b>10</b> (3): p. 613-625.
1206	159.	Rovira-Llopis, S., et al., Mitochondrial dynamics in type 2 diabetes: Pathophysiological
1207	1.00	
1208	160.	Wada, J. and A. Nakatsuka, Mitochondrial Dynamics and Mitochondrial Dysfunction in
1209		<i>Diabetes.</i> Acta Med Okayama, 2016. <b>70</b> (3): p. 151-8.
1210	161.	Luo, J.S., et al., The Role of Mitochondrial Quality Control in Cognitive Dysfunction in
1211		<i>Diabetes.</i> Neurochem Res, 2022. <b>47</b> (8): p. 2158-2172.
1212	162.	Cheng, H., et al., Mitochondrial dysfunction plays a key role in the development of
1213		<i>neurodegenerative diseases in diabetes.</i> Am J Physiol Endocrinol Metab, 2020. <b>318</b> (5): p.
1214		E750-E764.
1215	163.	Gusdon, A.M., et al., Exercise increases mitochondrial complex I activity and DRP1 expression
1216		in the brains of aged mice. Exp Gerontol, 2017. <b>90</b> : p. 1-13.
1217	164.	Amar, D., et al., The mitochondrial multi-omic response to exercise training across tissues.
1218		bioRxiv, 2023.
1219	165.	Mendham, A.E., et al., Exercise training improves mitochondrial respiration and is associated
1220		with an altered intramuscular phospholipid signature in women with obesity. Diabetologia,
1221		2021. <b>64</b> (7): p. 1642-1659.
1222	166.	Meex, R.C., et al., Restoration of muscle mitochondrial function and metabolic flexibility in
1223		type 2 diabetes by exercise training is paralleled by increased myocellular fat storage and
1224		improved insulin sensitivity. Diabetes, 2010. 59(3): p. 572-9.
1225	167.	Hey-Mogensen, M., et al., Effect of physical training on mitochondrial respiration and
1226		reactive oxygen species release in skeletal muscle in patients with obesity and type 2
1227		<i>diabetes.</i> Diabetologia, 2010. <b>53</b> (9): p. 1976-85.
1228	168.	Toledo, F.G., et al., Effects of physical activity and weight loss on skeletal muscle
1229		mitochondria and relationship with glucose control in type 2 diabetes. Diabetes. 2007. 56(8):
1230		p. 2142-7.

1231	169.	Baekkerud, F.H., et al., High Intensity Interval Training Ameliorates Mitochondrial Dysfunction
1232		in the Left Ventricle of Mice with Type 2 Diabetes. Cardiovasc Toxicol, 2019. 19(5): p. 422-
1233		431.
1234	170.	Ruegsegger, G.N., et al., Exercise and metformin counteract altered mitochondrial function in
1235		the insulin-resistant brain. JCI Insight, 2019. <b>4</b> (18).
1236	171.	American College of Sports, M., et al., American College of Sports Medicine position stand.
1237		<i>Exercise and physical activity for older adults.</i> Med Sci Sports Exerc, 2009. <b>41</b> (7): p. 1510-30.
1238	172.	Cassilhas, R.C., et al., The impact of resistance exercise on the cognitive function of the
1239		<i>elderly.</i> Med Sci Sports Exerc, 2007. <b>39</b> (8): p. 1401-7.
1240	173.	Murman, D.L., The Impact of Age on Cognition. Semin Hear, 2015. 36(3): p. 111-21.
1241	174.	Peel, C. and K.A. Mossberg, Effects of cardiovascular medications on exercise responses. Phys
1242		Ther, 1995. <b>75</b> (5): p. 387-96.
1243	175.	Powles, A.C., The effect of drugs on the cardiovascular response to exercise. Med Sci Sports
1244		Exerc, 1981. <b>13</b> (4): p. 252-8.