

**Supplemental Table 1. Model Estimates****Model for post-treatment MADRS scores (placebo arm)**

	<b>Estimate (SE)</b>	<b>95% CI</b>	<b>t-value</b>	<b>p-value</b>
Intercept	-2.86 (10.1)	-22.7 to 17.0	-0.28	0.777
Group (baseline opioid use status)	10.1 (4.73)	0.81 to 19.4	2.13	0.033*
Time	-0.07 (0.17)	-0.40 to 0.27	-0.39	0.699
Group×Time	0.24 (0.29)	-0.32 to 0.79	0.83	0.409
Baseline MADRS score	0.70 (0.31)	0.09 to 1.31	2.26	0.024*
Baseline pain intensity	-1.34 (0.74)	-2.78 to 0.11	-1.81	0.070
Postoperative pain intensity	0.50 (0.42)	-0.33 to 1.33	1.18	0.239
Ethnicity	-9.21 (9.22)	-27.3 to 8.87	-1.00	0.318

**Model for % change in MADRS scores (placebo arm)**

	<b>Estimate (SE)</b>	<b>95% CI</b>	<b>t-value</b>	<b>p-value</b>
Intercept	-43.8 (12.7)	-68.7 to -19.0	-3.46	0.001*
Group (baseline opioid use status)	38.4 (15.2)	8.59 to 68.2	2.53	0.012*
Time	-0.15 (0.56)	-1.25 to 0.95	-0.27	0.789
Group×Time	0.54 (0.95)	-1.33 to 2.41	0.57	0.570
Baseline pain intensity	-3.81 (2.40)	-8.51 to 0.89	-1.59	0.112
Postoperative pain intensity	1.75 (1.42)	-1.02 to 4.52	1.24	0.216
Ethnicity	-34.4 (30.0)	-93.1 to 24.3	-1.15	0.251

**Model for post-treatment MADRS scores (ketamine arm)**

	<b>Estimate (SE)</b>	<b>95% CI</b>	<b>t-value</b>	<b>p-value</b>
Intercept	-2.30 (7.15)	-16.3 to 11.7	-0.32	0.748
Group (baseline opioid use status)	2.63 (3.87)	-4.96 to 10.2	0.68	0.497
Time	0.31 (0.20)	-0.09 to 0.70	1.52	0.127
Group×Time	-0.11 (0.28)	-0.67 to 0.44	-0.40	0.686
Baseline MADRS score	0.50 (0.23)	0.05 to 0.95	2.19	0.029*
Baseline pain intensity	-0.87 (1.22)	-3.26 to 1.52	-0.72	0.474
Postoperative pain intensity	1.11 (0.63)	-0.13 to 2.34	1.75	0.079
Ethnicity	4.99 (6.14)	-7.03 to 17.0	0.81	0.416

**Model for % change in MADRS scores (ketamine arm)**

	<b>Estimate (SE)</b>	<b>95% CI</b>	<b>t-value</b>	<b>p-value</b>
Intercept	-57.9 (21.5)	-100 to -15.7	-2.69	0.007*
Group (baseline opioid use status)	2.92 (14.5)	-25.5 to 31.3	0.20	0.840
Time	0.47 (0.84)	-1.18 to 2.11	0.55	0.579
Group×Time	0.29 (1.18)	-2.02 to 2.60	0.25	0.806
Baseline pain intensity	-2.66 (4.25)	-11.0 to 5.68	-0.62	0.532
Postoperative pain intensity	4.43 (2.58)	-0.63 to 9.49	1.72	0.086
Ethnicity	19.2 (22.5)	-24.9 to 63.4	0.85	0.393

**Model for post-treatment HADS scores (placebo arm)**

	<b>Estimate (SE)</b>	<b>95% CI</b>	<b>t-value</b>	<b>p-value</b>
Intercept	14.3 (6.93)	0.71 to 27.9	2.06	0.039*
Group (baseline opioid use status)	2.41 (3.60)	-4.64 to 9.46	0.67	0.502
Time	-0.36 (0.10)	-0.55 to -0.16	-3.61	<0.001*
Group×Time	0.41 (0.17)	0.08 to 0.74	2.45	0.014*

Baseline HADS	0.24 (0.27)	-0.29 to 0.76	0.88	0.379
Baseline pain intensity	-0.36 (0.58)	-1.50 to 0.78	-0.62	0.533
Postoperative pain intensity	0.47 (0.25)	-0.02 to 0.96	1.90	0.058
Ethnicity	-3.23 (7.06)	-17.1 to 10.6	-0.46	0.648
<b>Model for % change in HADS scores (placebo arm)</b>				
	<b>Estimate (SE)</b>	<b>95% CI</b>	<b>t-value</b>	<b>p-value</b>
Intercept	36.9 (25.2)	-12.6 to 86.3	1.46	0.144
Group (baseline opioid use status)	16.1 (13.2)	-9.80 to 42.1	1.22	0.223
Time	-1.50 (0.45)	-2.38 to -0.63	-3.36	0.001*
Group×Time	1.52 (0.76)	0.04 to 3.01	2.01	0.045*
Baseline pain intensity	-1.11 (2.13)	-5.28 to 3.06	-0.52	0.601
Postoperative pain intensity	2.15 (1.12)	-0.05 to 4.34	1.92	0.055
Ethnicity	-21.9 (25.7)	-72.1 to 28.4	-0.85	0.394

\*Statistical significance defined by  $p < 0.05$  without corrections. SE, standard error. CI, confidence interval. MADRS, Montgomery-Asberg Depression Rating Scale. HADS, Hospital Anxiety and Depression Scale. Pain intensity was rated on a 0-10 numeric scale. Baseline values were obtained during screening. Post-treatment values were obtained on postoperative days 1, 2, 3, 5, 7 and 14.

### **Caption for Supplemental Table 1. Model Estimates**

Model estimates for post-treatment MADRS and HADS scores and percent change in scores for placebo and ketamine arms, detailing the effects of baseline opioid use status, time, and additional predictors on the outcomes.